JEFFERSON MATH PROJECT REGENTS BY CHAPTER

794 NY Math Regents Exam Questions Sorted by Amsco Chapter INTEGRATED ALGEBRA (Answer Key)

www.jmap.org

Dear Sir

I have to acknolege the reciept of your favor of May 14. in which you mention that you have finished the 6. first books of Euclid, plane trigonometry, surveying & algebra and ask whether I think a further pursuit of that branch of science would be useful to you. there are some propositions in the latter books of Euclid, & some of Archimedes, which are useful, & I have no doubt you have been made acquainted with them. trigonometry, so far as this, is most valuable to every man, there is scarcely a day in which he will not resort to it for some of the purposes of common life. the science of calculation also is indispensible as far as the extraction of the square & cube roots; Algebra as far as the quadratic equation & the use of logarithms are often of value in ordinary cases: but all beyond these is but a luxury; a delicious luxury indeed; but not to be indulged in by one who is to have a profession to follow for his subsistence. in this light I view the conic sections, curves of the higher orders, perhaps even spherical trigonometry, Algebraical operations beyond the 2d dimension, and fluxions.

Letter from Thomas Jefferson to William G. Munford, Monticello, June 18, 1799.

	nap.org
[1]	<u>D</u>
[2]	<u>D</u>
[3]	<u>C</u>
[4]	<u>B</u>
[5]	<u>A</u>
[6]	<u>C</u>
[7]	<u>B</u>
[8]	<u>A</u>
[9]	[2] No, and a correct justification is given. [1] No, but an incomplete or partially incorrect explanation is given. [0] No, but no explanation is given. or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.
[10]	<u>A</u>
[11]	<u>D</u>
	[2] An irrational number is written, and an appropriate explanation is written, such as an irrational number cannot be written as a fraction or as a repeating or terminating decimal. [1] An irrational number is written, such as π or the square root of a nonperfect square, but no explanation or an inappropriate explanation is written. or [1] A correct definition of an irrational number is written, but the example is missing or is inappropriate.
	[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
[12]	incorrect procedure.
[13]	<u>A</u>
[14]	В

[15] <u>A</u>

[16]	<u>B</u>
[17]	<u>B</u>
[18]	<u>A</u>
[19]	 [2] √196, and an appropriate explanation is given. [1] An incorrect answer is chosen, but an appropriate explanation is given. or [1] √196, but no explanation or an incorrect explanation is given. [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.
[20]	<u>A</u>
	[2] $2\frac{4}{5}$, $\sqrt{8}$, $3.\overline{1}$, π , $2\sqrt{3}$ and appropriate work is shown, such as converting each value to a decimal equivalent. [1] All values are correctly converted to decimal equivalents, but the order is not indicated or is indicated incorrectly. or [1] One or two computational errors are made in finding decimal equivalents, but the appropriate order is indicated. or [1] Appropriate work is shown, but one conceptual error is made, such as indicating the order from greatest to least. or [1] $2\frac{4}{5}$, $\sqrt{8}$, $3.\overline{1}$, π , $2\sqrt{3}$, but no work is shown. [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct
[21]	response that was obtained by an obviously incorrect procedure.
[22]	A
[23]	<u>A</u>
	<u>A</u>

[25] <u>A</u>

[26] <u>A</u>

- [27] D
- [28] B
- [29] B
- [30] A
- [31] D
- [32] <u>C</u>
- [33] B
- [34] B

[1]	<u>B</u>	[15] <u>B</u>
[2]	<u>B</u>	[16] <u>A</u>
[3]	<u>D</u>	[17] <u>A</u>
[4]	<u>D</u>	[18] <u>A</u>
	[3] All three examples are illustrated under	[19] <u>A</u>
	division correctly, such as $2 \div 0$, $-2 \div 4$, $-2 \div -4$, and correct explanations are given.	[20] <u>C</u>
	[2] Only two of the three examples are	[21] <u>C</u>
	illustrated and explained correctly. or [2] All three examples are illustrated	[22] <u>D</u>
	correctly, but only one explanation is given or is correct.	[23] A
	or [2] The division examples and explanations are correct, but at most two incorrect	[24] B
	examples are also shown, such as examples	[25] A
	for addition, subtraction, or multiplication. [1] The division examples and explanations	[26] B
	are correct, but more than two incorrect examples are shown, such as examples for	[27] C
	addition, subtraction, or multiplication.	[28] B
	or [1] All three examples are illustrated correctly, but no correct explanation is given.	[29] D
[5]	or [1] Only one correct example with a correct explanation is given. [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.	[2] 0, and an appropriate explanation is given, such as 0 is the number that when added to any number results in that number or does not change it, or $1 + 0 = 1$, $2 + 0 = 2$, and $3 + 0 = 3$.
[6]	C	[1] 0, but no explanation or an incorrect explanation is given.
[7]	C	[0] A zero response is completely incorrect,
[8]	C	irrelevant, or incoherent or is a correct response that was obtained by an obviously
[9]	<u> </u>	[30] incorrect procedure.
[10]	 D	[31] <u>B</u>
[11]	 D	[32] <u>B</u>
[12]	 D	[33] <u>C</u>
[13]	<u> </u>	[34] <u>B</u>
[14]		
[- 1]	-	

- [2] 1, and an appropriate explanation is given, such as when 1 is added to 3, the result is the identity element, 4; therefore 1 is the inverse of 3.
- [1] 1 + 3 = 4, but the identity element is not identified.
- or [1] 4 is identified as the inverse because the identity element and inverse element are confused.
- or [1] 1, but no explanation or an incorrect explanation is given.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [35] incorrect procedure.
- [36] C
- [37] A

[1]	<u>D</u>
[2]	<u>D</u>
[3]	<u>A</u>
[4]	<u>C</u>
[5]	<u>A</u>
[6]	<u>B</u>
[7]	[2] No, and an appropriate explanation is given or the expression is evaluated correctly. [1] No, and the correct order of operations is used to evaluate $2(3)^2 + 5$, but one computational error is made. or [1] One conceptual error is made in evaluating the expression, but the question is answered appropriately. or [1] Appropriate work is shown, but the question is not answered. [0] No, but no explanation or an inappropriate explanation is given. or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.
[8]	<u>B</u>
[9]	<u>B</u>
[10]	<u>B</u>
[11]	<u>D</u>
[12]	<u>B</u>

ww.ji	nap.org		
[1]	<u>D</u>		[2] 21 and the student shows an appropriate
[2]	<u>D</u>		solution, such as the equation $x + x + 1 + x + 2 = 63$ or trial and error.
[3]	<u>A</u>		[1] Appropriate work is shown, but an incorrect answer is found.
[4]	<u>B</u>		or [1] An incorrect equation is shown, but it is solved appropriately to find an answer, such
	 [2] 6, and appropriate work is shown, such as solving the equation 2x + 3 = 15 or trial and error with at least three trials and appropriate checks. [1] Appropriate work is shown, but one 		as $x + x + 2 + x + 4 = 63$. or [1] 21 but no work is shown. [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
	computational error is made. or [1] Appropriate work is shown, but one	[10]	incorrect procedure.
	conceptual error is made.	[11]	<u>D</u>
	or [1] A correct equation is written, but no further correct work is shown.	[12]	<u>B</u>
[5]	or [1] The trial-and-error method is attempted and at least six systematic trials and appropriate checks are shown, but no solution is found. or [1] 6, but no work or fewer than three trials and appropriate checks are shown. [0] A zero response is completely incorrect, irrelevant or incoherent or is a correct response that was obtained by an obviously	[13]	[2] 4, and appropriate work is shown. [1] Appropriate work is shown, but one computational error is made. or [1] Appropriate work is shown, but one conceptual error is made. or [1] 4, but no work is shown. [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.
[6]	C	[14]	В
[7]	C		
[8]	<u> </u>	[15]	
[o]		[16]	<u>B</u>
	[4] \$167.50, and appropriate work is shown, such as $350x + (150)(130) = 1.25(62,500)$ or trial and error with at least three trials with appropriate checks. [3] Appropriate work is shown, but one computational error is made. [2] Appropriate work is shown, but more than one computational error is made. or [2] \$167.50, but only one trial with an appropriate check is shown. [1] \$167.50, but no work is shown. [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously	[17]	<u>D</u>
[9]	incorrect procedure.		

- [2] 10, and appropriate work is shown, such as solving the equation or trial and error with at least three trials and appropriate checks.
- [1] Appropriate work is shown, but one computational error is made.
- or [1] Appropriate work is shown, but one conceptual error is made.
- or [1] The trial-and-error method is attempted and at least six systematic trials and appropriate checks are shown, but no solution is found.
- or [1] 10, but no work or fewer than three trials and appropriate checks are shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [18] incorrect procedure.
- [19] B
- [20] C
- [21] D
 - [2] 18 and correct substitution, $F = \frac{9}{5}(-8) +$
 - 32, is shown.
 - [1] A correct substitution method is shown, but one computational error is made.
 - or [1] The answer is not rounded to the nearest integer, such as 17.6 or 17.
 - or [1] The student substitutes -8 for F, but then solves appropriately for C.
 - or [1] The student substitutes +8 for C, but then solves appropriately for F.
 - or [1] 18 but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [22] incorrect procedure.

- [2] 50, and appropriate work is shown, such as solving the equation $10 = \frac{5}{9}(F 32)$.
- [1] Appropriate work is shown, but one computational error is made.
- or [1] Appropriate work is shown, but one conceptual error is made.
- or [1] Correct substitution is made into the equation, but no further correct work is shown.
- or [1] 50, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [23] incorrect procedure.
- [24] D
 - [3] 3 hours and an appropriate method or equation is shown, such as 45(x + 1) = 60x. [2] An appropriate method is shown, but an
 - incorrect answer is found, such as 4 hours (the truck's time) or 180 miles traveled.
 - [1] An appropriate equation or method is shown, but no answer is found, such as showing an equation that reflects a one-hour difference in time but it is not solved. or [1] 3 hours and no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [25] incorrect procedure.
- [26] A

- [2] Bob, and appropriate work is shown, such as using the distance formula to calculate the two travel times or setting up a proportion.
- [1] Appropriate work is shown, but one computational or conceptual error is made, but an appropriate answer is found.
- or [1] Appropriate work is shown, but no answer or an incorrect answer is found.
- [0] Bob, but no work or inappropriate work is shown.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an abviously incorrect presendant.
- [27] obviously incorrect procedure.
 - [2] 5, and appropriate work is shown, such as solving the linear equation 80x + 100x = 900, using a diagram or proportion or trial and error.
 - [1] Appropriate work is shown, but one computational error is made.
 - or [1] 5, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.
- [28] incorrect procedure.
 - [2] 20, and appropriate work is shown, such as $\frac{15}{150} = \frac{2}{x}$.
 - [1] Appropriate work is shown, but one computational error is made.
 - or [1] Appropriate work is shown, but one conceptual error is made, such as expressing

the answer as $\frac{1}{3}$ hour.

- or [1] 20, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [29] incorrect procedure.
- [30] D
- [31] B

- [3] 4 and an appropriate method is shown, such as calculating A at 6 mph and B at 2 mph through arithmetic, formula, or extending the graph to 60 minutes.
- [2] The speeds of 6 and 2 are found but not their difference.
- or [2] Their difference is found but not in miles per hour.
- [1] Only distances of 4.5 miles and 1.5 miles are found.
- or [1] The speeds found are incorrect but then are subtracted appropriately.
- or [1] 3 times as fast and no appropriate explanation is given.
- or [1] 4 and no appropriate explanation is given.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously [32] incorrect procedure.
 - [3] 12, and appropriate work is shown, such as finding the rates of both vehicles and then subtracting 48 from 60.
 - [2] Appropriate work is shown, but one computational error is made.
 - or [2] The rates of both vehicles are found correctly, and appropriate work is shown, but they are not subtracted.
 - or [2] The rates of both vehicles are found correctly, and the correct difference is found, but no work is shown.
 - [1] Appropriate work is shown, but two or more computational errors are made.
 - or [1] Appropriate work is shown, but one conceptual error is made.
 - or [1] The rates of both vehicles are found correctly, but no work is shown, and the difference is not found.
 - or [1] 12, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [33] incorrect procedure.

- [3] 50, 1.5, and 10, and appropriate work is shown.
- [2] Appropriate work is shown, but one computational error is made.
- [1] Appropriate work is shown, but one conceptual error is made.
- or [1] Appropriate work is shown, but two or more computational errors are made.
- or [1] 50, and appropriate work is shown, but no further correct work is shown.
- or [1] 1.5, and appropriate work is shown, but no further correct work is shown.
- or [1] 10, and appropriate work is shown, but no further correct work is shown.
- or [1] 50, 1.5, and 10, but no work is shown.
- [0] 50 or 1.5 or 10, but no work is shown.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an
- [34] obviously incorrect procedure.
- [35] C
- [36] B
- [37] C
- [38] B
- [39] B

- [2] 6x-2 or an equivalent expression, and appropriate work is shown, such as 2(2x+3)+2(x-4)=6x-2.
- [1] The length is represented correctly as 2x+3 and the width as x-4, but the representation of the perimeter is determined incorrectly.
- or [1] The length, the width, and the perimeter are represented appropriately, but by a variable other than x.
- or [1] One or both dimensions are represented incorrectly, but the perimeter is represented appropriately.
- [0] One or both dimensions are represented incorrectly, and the perimeter is not determined.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an [40] obviously incorrect procedure.
 - [3] 18, and appropriate work is shown.
 - [2] Appropriate work is shown, but one computational error is made.
 - or [2] Appropriate work is shown, and the value of x is found, but no further correct work is shown.
 - [1] Appropriate work is shown, but two or more computational errors are made.
 - or [1] Appropriate work is shown, but one conceptual error is made.
 - or [1] A correct expression is written for the perimeter of each figure, but no further correct work is shown.
 - or [1] 18, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [41] incorrect procedure.
- [42] B
- [43] C
- [44] C
- [45] D
- [46] <u>C</u>

- [47] A
- [48] D
- [49] D
- [50] C
- [51] <u>A</u>
- [52] C
 - a [1] $\frac{S+24}{3}$ or $\frac{S}{3}+8$
 - b [1] 11.5
 - or [1] Correct substitution into an incorrect part a is shown, and the answer is given to the nearest tenth of an inch.

a and b

- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [53] incorrect procedure.
- [54] A
- [55] B
- [56] B
- [57] D
- [58] C

- [3] -5, -3, -1, and appropriate work is shown, such as solving the inequality or trial and error with at least three trials and appropriate checks.
- [2] Appropriate work is shown, but one computational error is made.
- or [2] Appropriate work is shown, and the

inequality $x \ge -5\frac{1}{3}$ is written, but no further

correct work is shown.

- or [2] The trial-and-error method is used to find the correct solutions, but only two trials and appropriate checks are shown.
- [1] Appropriate work is shown, but two or more computational errors are made.
- or [1] Appropriate work is shown, but one conceptual error is made.
- or [1] The trial-and-error method is attempted and at least six systematic trials and appropriate checks are shown, but the solutions are not found.
- or [1] -5, -3, -1, but no work or only one trial with an appropriate check is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [59] incorrect procedure.
- [60] D
- [61] C
- [62] A
- [63] D

- [2] A correct graph is drawn on the number line, with a closed circle at the left end and an open circle at the right end.
- [1] Appropriate work is shown, but one graphing error is made, such as writing an incorrect scale on the number line.
- or [1] Appropriate work is shown, but one conceptual error is made, such as using a closed circle instead of an open circle.
- or [1] A correct inequality is written, but the graph is not drawn.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously [64] incorrect procedure.
 - [2] 65, and appropriate work is shown, such as solving the inequality $15x \ge 225 + 750$ or trial and error with at least three trials and appropriate checks.
 - [1] Appropriate work is shown, but one computational error is made.
 - or [1] Appropriate work is shown, but one conceptual error is made.
 - or [1] The trial-and-error method is attempted and at least six systematic trials and appropriate checks are shown, but no solution is found.
 - or [1] 65, but no work or fewer than three trials and appropriate checks are shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [65] incorrect procedure.

- [3] 7, and appropriate work is shown, such as solving the inequality $15x + 22 \ge 120$, solving an equation, or trial and error with at least three trials and appropriate checks.
- [2] Appropriate work is shown, but one computational or rounding error is made.
- or [2] The trial-and-error method is used to find a correct solution, but only two trials and appropriate checks are shown.
- [1] Appropriate work is shown, but two or more computational or rounding errors are made.
- or [1] Appropriate work is shown, but one conceptual error is made.
- or [1] An incorrect equation of equal difficulty is solved appropriately.
- or [1] A correct inequality or equation is written, but no further correct work is shown.
- or [1] The trial-and-error method is attempted and at least six systematic trials and appropriate checks are shown, but no solution is found.
- or [1] 7, but no work or only one trial with an appropriate check is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [66] incorrect procedure.
- [67] D
- [68] C
 - [2] 6, and appropriate work is shown, such as $0.70x + 0.30 \le 5.00$ or trial and error with three trials and appropriate checks.
 - [1] The inequality is solved correctly, but the number of doughnuts is not found.
 - or [1] The trial-and-error method is used to find a correct solution, but fewer than three trials are shown.
 - or [1] 6, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [69] incorrect procedure.

- [3] 15 and an appropriate method or explanation is shown, such as trial and error or the inequality $6x + 15 \ge 100$.
- [2] An appropriate method is shown, but it stops at 14.
- [1] An appropriate method is shown, but no answer is found.
- or [1] 15 and no explanation is given.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [70] incorrect procedure.
- [71] B

- [1] A
- [2] A
- [3] D
- [4] A
- [5] B
- [6] B
- [7] C
- [8] D
- [9] A
- [10] D
- [11] D
- [12] A
- [13] <u>B</u>
 - [2] $4x^2 + 10x + 2$, and appropriate work is shown, such as $(9x^2 + 3x 4) (5x^2 7x 6)$.
 - [1] The setup is correct, but the distribution of the negative sign is incorrect.
 - or [1] $14x^2 4x 10$, but appropriate work is shown.
 - or [1] $4x^2 + 10x + 2$, but no work is shown. [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [14] incorrect procedure.
- [15] C
- [16] <u>C</u>
- [17] C
- [18] D
- [19] B
- [20] C
- [21] D
- [22] A

- [23] A
- [24] B
- [25] A
- [26] D
- [27] D
- [28] C
- [29] D
- [30] B
- [31] C
- [32] B
- [33] B
- [34] D
- [35] A
- [36] D
- [37] A
- [38] D
- [39] A
- [40] B
- [41] A
- [42] A
- [43] <u>C</u>
- [44] <u>D</u>
- [45] D
- [46] <u>D</u>
- [47] B
- [48] <u>D</u>
- [49] <u>A</u>
- [50] <u>C</u>

- [51] A
- [52] C
 - [3] 499 days and appropriate work is shown,

such as
$$\frac{17,000,000 \, miles}{1420 \frac{miles}{hour} \times 24 \frac{hours}{day}}.$$

- [2] Appropriate work is shown, but one computational error is made or the student incorrectly calculates 1.7×10^7 by one decimal place.
- or [2] Appropriate work is shown, but the answer is rounded incorrectly or is not rounded.
- [1] $1.7 \times 10^7 = 17,000,000$ is shown.
- or [1] $\frac{1.7 \times 10^7}{1420}$ = 11,971.831 hours is shown.
- or [1] 34,080 miles in 1 day is shown.
- or [1] 499 but no work is shown.
- [0] The student does not understand scientific notation.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an
- [53] obviously incorrect procedure.
- [54] <u>B</u>
- [55] C
- [56] D
- [57] D
- [58] A
- [59] C
- [60] D
- [61] A
- [62] A
- [63] A
- [64] <u>D</u>

vw.jr	nap.org
[1]	<u>B</u>
[2]	<u>B</u>
[3]	<u>A</u>
[4]	<u>C</u>
[5]	[4] \$52,950, \$35,300, and \$88,250 and an appropriate method is shown, such as $3x + 2x + 5x = $176,500$. [3] A correct equation is set up or multiplied by correct fractional values $\frac{3}{10}$, $\frac{2}{10}$, and $\frac{5}{10}$, but a computational mistake is made, and three appropriate values are found. or [3] An appropriate method is shown, but not all three values are found. [2] The equation is set up correctly, but numerous computational mistakes are made, and three appropriate values are found. or [2] An incorrect equation is shown, but three appropriate values are found. or [2] An appropriate equation is shown but is solved only for x (17,650). [1] The equation is set up correctly, but no appropriate values are found. or [1] Three correct answers are found, and no work is shown. [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.
[6]	<u>D</u>
[7]	<u>D</u>

- [4] \$68,000, and appropriate work is shown. [3] \$119,000 and \$51,000, and appropriate work is shown, but the answers are not subtracted to find the difference. or [3] Appropriate work is shown, but one computational error is made. [2] Appropriate work is shown, but more than one computational error is made. [1] The value for one share (\$17,000) is found, but no further correct work is shown. or [1] \$68,000, but no work is shown. [0] \$17,000 or \$119,000 or \$51,000, and no work is shown. or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an
- [9] D___

[8] obviously incorrect procedure.

- [10] <u>B</u>
- [11] B
- [12] <u>D</u>
 - [2] \$178.50, and appropriate work is shown, such as solving a proportion, using a table, or trial and error with at least three trials and appropriate checks.
 - [1] Appropriate work is shown, but one computational error is made.
 - or [1] An appropriate proportion is set up, but no solution or an incorrect solution is found. or [1] An incorrect proportion is set up, but an
 - or [1] An incorrect proportion is set up, but ar appropriate solution is found.
 - or [1] \$178.50, but no work is shown or fewer than three trials with appropriate checks are shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [13] incorrect procedure.

[2] $6\frac{2}{3}$ or 6 hr 40 min or $6.6\overline{6}$ or an

equivalent answer, and appropriate work is shown.

- [1] 400 min, but the answer is not converted into hours.
- or [1] Appropriate work is shown, but one computational error is made.
- or [1] Appropriate work is shown, but the answer is rounded to the nearest hour.

or [1]
$$6\frac{2}{3}$$
 or 6 hr 40 min or $6.6\overline{6}$ or an

equivalent answer, but no work is shown. [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously

- [14] incorrect procedure.
- [15] B
- [16] C
- [17] B
 - [2] 319, and appropriate work is shown.
 - [1] A correct proportion is shown, but no solution or an incorrect solution is found. or [1] An incorrect proportion of equal difficulty is solved appropriately. or [1] Appropriate work is shown, but one computational error is made.
 - or [1] 319, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [18] <u>incorrect procedure.</u>
- [19] B
 - [2] 45, and appropriate work is shown, such as a diagram or $\frac{1.2}{2} = \frac{x}{75}$.
 - [1] Appropriate work is shown, but no answer or an incorrect answer is found.
 - or [1] 45, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect precedure.
- [20] incorrect procedure.

[2] 15, and any equivalent proportion, equation, or fraction conversion is shown,

such as
$$\frac{12}{16} = \frac{x}{20}$$
.

- [1] An appropriate proportion, equation, or fraction conversion is shown, but one computational or conceptual error is made. or [1] An incorrect proportion, equation, or fraction conversion is shown, but an appropriate answer is found for the incorrect proportion.
- or [1] 15, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [21] incorrect procedure.
 - [2] 78.6%, and appropriate work is shown.
 - [1] Appropriate work is shown, but one computational or rounding error is made. or [1] Appropriate work is shown, but one
 - conceptual error is made. or [1] 78.6%, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, incoherent or is a correct response that was obtained by an obviously incorrect
- [22] procedure.
- [23] D
 - [2] 20, and appropriate work is shown, such as $(180 \div 0.9) 180$.
 - [1] A partial answer is found, such as 200 students are enrolled, but 180 is not subtracted from the answer.
 - or [1] An appropriate equation is shown, but one computational error is made, but 180 is subtracted.
 - or [1] An answer of 18 is found by subtracting 180 x 0.9 from 180.
 - or [1] 20, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [24] incorrect procedure.
- [25] A

- [3] No, it will not differ and the student shows that both methods lead to \$47.08, such as \$55 x .80 = \$44, \$44 x 1.07 = \$47.08, \$55 x 1.07 = \$58.85, and \$58.85 x .80 = \$47.08.
- [2] Both ways are computed, one computational mistake is made, and an appropriate answer is found.
- or [2] Both ways are computed correctly, but no comparison is found.
- [1] At least one way is computed correctly, but no comparison is found.
- or [1] Both ways are computed incorrectly, but an appropriate comparison is found.
- [0] Both ways are computed incorrectly, and no comparison is found.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.
- - [2] \$40, and appropriate work is shown.
 - [1] Appropriate work is shown, but one computational error is made.
 - or [1] \$40, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure
- [27] incorrect procedure.

[26]

- [3] \$800, and appropriate work is shown, such as 0.15x + 50 = 170 or a table of values or trial and error with at least three trials and appropriate checks.
- [2] Appropriate work is shown, but one computational error is made.
- or [2] The trial-and-error method is used to find the correct solution, but only two trials and appropriate checks are shown.
- [1] Appropriate work is shown, but two or more computational errors are made.
- or [1] Appropriate work is shown, but one conceptual error is made.
- or [1] Appropriate work is shown, but the \$50 per day is not included in his pay, resulting in an answer of \$1,133.33.
- or [1] The trial-and-error method is attempted and at least six systematic trials and appropriate checks are shown, but no solution is found.
- or [1] \$800, but no work or only one trial with an appropriate check is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [28] incorrect procedure.
 - [2] 44, and appropriate work is shown, such as 0.8(200 145).
 - [1] Appropriate work is shown, but one computational or conceptual error is made. or [1] 44, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [29] incorrect procedure.

- [2] 20.7, and appropriate work is shown, such as $\frac{141288}{683748} = \frac{x}{100}$.
- [1] Appropriate work is shown, but one computational or rounding error is made. or [1] Appropriate work is shown, but one conceptual error is made.
- or [1] 20.7, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [30] incorrect procedure.
- [31] C
- [32] C
 - [3] 1,095 and 1,209, and appropriate work is shown.
 - [2] Appropriate work is shown, but one computational error is made.
 - or [2] Appropriate work is shown, but a whole-number solution is not found. or [2] 5% of CD cases is rounded to 58, but 58 is added to or subtracted from 1,152 appropriately.
 - or [2] Appropriate work is shown, but only one correct solution is found.
 - [1] Appropriate work is shown, but more than one computational error is made.
 - or [1] 5% of CD cases is rounded to 58, but 58 is added to or subtracted from 1,152, but one computational error is made.
 - or [1] 5% of 1,152 is found, but no further work is shown.
 - or [1] 1,095 and 1,209, but no work is shown. [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [33] incorrect procedure.

- [2] 42.85714286 or an equivalent answer, and appropriate work is shown.
- [1] Appropriate work is shown, but one computational or rounding error is made. or [1] An answer of 30 is found by dividing 1.8 by 6.
- or [1] An answer of 70 is found by dividing 4.2 by 6.
- or [1] 42.85714286 or an equivalent answer, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [34] incorrect procedure.
- [35] D
- [36] A
- [37] A
- [38] C
 - [2] 70.92, and appropriate work is shown, such as a proportion.
 - [1] Appropriate work is shown, but one computational or rounding error is made.
 - or [1] Appropriate work is shown, but one conceptual error is made.
 - or [1] 70.92, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [39] incorrect procedure.

[1]	D

[2] D

[2] 75, and appropriate work is shown.

[1] Appropriate work is shown, but one computational error is made.

or [1] An incorrect equation of equal difficulty, such as x + 5x = 180, is solved appropriately, and an appropriate angle measure is found.

or [1] A correct equation is written and solved for x, but no further correct work is shown. or [1] 75, but no work is shown.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously

- [3] incorrect procedure.
- [4] D
- [5] C
- [6] A

[3] 120, and appropriate work is shown, such as 6t + 30 + 8t - 60 = 180.

[2] The student finds correctly the unknown, t = 15, but does not find the measure of angle 4.

or [2] Appropriate work is shown, but one computational error is made.

[1] The student forms an incorrect equation, such as setting the two angles equal, and arrives at t = 45 and an angle of 300.

or [1] 120, but no work is shown.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously

[7] incorrect procedure.

- [3] 95, and appropriate work is shown, such as 3x-20+x+60=180.
- [2] Appropriate work is shown, but one computational error is made.
- or [2] A correct equation is written and solved for x, but $m \angle ROY$ is not found.
- [1] Appropriate work is shown, but two or more computational errors are made.
- or [1] Appropriate work is shown, but one conceptual error is made, such as writing the equation x + 60 = 3x 20, but an appropriate answer is found.
- or [1] A correct equation is written, but no further correct work is shown,
- or [1] 95, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [8] incorrect procedure.
- [9] A
- [10] A

[3] 80, and appropriate work is shown.

[2] x = 30 is shown, but the student fails to substitute to find $m \angle AEC$.

or [2] x = 30 is shown, but the student states that the answer is 100° , by finding the supplement of $\angle AEC$.

or [2] The student makes one computational error in the solution of the correct equation 4x - 40 = x + 50 but appropriately substitutes the incorrect value to solve for $m \angle AEC$.

[1] The student makes one computational error in the solution of the correct equation 4x - 40 = x + 50 and fails to substitute to find $m\angle AEC$.

or [1] The student makes more than one computational error in the solution of the correct equation 4x - 40 = x + 50, but appropriately substitutes the incorrect value to solve for $m \angle AEC$.

or [1] 80, but no work is shown.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously

[11] incorrect procedure.

- [4] 112.5, and appropriate work is shown, such as solving the equation 5x 20 = x + 50.
- [3] Appropriate work is shown, but one computational error is made.
- or [3] $m \angle BED = 67.5$ or $m \angle AEC = 67.5$, but no further correct work is shown.
- [2] Appropriate work is shown, but two or more computational errors are made.
- or [2] Appropriate work is shown, but one conceptual error is made, but an appropriate measure for $\angle CEB$ is found.
- or [2] A correct equation is written and solved for *x*, but no further correct work is shown.
- [1] Appropriate work is shown, but one conceptual error and one computational error are made.
- or [1] 112.5, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [12] incorrect procedure.
- [13] B
- [14] D
- [15] A
 - [3] 65, and appropriate work is shown, such as setting the given angles equal to each other and determining the value of x to be 16, and correct substitution is shown.
 - [2] The given angles are set equal to each other, the correct value of x is determined, but no substitution is shown.
 - or [2] The given angles are set equal to each other, and substitution is shown, but one computational or substitution error is made.
 - [1] The given angles are set equal to each other, but no further work is shown.
 - or [1] An incorrect equation is solved appropriately, such as 5x 15 + 2x + 33 = 180.
 - or [1] 65, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [16] incorrect procedure.

- [17] B
- [18] A
- [19] A
- [20] D
 - [2] 57°, and appropriate work is shown, such as determining that $m\angle FYD \cong m\angle BXY$ and $\angle AXY$ is supplementary to $\angle BXY$.
 - or [2] 57°, and a correctly labeled diagram with appropriate angles is shown.
 - [1] $\angle CYX$ or $\angle BXY$ is determined, but one computational error is made in subtracting to find $m\angle AXY$.
 - or [1] An angle is determined incorrectly, but an appropriate solution is found.
 - or [1] 57°, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [21] incorrect procedure.
 - [2] 31, and appropriate work is shown, such as 5x + 25 = 180.
 - [1] Appropriate work is shown, but one computational error is made.
 - or [1] Appropriate work is shown, but one conceptual error is made, such as setting the given angles equal to each other.
 - or [1] A correct equation is written, but no further correct work is shown.
 - or [1] 31, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [22] <u>incorrect procedure.</u>

- [4] 146, and appropriate work is shown, such as solving the equation 2x = 5x 51.
- [3] Appropriate work is shown, but one computational error is made.
- or [3] The measure of $\angle FHB$ or $\angle DGH$ is found to be 34, and appropriate work is shown, but no further correct work is shown.
- [2] Appropriate work is shown, but two or more computational errors are made.
- or [2] Appropriate work is shown, but one conceptual error is made, such as solving the equation 2x + 5x 51 = 180.
- or [2] The correct equation is solved for x = 17, but no further correct work is shown.
- [1] Appropriate work is shown, but one conceptual error and one computational error are made.
- or [1] The correct equation is written, but no further correct work is shown.
- or [1] 146, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect precedure.
- [23] incorrect procedure.
- [24] B

- [4] $m\angle A = 20$, $m\angle B = 59$, and $m\angle C = 101$, and appropriate work is shown.
- [3] Appropriate work is shown, but one computational error is made.
- or [3] A correct equation is written and solved, and the correct measures for the angles are found, but they are not labeled or are labeled incorrectly.
- [2] Appropriate work is shown, but two or more computational errors are made.
- or [2] Appropriate work is shown, but one conceptual error is made.
- or [2] A correct equation is written and solved for x, but the measures of the angles are not found.
- or [2] An incorrect equation of equal difficulty is solved appropriately, and the three angles are found.
- [1] Appropriate work is shown, but one conceptual error and one computational error are made.
- or [1] A correct equation is written, but no further correct work is shown.

or [1]

 $m\angle A = 20$, $m\angle B = 59$, and $m\angle C = 101$., but no work is shown.

- [0] $m\angle A = 20$ or $m\angle B = 59$ or $m\angle C = 101$, but no work is shown.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an
- [25] obviously incorrect procedure.
- [26] B
- [27] B
- [28] C
- [29] C
- [30] C
- [31] B
- [32] C
- [33] D

- [2] 29, and appropriate work is shown, such as 92 63 = 29.
- [1] The correct application of the exterior angle theorem is shown, but one or more computational errors are made.
- or [1] The correct application of supplementary angles and the sum of the angles of a triangle are shown, but one or more computational errors are made.
- or [1] $m \angle BCA$ is calculated incorrectly, but the sum of the angles in a triangle is used appropriately.
- or [1] 29, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [34] incorrect procedure.
 - [2] 120, and appropriate work is shown, such as $m\angle CDB = 180 130 = 150$ and $m\angle CBA = 70 + 50 = 120$ or correctly labeled angles in a diagram.
 - [1] Appropriate work is shown, but one computational error is made.
 - or [1] Appropriate work is shown, but one conceptual error is made.
 - or [1] $m\angle CBD = 60$ is found, but no further correct work is shown.
 - or [1] 120, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [35] incorrect procedure.
- [36] D

- [2] 32, and appropriate work is shown, such as a diagram or "let" statements and an appropriate equation, such as 5x + 20 = 180. or [2] 32, and an appropriate trial-and-error method with at least two trials and appropriate checks are shown.
- [1] Appropriate work is shown, but one computational error is made.
- or [1] An incorrect equation set equal to 180° is shown, but it is solved appropriately, such as 4x + 20 = 180; or an incorrect equation set equal to 360° is shown, such as 5x + 20 = 360.
- or [1] 32, and an appropriate trial-and-error method with less than two trials and appropriate checks are shown.
- or [1] 32, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.
 - [2] An isosceles triangle that is not acute is drawn, and its three angles are labeled, such as 20, 20, 140 or 45, 45, 90.
 - [1] An isosceles triangle is drawn that shows an angle that is not acute, but the base angles are not labeled.
 - or [1] The three angles are stated correctly, but no triangle is drawn.
 - [0] The triangle that is drawn and labeled is not isosceles or is acute.
 - or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an
- [38] obviously incorrect procedure.

- [3] The student draws an obtuse triangle and all sides and all angles are correctly calculated, such as by using 120°, 30°, and 30° and sides 4, 4, and 10.
- [2] The student has the angles correctly indicated and the two congruent sides marked, but the length of the longest side is incorrect or is missing.
- or [2] All sides are correctly marked, but the angles do not add to 180°, but an obtuse angle and two congruent angles are shown.
- [1] Only the angles are correctly shown.
- or [1] Only the sides are correctly shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [39] incorrect procedure.
 - [3] 135 and appropriate work is shown.
 - [2] The two correct angles of 65° and 70° are found, but their sum is not identified as the answer to the question.
 - or [2] 65° or 70° and an appropriate sum is found.
 - [1] Either the 65° or the 70° is correctly identified.
 - or [1] Two incorrect angle measures are found, but they are added correctly. or [1] 135 and no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [40] incorrect procedure.
 - [2] 30, and appropriate work is shown or an appropriate explanation is given.
 - [1] Angles of the equilateral triangle are shown to be 60° , but *x* is not determined or is determined incorrectly.
 - or [1] 30, but no work is shown or no explanation is given.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [41] incorrect procedure.
- [42] A

- [2] 40, and appropriate work is shown, such as x = 180 (70 + 70) or correctly labeling all the angles in the diagram.
- [1] Appropriate work is shown, but one computational error is made.
- or [1] Appropriate work is shown, but one conceptual error is made.
- or [1] A correct equation is written, but no further correct work is shown.
- or [1] The measures of $\angle ACB$ and $\angle ABC$ are both found to be 70°, but no further correct work is shown.
- or [1] An incorrect equation of equal difficulty is solved appropriately.
- or [1] 40, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [43] incorrect procedure.
- [44] A
- [45] D
 - [2] A correct construction is drawn to find the midpoint of \overline{BC} , showing both sets of arcs and a line connecting A with the midpoint. [1] A correct construction is drawn to find the
 - midpoint of \overline{BC} , but the median is not drawn. or [1] The construction is appropriate, but a compass and a straightedge are not used.
 - [0] No construction arcs are shown.
 - or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an
- [46] obviously incorrect procedure.
 - [2] A correct construction is drawn, showing the arcs intersecting above and below \overline{AB} , and the perpendicular line is drawn.
 - [1] All of the construction arcs are drawn, but the perpendicular line is not drawn.
 - [0] A drawing that is not an appropriate construction is shown.
 - or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an
- [47] obviously incorrect procedure.

- [2] A correct construction is drawn, showing the arcs intersecting above and below \overline{AB} , and line c is drawn.
- [1] A correct construction is drawn, but line c is not labeled.
- [0] A drawing that is not a construction is shown with arc marks sketched.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an abviously incorrect presendant.
- [48] obviously incorrect procedure.
- [49] A
- [50] C
- [51] C
- [52] A
- [53] C
- [54] A
- [55] D
- [56] D
 - [2] The student draws a parallelogram, which is not a rectangle, with four sides and four angles labeled, such as angles of 60, 120, 60, and 120 and sides of 4, 6, 4, and 6.
 - [1] A parallelogram or rhombus, not a square, is drawn, which does not have measures for all lengths or angles.
 - [0] Angles and/or lengths are not appropriate for a parallelogram.
 - or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.
- [57] obviously incorrect procedure.
- [58] <u>D</u>

- [3] 102, and appropriate work is shown, such as using the equation 2x + 10 + 3x = 180 or an equivalent equation.
- [2] The equation 2x + 10 + 3x = 180 is solved correctly for x, but $m \angle B$ is not determined or is determined incorrectly.
- [1] Appropriate work is shown, but one computational error is made or *x* is not determined.
- or [1] The equation 2x + 10 + 3x = 360 is solved correctly, and an answer of 210 is found.
- or [1] 102, but no work is shown.
- [0] The equation 2x + 10 = 3x where x = 10 is given.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an
- [59] obviously incorrect procedure.
 - [3] 120, and appropriate work is shown, such as solving the equation 5x + 2x + 12 = 180.
 - [2] Appropriate work is shown, but one computational error is made.
 - or [2] The correct equation is solved for *x*, but no further correct work is shown.
 - [1] Appropriate work is shown, but two or more computational errors are made.
 - or [1] Appropriate work is shown, but one conceptual error is made.
 - or [1] A correct equation is written, but no further correct work is shown.
 - or [1] An incorrect equation of equal difficulty is solved appropriately, and an appropriate measure is found for $\angle D$.
 - or [1] 120, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [60] incorrect procedure.

- [2] 96, and appropriate work is shown, such as an algebraic solution or a correctly labeled diagram.
- [1] Appropriate work is shown, but one computational error is made.
- or [1] Appropriate work is shown, but one conceptual error is made.
- or [1] 96, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [61] incorrect procedure.
- [62] B
 - [4] 12 and an appropriate method is shown, such as $(AB)^2 = 9^2 + 8^2$.
 - [3] An incorrect length is found for AE, but then it is used to correctly complete the problem.
 - or [3] An appropriate method is shown, but one computational mistake is made.
 - or [3] An appropriate method is shown, but the answer is not given to the nearest foot, such as $\sqrt{145}$.
 - [2] AE = 8 and one computational mistake is made using the Pythagorean theorem.
 - or [2] An incorrect length is found for AE, but then it is used to complete the problem correctly, but the answer is not rounded.
 - [1] AE = 8 is found, but the Pythagorean theorem is not used.
 - or [1] 12 and no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [63] incorrect procedure.
- [64] <u>B</u>
- [65] B

- [4] \$148.54, and appropriate work is shown.
- [3] The correct pre-tax amount of \$137.54 is found, but no tax or an incorrect tax is shown. or [3] Appropriate work is shown, but one computational error is made.
- [2] The correct area of 46 ft² is found, but no cost is shown.
- or [2] Appropriate work is shown, but more than one computational error is made.
- or [2] An incorrect area is determined, such as by adding or multiplying all sides, but then a final cost including tax is determined appropriately.
- [1] An incorrect area is shown, and one computational error is made.
- or [1] \$148.54, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously[66] incorrect procedure.
 - [4] The side equals 2.3 and the area equals 25.5, and appropriate work is shown.
 - [3] Appropriate work is shown, but one computational or rounding error is made.
 - [2] Appropriate work is shown, but one incorrect formula is used, such as using an incorrect trigonometric function, but appropriate answers are found.
 - or [2] Appropriate work is shown to find the correct side, but no further correct work is shown.
 - [1] The radius equals 3 and the central angle equals 45°, but no further correct work is shown.
 - or [1] The side equals 2.3 and the area equals 25.5, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [67] incorrect procedure.
- [68] D
- [69] D
- [70] <u>A</u>
- [71] B

- [72] C
- [73] B
- [74] C
- [75] C
 - a [1] Either (x-2)(x+1)(2x) = V or the same expression without "= V" is shown.
 - or [1] $2x^3 2x^2 4x$ or an equivalent expression is shown.
 - b [1] 864
 - or [1] The student substitutes appropriately into an incorrect part a equation.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously[76] incorrect procedure.
- [3] 3, 12, and 30 and an appropriate arithmetic method or equation is shown, such as $40x^3 = 1080$.
 - [2] An appropriate equation or method is shown, but not all three dimensions are found. or [2] An appropriate method is shown, and although one computational mistake is made, the student does find three dimensions based on this mistake, such as dividing 1080 by 40 incorrectly.
 - [1] The student shows that multiplication is required to find volume but sets up an incorrect method and does not find three dimensions.
 - or [1] 3, 12, and 30 and no work is shown. [0] The sum is used instead of the product, or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.
- [77] obviously incorrect procedure.

- [2] 20, and appropriate work is shown, such as $3,360 \div (14 \times 12)$.
- [1] Appropriate work is shown, but one computational error is made.
- or [1] 20, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect proceed was
- [78] incorrect procedure.
 - [3] 27 and an appropriate method or explanation is shown, such as

$$(\frac{1}{6})(\frac{1}{3})(\frac{2}{3}) = \frac{1}{27}$$
 of a cubic foot, thus 27

bricks needed or, in inches, $\frac{1728}{64} = 27$. A

labeled drawing is an acceptable explanation.
[2] An appropriate method for finding volume is shown, but one computational mistake is made.

- [1] Correct conversion into feet is shown. or [1] The volume of 64 cubic inches is found.
- or [1] 27 and no explanation is given.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.
 - [3] 64, and appropriate work is shown, such as calculating $\frac{(36 \times 144)}{(9 \times 9)}$ or drawing a

labeled diagram.

- [2] Appropriate work is shown, but one computational error is made.
- [1] Appropriate work is shown, but more than one computational error is made.
- or [1] Appropriate work is shown, but one conceptual error is made.
- or [1] 64, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [80] incorrect procedure.

- [3] 5-inch box and appropriate work is shown, including showing a diameter between 4 and 5.
- [2] The correct diameter is shown, but the wrong box size is chosen.
- or [2] The correct radius is shown, but the 3inch box is chosen.
- [1] The correct diameter or radius is shown, but no box is chosen.
- or [1] An appropriate radius between 2 and 3 is shown, using the incorrect formula

 $A = \pi r^2$, and the 3-inch box is chosen.

or [1] An appropriate diameter, using

 $A = \pi r^2$, is shown, but the appropriate box is chosen.

- or [1] An appropriate radius, using $A = \pi r^2$, is shown, but no box is chosen.
- or [1] The 5-inch box is chosen, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [81] incorrect procedure.
 - [3] 47, and appropriate work is shown.
 - [2] Appropriate work is shown, but one computational or rounding error is made. or [2] The correct numerical value of the volume of the cup (20π or its equivalent) and the volume of the tank (3,000) are shown, but the solution is not completed.
 - [1] The correct volume of only the cup or only the tub is shown.
 - or [1] 47, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously [82] incorrect procedure.

- [3] 2.6, and appropriate work is shown, such as $(5 \bullet 5 \bullet 5) = (7 \bullet 7)h$.
- [2] Appropriate work is shown, but one computational or rounding error is made.
- [1] Appropriate work is shown, but two or more computational or rounding errors are made.
- or [1] Appropriate work is shown, but one conceptual error is made, such as using an incorrect formula.
- or [1] The volume of both of the cubes is found correctly, but no further correct work is
- or [1] 2.6, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [83] incorrect procedure.
 - [3] 12, and appropriate work is shown, such as calculating volume = 5.760 in^3 and dividing by 500 in³.
 - [2] Appropriate work is shown, but one computational or rounding error is made. or [2] The volume is found incorrectly by multiplying $24 \times 16 \times 18$, but it is divided by 500 and rounded appropriately, resulting in an answer of 14.
 - [1] Appropriate work is shown, but two or more computational or rounding errors are made.
 - or [1] Appropriate work is shown, but one conceptual error is made.
 - or [1] The volume of 5,760 is found correctly, but no further correct work is shown.
 - or [1] 12, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [84] incorrect procedure.

- [4] 11, and appropriate work is shown, such as solving the quadratic equation 3x(x+5) = 150 or trial and error with at least three trials and appropriate checks.
- [3] Appropriate work is shown, but one computational error is made.
- or [3] Appropriate work is shown to determine that 5 is the shorter side of the box, but the shorter side of the original sheet is not found or is found incorrectly.
- or [3] An incorrect quadratic equation of equal difficulty is solved appropriately, and an appropriate shorter side of the original sheet is found.
- [2] Appropriate work is shown, but more than one computational error is made.
- or [2] Appropriate work is shown, but one conceptual error is made.
- or [2] An incorrect quadratic equation of equal difficulty is solved appropriately, but the shorter side of the original sheet is not found.
- or [2] A correct quadratic equation is set equal to zero, but no further correct work is shown.
- or [2] The trial-and-error method is used to find a correct solution, but only two trials and appropriate checks are shown.
- [1] Appropriate work is shown, but one conceptual error and one computational error are made.
- or [1] One conceptual error is made in finding the shorter side of the box, and the corresponding shorter side of the original sheet is not found or is found incorrectly. or [1] A correct quadratic equation is written, but it is not set equal to zero, and no further correct work is shown.
- or [1] 11, but no work or only one trial with an appropriate check is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously [85] incorrect procedure.

- [4] 21 by 23, and appropriate work is shown, such as solving the equation 765 = 3(x-4)(x-6).
- [3] Appropriate work is shown, but one computational error is made.
- or [3] Appropriate work is shown, but only one dimension is found.
- [2] Appropriate work is shown, but two or more computational errors are made.
- or [2] Appropriate work is shown, but one conceptual error is made.
- or [2] An incorrect equation of equal difficulty is solved appropriately, and appropriate dimensions are found.
- or [2] A correct quadratic equation is written in standard form, but no further correct work is shown.
- [1] Appropriate work is shown, but one conceptual error and one computational error are made.
- or [1] An incorrect equation of equal difficulty is written, and one computational error is made, but appropriate dimensions are found.
- or [1] An incorrect equation of equal difficulty is solved appropriately, but one computational error is made when finding the length.
- or [1] 21 by 23, but no work is shown.
- [0] 21 or 23, but no work is shown.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an
- [86] obviously incorrect procedure.

- [2] 2.6, and appropriate work is shown, such as solving the equation $(10+x)^3 = 2000$.
- [1] Appropriate work is shown, but one computational or rounding error is made.
- or [1] Appropriate work is shown, but one conceptual error is made.
- or [1] The equation $(10+x)^3 = 2000$ is written, but no further correct work is shown.
- or [1] An incorrect equation of equal difficulty is solved appropriately.
- or [1] 2.6, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [87] incorrect procedure.

vw.ji	nap.org		
[1] [2]	<u>A</u> <u>A</u>		[2] 8 and the use of trigonometry, the Pythagorean theorem, or Pythagorean triple is shown.
[3]	<u>C</u>		 [1] The Pythagorean theorem or trigonometry is used, but a computational mistake is made or substitution is incorrect, such as 6² = 10² + x². [1] 8 and no work is shown. [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.
[4]	<u>D</u>		
[5]	<u>C</u>		
[6]	[3] $14\sqrt{2}$, and appropriate work is shown, such as using the Pythagorean theorem or drawing a correctly labeled diagram that shows the isosceles right triangle. [2] Appropriate work is shown, but one computational error is made. or [2] Appropriate work is shown, but the answer is expressed as a decimal or the radical is not simplified. [1] Appropriate work is shown, but two or more computational errors are made. or [1] Appropriate work is shown, but one computational error is made, and the answer is not expressed as a radical in simplest form. or [1] Appropriate work is shown, but one conceptual error is made. or [1] 14, the side of the square is found correctly, but no further correct work is shown. or [1] $14\sqrt{2}$, but no work is shown. [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.		
		[9]	
		[10]	[2] 16, and appropriate work is shown, such as the Pythagorean theorem, the Pythagorean triple, or trigonometry. [1] Appropriate work is shown, but one computational error is made. or [1] Appropriate work is shown, but one conceptual error is made, such as using an incorrect trigonometric function. or [1] 16, but no work is shown. [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.
[7]			[2] 15, and appropriate work is shown, such as using the Pythagorean theorem, Pythagorean triples, or trigonometric functions. [1] The data are substituted incorrectly, but an appropriate answer is found and is rounded correctly. or [1] Appropriate work is shown, but one or
[8]	С		more computational errors are made. or [1] 15, but no work is shown.
			[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
		[11]	incorrect procedure.
		[12]	<u>C</u>
		[13]	A

- [2] 2.8, and appropriate work is shown, such as $3^2 = 1^2 + (BC)^2$.
- [1] Appropriate work is shown, but one computational or rounding error is made. or [1] Appropriate work is shown, but one conceptual error is made.
- or [1] The length of BD is found to be 3, but no further correct work is shown.
- or [1] 2.8, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- incorrect procedure.
 - [3] 3, and appropriate work is shown, such as using a 3:4:5 right triangle, correct proportions, or the Pythagorean theorem with a proportion.
 - [2] Appropriate work is shown, and the value of the side is determined to be 5, but n = 3 is not found.
 - [1] A correct proportion is set up, but no answer or an incorrect answer is found. or [1] 3, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [15] incorrect procedure.
- [16] A
 - [2] An appropriate explanation is written, such as defining special isosceles right triangles, or appropriate work is shown, such as using legs of six and finding the hypotenuse.
 - [1] Appropriate work is shown, but one computational error is made.
 - or [1] Appropriate work is shown, but one conceptual error is made.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- incorrect procedure. [17]

- [2] $\sqrt{171}$ or 13 or 13.1 or 13.08 or an equivalent answer, and appropriate work is shown, such as the use of the equation of a circle $(x^2 + y^2 = r^2)$ or the Pythagorean
- [1] Appropriate work is shown, but one computational error is made.
- or [1] Incorrect analysis is shown, such as x =5 and y = 14, but the work is concluded appropriately.
- or [1] A correct answer is found, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously [18] incorrect procedure.
 - [4] 9.4, and appropriate work is shown, such as the use of the Pythagorean theorem.
 - [3] Appropriate work is shown, but one computational or rounding error is made.
 - [2] Appropriate work is shown, but more than one computational or rounding error is made.
 - or [2] Appropriate work is shown, but one conceptual error is made.
 - or [2] An incorrect diagonal of the base is found, but an appropriate solution is found. or [2] Only the diagonal of the base is found correctly, but appropriate work is shown, such as $3^2 + 4^2 = d^2$ or use of 3-4-5 right triangles.
 - [1] Appropriate work is shown, but one conceptual error and one computational or rounding error are made.
 - or [1] The Pythagorean theorem is used to find the length of the straw, but the appropriate legs are not used. or [1] 9.4, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [19] incorrect procedure.
- [20] D
- [21] A
- [22] D

[23] A

[3] 109 meters and appropriate work is shown by using an appropriate trigonometric ratio,

such as
$$\tan 32^\circ = \frac{y}{175}$$
.

[2] 109 meters but one rounding error is made.

or [2] The student uses an appropriate trigonometric function with an inverted ratio,

such as
$$\tan 32^\circ = \frac{175}{y}$$
, but completes the

calculation appropriately, such as showing 280 meters.

[1] The student uses an incorrect trigonometric ratio but completes the calculation appropriately.

or [1] The student uses an inverted tangent ratio and makes one computational or rounding error.

or [1] The student uses the correct trigonometric ratio but solves it incorrectly or does not solve it at all.

or [1] 109 meters but no work or explanation is shown.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously [24] incorrect procedure.

[4] 116 and an appropriate method is shown.

[3] An appropriate method is shown, but the answer is left in an inappropriate form, such as 116.2.

or [3] An appropriate method is shown, but 3 feet is not added, and the answer is left 113.

or [3] Tangent function is used, but computational mistakes are made, but 3 feet is added to the incorrect value and the answer is found correctly.

[2] An incorrect trigonometric function is used, 3 feet is added, and the answer is rounded correctly.

or [2] Tangent function is used, but computational mistakes are made, and 3 feet is not added to an incorrect answer.

[1] 116 and no work is shown.

or [1] An incorrect trigonometric function is used, and 3 feet is added to the incorrect answer, but the answer is rounded incorrectly.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously

[25] incorrect procedure.

[26] B

[4] 28.2, and an appropriate equation is shown, such as $\tan 62 = \frac{x}{15}$.

[3] Appropriate work is shown, but the answer is rounded incorrectly.

or [3] The student uses the correct tangent function and rounds the answer, but makes one computational error.

[2] The student uses the correct tangent function, but makes several errors.

or [2] An incorrect trigonometric function is used, but appropriate work is shown.

[1] The tangent function is indicated, but the ratio is set up incorrectly.

or [1] 28.2, but no work is shown.


[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously

[27] incorrect procedure.

- [2] 117.6, and appropriate work is shown, such as $\tan 78^\circ = \frac{x}{25}$.
- [1] Appropriate work is shown, but one computational or rounding error is made. or [1] Appropriate work is shown, but one conceptual error is made, such as using an incorrect trigonometric function, but an appropriate solution is found. or [1] A correct trigonometric equation is written, but no further correct work is shown. or [1] 117.6, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure
- [28] incorrect procedure.
 - [2] 10, and appropriate work is shown. [1] Appropriate work is shown, but one
 - computational or rounding error is made. or [1] Appropriate work is shown, but one
 - conceptual error is made, such as using an incorrect trigonometric function.
 - or [1] Appropriate work is shown, but the length of the ladder is found.
 - or [1] 10, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.
- [29]

- [3] 45, and appropriate work is shown, such as $\tan 66^\circ = \frac{x}{20}$.
- [2] A correct trigonometric ratio is used, and values are substituted correctly, but one computational or rounding error is made, or the calculator is left in radian mode.
- [1] Appropriate work is shown, but two or more computational or rounding errors are made.
- or [1] Appropriate work is shown, but one conceptual error is made, such as using an incorrect trigonometric ratio.
- or [1] An incorrect diagram is drawn, but an appropriate solution is found.
- or [1] A correctly labeled diagram is drawn, but no further correct work is shown.
- or [1] A correct trigonometric ratio is written, but no further correct work is shown.
- or [1] 45, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.
- [30] incorrect procedure.

[4] 2,058, and appropriate work is shown, such as the accompanying diagram and equation.



- [3] Appropriate work is shown, including a correct diagram and the use of the tangent function, but one computational error is made. or [3] Appropriate work is shown, including a correct diagram and the use of the tangent function, but the answer is not rounded or is rounded incorrectly.
- [2] A correct diagram is drawn, but an incorrect trigonometric function is selected, but it is solved and rounded appropriately. or [2] A correct diagram is drawn and the tangent function is selected, but no further work is shown.
- or [2] An incorrect diagram is drawn, but the appropriate trigonometric function, based on the drawing, is selected, solved, and rounded appropriately.
- [1] An incorrect diagram is drawn and an incorrect trigonometric function is selected, but it is solved and rounded appropriately.
- or [1] Only a correct diagram is drawn.
- or [1] 2,058, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure
- [31] incorrect procedure.

[4] 114" (9 feet 6 inches) and 37" (3 feet 1 inch) and appropriate work is shown, such as

$$\sin 72^\circ = \frac{x}{10}$$
 and $\cos 72^\circ = \frac{y}{10}$ or use of the

Pythagorean theorem.

- [3] An incorrect diagram is drawn, but appropriate work and an appropriate solution for that diagram are shown.
- or [3] Appropriate work is shown, but the answers are rounded to the nearest foot and then converted to inches, arriving at 120" and 36".
- or [3] The setup is correct, but the answers are not converted to the nearest inch.
- [2] One correct dimension is shown, such as 114" (9 feet 6 inches) or 37" (3 feet 1 inch). or [2] Only one error involving interchanging sine and cosine is made.
- or [2] An incorrect diagram is drawn, and the solution is appropriate for the diagram but is not rounded to the nearest inch.
- [1] The student switches sine and cosine and does not round to the nearest inch.
- or [1] The student uses the correct trigonometric function to compute one side correctly but does not convert it to the nearest inch.
- or [1] 114" (9 feet 6 inches) and 37" (3 feet 1 inch) but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [32] incorrect procedure.

[4] x = 19.62990915 and y = 9.814954576 or equivalent answers, and appropriate work is shown, such as $\sin 60^\circ = \frac{17}{x}$ and

$$\tan 60^\circ = \frac{17}{y}$$
 or the Pythagorean theorem.

[3] Appropriate work is shown, but one computational or rounding error is made. or [3] Appropriate work is shown, and the correct answers are found, but not identified. [2] Appropriate work is shown, but one conceptual error is made, such as

$$\sin 60^\circ = \frac{x}{17}.$$

- or [2] Appropriate work is shown, but more than one computational or rounding error is made.
- [1] Appropriate work is shown, but two conceptual errors are made, such as

$$\sin 60^{\circ} = \frac{x}{17}$$
 and $\tan 60^{\circ} = \frac{y}{17}$.

or [1] x = 19.62990915 and y = 9.814954576 or equivalent answers, but no work is shown. [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.

[4] Length of ladder =11 and distance from the base of the ladder to the wall = 4, and appropriate work is shown, such as using sine and then tangent or the Pythagorean theorem.

[3] Appropriate work is shown, but one computational or rounding error is made.

- or [3] Appropriate work is shown, but the correct answers are not labeled or are labeled incorrectly.
- [2] Appropriate work is shown, but two or more computational or rounding errors are made.
- or [2] Appropriate work is shown, but one conceptual error is made, such as using one incorrect trigonometric ratio.
- or [2] Appropriate work is shown, but only the length of the ladder or the distance from the base of the ladder to the wall is found. or [2] Two correct trigonometric equations are written, but no further correct work is shown.
- [1] Appropriate work is shown, but one conceptual error and one computational or rounding error are made.
- or [1] Only one correct trigonometric equation is written, and no further correct work is shown.
- or [1] Length of ladder = 11 and distance from the base of the ladder to the wall = 4, but no work is shown.
- [0] Length of ladder = 11 or distance from the base of the ladder to the wall = 4, but no work is shown.
- or [0] 11 and 4, but no work is shown, and the solutions are not labeled.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an

[34] obviously incorrect procedure.

[3] 12 and the equation
$$\tan x = \frac{420}{2000} = .21$$
 is shown.

or [3] 12 and the Pythagorean theorem and an appropriate trigonometric function are correctly used.

[2] Tan function is correctly used, but the answer is not rounded, such as 11.859. or [2] The setup is correct, but one computational mistake is made, and an appropriate angle is found.

or [2] The answer is incorrectly expressed, such as $\tan x = 12$.

[1] The tan function is set up correctly, but the angle is not computed.

or [1] 12 and no work is shown.

or [1] 12 and
$$\sin x = \frac{420}{2000}$$
 is used.

or [1] 78 and
$$\cos x = \frac{420}{2000}$$
 is used.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously [35] incorrect procedure.

> [2] 25.4, and appropriate work is shown, such as solving the equation $\sin x = \frac{3}{7}$.

[1] Appropriate work is shown, but one computational or rounding error is made. or [1] Appropriate work is shown, but one conceptual error is made, such as using an incorrect trigonometric function. or [1] A correct trigonometric equation is written, but no further correct work is shown. or [1] 25.4, but no work is shown. [0] A zero response is completely incorrect,

irrelevant, or incoherent or is a correct response that was obtained by an obviously

incorrect procedure. [36]

a [2] 59, and the equation
$$\tan x = \frac{280}{170}$$
 is

shown, or the Pythagorean theorem is used first to find the hypotenuse, and either sine or cosine is used correctly to find x.

[1] Appropriate work is shown, but one computational or rounding error is made. or [1] 59, but no work is shown.

b [2] 122, if the Pythagorean theorem is used or if a trigonometric function of the angle is used before it was rounded to 59°.

or [2] 120, if
$$\cos 59 = \frac{170}{hyp}$$
 is used.

or [2] 123, if
$$\sin 59 = \frac{170}{hyp}$$
 is used.

[1] Appropriate work is shown, but one computational or rounding error is made. or [1] 122 or 120 or 123, but no work is shown.

a and b [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.

[37]

a [2] 56, and appropriate work is shown, such as $\tan A = \frac{6}{4}$ or finding the hypotenuse and

then using sine or cosine or using proportional sides of similar triangles.

- [1] Appropriate work is shown, but one computational or rounding error is made. or [1] Appropriate work is shown, but one
- conceptual error is made.
- or [1] The length of the hypotenuse is found correctly, but no further correct work is shown.
- or [1] 56, but no work is shown.
- b [2] 12, and appropriate work is shown, such

as
$$\sin 56 = \frac{h}{15}$$
.

- or [2] An appropriate answer is found based on an incorrect angle found in part a.
- [1] Appropriate work is shown, but one computational or rounding error is made.
- or [1] Appropriate work is shown, but one conceptual error is made.
- or [1] 12, but no work is shown. a and b [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.

[4] 32, and appropriate work is shown, such

as
$$12^2 + 16^2 = r^2$$
, $50 - r = s$, and $\sin x = \frac{16}{30}$.

- [3] Appropriate work is shown, but one computational error is made.
- or [3] Appropriate work is shown to find r = 20 and s = 30 and the trigonometric equation

$$\sin x = \frac{16}{30}$$
 is written, but it is not solved or is

solved incorrectly.

- [2] Appropriate work is shown, but two or more computational errors are made.
- or [2] Appropriate work is shown, but one conceptual error is made, such as using an incorrect trigonometric function to find the angle.
- or [2] The lengths of rand s are found correctly, but no further correct work is shown.
- or [2] Incorrect lengths are found for rand s, but the sine function is used correctly to find an appropriate angle.
- [1] Appropriate work is shown, but one conceptual error and one computational error are made.
- or [1] The length of r is found correctly, but no further correct work is shown.
- or [1] 32, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [39] incorrect procedure.

[4] 153, and appropriate work is shown, such

as
$$\sin 50^{\circ} = \frac{x}{200}$$
.

[40]

- [3] An appropriate analysis is shown, but one computational or rounding error is made.
- [2] An incorrect trigonometric function is

used, such as $\cos 50^\circ = \frac{x}{200}$, but it is carried

to an appropriate final answer and is rounded correctly.

- [1] An incorrect trigonometric function is used and solved appropriately, but it is rounded incorrectly.
- or [1] Only an appropriate diagram is shown.
- or [1] 153, but no work is shown.
 [0] Use of the Pythagorean theorem, such as
- [0] Use of the Pythagorean theorem, such as $200^2 = 50^2 + x^2$, is shown.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.

[4] 79.4, and appropriate work is shown, such

as
$$\tan 52 = \frac{x}{62}$$
.

- [3] Appropriate work is shown, but one computational or rounding error is made. or [3] An incorrectly labeled diagram is drawn, but the appropriate trigonometric function is used, and an appropriate answer is found.
- [2] Appropriate work is shown, but two or more computational or rounding errors are made.
- or [2] Appropriate work is shown, but one conceptual error is made, such as using an incorrect trigonometric function or ratio.
- [1] Appropriate work is shown, but one conceptual error and one computational or rounding error are made.
- or [1] A correctly labeled diagram is drawn, but no further correct work is shown.
- or [1] A correct equation is written, but no further correct work is shown.
- or [1] An incorrectly labeled diagram is drawn, but an appropriate equation is written, but no further correct work is shown.
- or [1] 79.4, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [41] incorrect procedure.

- [4] 41.4, and appropriate work is shown, such as 200 tan 28°-200 tan 18°.
- [3] Appropriate work is shown, but one computational or rounding error is made. or [3] Appropriate work is shown to find the correct height of the cliff and the correct combined height of the lighthouse and the cliff, but they are not subtracted.
- [2] Appropriate work is shown, but two or more computational or rounding errors are made.
- [2] Appropriate work is shown, but one conceptual error is made, such as using an incorrect trigonometric function.
- or [2] Appropriate work is shown to find the correct height of the cliff or the correct combined height of the lighthouse and the cliff, but no further correct work is shown.
- [1] Appropriate work is shown, but one conceptual error and one computational or rounding error are made.
- or [1] A correct equation is written to find the height of the lighthouse, but no further correct work is shown.
- or [1] 41.4, but no work is shown.
- [0] The correct height of the cliff or the correct combined height of the lighthouse and cliff is found, but no work is shown. or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an
- [42] obviously incorrect procedure.

- [1] D
- [2] B
- [3] D
- [4] C
- [5] B
- [6] A
- [7] A
- [8] A
- [9] A
- [10] B
- [11] C
- [12] D
- [13] B
- [14] D
 - [2] 12 and an appropriate explanation is given.
 - [1] The student uses an appropriate method, such as showing $\frac{k-2}{3-1} = 5$ or graphing of a

line through (1,2) having a slope of 5, but the correct answer is not found.

- or [1] 12 and no explanation is given.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [15] incorrect procedure.
- [16] <u>C</u>
- [17] <u>D</u>
- [18] A
- [19] B
- [20] <u>C</u>
- [21] B

- [22] C
- [23] D
- [24] B
 - [2] A correct equation is written, such as $y = -\frac{3}{2}x + 4$ or $(y-4) = -\frac{3}{2}(x-0)$.
 - [1] An appropriate equation is written, but one computational error is made or one incorrect substitution is made.
 - [1] An appropriate equation is written, but one conceptual error is made, such as writing an equation for a parallel line going through (0,4) or for a perpendicular line that does not go through (0,4).
 - or [1] The slope is identified correctly as $-\frac{3}{2}$

or the *y*-intercept as 4, but no equation or an incorrect equation is written.

- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [25] incorrect procedure.
 - [3] Three correct equations are shown, such as y = x + 7, y = -x 6, and 2y = 2x 12.
 - [2] Only two correct equations are shown.
 - [1] Only one correct equation is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [26] incorrect procedure.
- [27] C
- [28] A
- [29] <u>A</u>
- [30] <u>C</u>
- [31] A
- [32] C
- [33] D
- [34] B

vww.jr	nap.org		- ··· - ··· · · ·	
[35]	C			[2] 5,279.61, and ap
[36]	В			such as $3,500(1+\frac{0.0}{1})$
[37]	<u>A</u>			[1] Appropriate work
[38]	<u>A</u>			or [1] 5,279.61, but 1 [0] A zero response
[39]	<u>B</u>			irrelevant, or incohe
[40]	В		[50]	response that was ob- incorrect procedure.
[41]	В			[2] 65, and appropria
[42]	C			as $P(10) = 80(0.98)^{10}$
[43]	A			[1] Appropriate work
[44]	D			or [1] Appropriate w
[45]	Α			or [1] 65, but no wor
[46]	A			[0] A zero response irrelevant, or incohe
[47]	A			response that was ob
		:	[51]	incorrect procedure.
[48]	C		[52]	<u>A</u>
		48, and appropriate work is shown, providing a correctly labeled table or	[53]	<u>B</u>
	solving	the equation $(1.39)(1.005)^{12} = C$.		[2] 7,800, and appro
		propriate work is shown, but one		[1] Appropriate work
	-	ational or rounding error is made.		computational or rou
		Appropriate work is shown, but one		or [1] 7,800, but no
	-	tual error is made, such as using 1.05 or using an incorrect exponent.		[0] A zero response irrelevant, or incoher
		A correct equation is written, but no		response that was ob
		correct work is shown.	[54]	incorrect procedure.
	or [1] A	An incorrect equation of equal		
		ty is solved appropriately.	[55]	<u>A</u>
		1.48, but no work is shown.		
	[0] A z	ero response is completely incorrect,		

irrelevant, or incoherent or is a correct response that was obtained by an obviously

[49] incorrect procedure.

[2] 5,279.61, and appropriate work is shown, such as $3,500(1+\frac{0.0825}{12})^{(12\times5)}$. [1] Appropriate work is shown, but one computational or substitution error is made. or [1] 5,279.61, but no work is shown. [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously [50] incorrect procedure. [2] 65, and appropriate work is shown, such as $P(10) = 80(0.98)^{10}$. [1] Appropriate work is shown, but one computational or rounding error is made. or [1] Appropriate work is shown, but one conceptual error is made. or [1] 65, but no work is shown. [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously [51] incorrect procedure. [52] A [53] B [2] 7,800, and appropriate work is shown. [1] Appropriate work is shown, but one computational or rounding error is made. or [1] 7,800, but no work is shown. [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously

[1] C

- [3] The student says the point does not lie on the line and an appropriate method is shown, such as slope of -2 does not work with the new point (-25,81) and either other point (0,4) or (2,0), or accurately shows a graph where (-25,81) is not on line ℓ .
- [2] The student says the point does not lie on the line but gives an inappropriate explanation of slope.
- or [2] The student tries to use slope concept but makes one computational mistake and gives an appropriate answer based on this mistake.
- [1] Only the slope of -2 is found.
- or [1] The correct diagram is drawn with no interpretation.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure
- [2] incorrect procedure.
 - [2] \$390 or 390 and appropriate work is shown, such as a numerical table or the equation y = 30x + 90 or the expression 90 + 30N.
 - [1] Appropriate work is shown, but one computational error is made.
 - or [1] \$300 or 300 or a slope of 30 but appropriate work is shown.
 - or [1] \$390 or 390 but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [3] incorrect procedure.
- [4] B
- [5] D
- [6] D
- [7] <u>A</u>
- [8] A
- [9] A
- [10] B

- [4] \$1.50 for one slice of pizza and \$0.75 for one cola, and appropriate work is shown, such as 3x + 2y = \$6 and 2x + 3y = \$5.25.
- [3] Appropriate work is shown, but one computational error is made.
- or [3] Appropriate work is shown, but only the price of one slice of pizza or the price of one cola is found correctly.
- [2] Appropriate work is shown, but more than one computational error is made.
- or [2] An incorrect system of equations of equal difficulty is solved appropriately to calculate the cost of one slice of pizza and one cola.
- [1] \$1.50 for one slice of pizza and \$0.75 for one cola, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.
- [11] incorrect procedure.
 - [4] Milk Chocolate bar = \$0.75 and Creamy Nougat bar = \$0.50, and appropriate work is shown, such as equations, a trial-and-error method with at least two trials and appropriate checks, or an algebraic or graphic solution.
 - [3] Appropriate work is shown, but one computational error is made.
 - [2] The cost of one candy bar is determined correctly with appropriate work shown, but no attempt is made to find the cost of the other candy bar.
 - or [2] Appropriate work is shown, but more than one computational error is made.
 - [1] Appropriate work is shown, but no answer is found.
 - or [1] Milk Chocolate bar = \$0.75 and Creamy Nougat bar = \$0.50, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [12] incorrect procedure.

- [4] One doughnut is \$0.75 and one cookie is \$0.60, and appropriate work is shown, such as a system of equations, trial and error with at least three trials and appropriate checks, or a table.
- [3] Appropriate work is shown, but one computational error is made.
- or [3] Appropriate work is shown, but only one correct answer is found, or two correct answers are found, but they are not identified clearly as doughnuts or cookies, or the doughnuts and cookies are labeled incorrectly.
- [2] Appropriate work is shown, but more than one computational error is made.
- or [2] Two equations are written, one correct and one incorrect, but two appropriate answers are found.
- or [2] The trial-and-error method is used to find a correct solution, but only two trials and appropriate checks are shown.
- [1] Two correct equations are written, but no further correct work is shown.
- or [1] One doughnut is \$0.75 and one cookie is \$0.60, but no work or only one trial with an appropriate check is shown.
- [0] One correct equation is shown, and no answer or only one appropriate answer is found.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an
- obviously incorrect procedure.

- [4] \$5 for the sprayer and \$10 for the generator, and appropriate work is shown, such as x = hourly cost of sprayer and y =hourly cost of generator, and an appropriate system of equations is solved or a trial-anderror method is used, showing at least two trials with appropriate checks.
- [3] Both correct equations are shown or an appropriate chart or trial-and-error method is used, but one computational error is made. or [3] Both correct equations are shown, and they are solved for one value, but no further work is shown.
- [2] Only one of the two equations is correct, but they are solved appropriately for both
- or [2] Both correct equations are shown, but more than one computational error is made. or [2] \$5 for the sprayer and \$10 for the generator, but only one trial is shown with appropriate checks.
- [1] Both equations are incorrect, but they are solved appropriately for both values.
- or [1] Both correct equations are shown, but they are not solved.
- or [1] \$5 for the sprayer and \$10 for the generator, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously

[14]	incorrect procedure

[15]	D

[16] D

[17] D

- a [2] An appropriate equation or system is shown, such as x + y = 148 and 12x + 9y =1410 or one equation such as 12(148 - x) + 9x= 1410 with variables identified.
- [1] The student shows appropriate equation(s), but variables are not defined. or [1] One mistake in equation(s) is made, or only one equation with two variables is shown, but variables are defined.
- b(1) [1] 26 and an appropriate method is shown, such as solving the equation or making a table.
- or [1] An appropriate answer is found based on incorrect equation(s) obtained in part a. b(2) [1] 122 and an appropriate method is shown, such as 148 - 26.
- or [1] An appropriate answer is found based on incorrect equation(s) obtained in part a. b (1) and b (2) [1] 26 and 122 and no work is shown.
- a and b [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.

[18]

- [3] Seth had 101, Jason had 51, and Raoul had 104, and appropriate work is shown, such as x + 25 = (2x - 1) - 25 or trial and error with at least three trials and appropriate checks.
- [2] Appropriate work is shown, but one computational error is made.
- or [2] 101, 51, and 104, and appropriate work is shown, but the solutions are not labeled or are labeled incorrectly.
- or [2] A correct equation is solved, but the number of CDs for only one boy is found.
- or [2] The trial-and-error method is used to find a correct solution, but only two trials and appropriate checks are shown.
- [1] Appropriate work is shown, but more than one computational error is made.
- or [1] Appropriate work is shown, but one conceptual error is made, but an appropriate number of CDs is found for each boy.
- or [1] A correct equation is written, but no further correct work is shown.
- or [1] Seth had 101, Jason had 51, and Raoul had 104, but no work or only one trial with an appropriate check is shown.
- [0] Seth had 101 or Jason had 51 or Raoul had 104, but no work is shown.
- or [0] 101, 51, and 104, but no work is shown and the solutions are not labeled or are labeled incorrectly.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an [19] obviously incorrect procedure.

- [3] 374 grasshoppers and 187 crickets, and appropriate work is shown.
- [2] An appropriate equation is solved or appropriate work is shown, but only one correct answer is found, or two correct answers are found but they are not identified clearly as grasshoppers or crickets, or the grasshoppers and crickets are labeled incorrectly.
- or [2] Appropriate work is shown, but one computational error is made.
- [1] Appropriate work is shown, but more than one computational error is made.
- or [1] An incorrect equation of equal difficulty is solved appropriately.
- or [1] 374 grasshoppers and 187 crickets, but no work is shown.
- [0] 374 and 187, but no work is shown, and the answers are not identified clearly as grasshoppers or crickets.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an [20] obviously incorrect procedure.
- [2] 4.5 and an appropriate method is shown, such as the equation 3x + x + 2 = 20 or some trial and error or arithmetic process.
 - [1] An appropriate method is shown, but the correct answer is not found.
 - or [1] 4.5 and no work is shown.
 - or [1] The student solves the equation
 - x + 3x 2 = 20 and answers 5.5.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [21] incorrect procedure.

- [2] 42 nickels and 21 dimes, and appropriate work is shown, such as 0.1x + (0.05)2x = 4.20 or a guess and a check with a minimum of two trials and appropriate checks or another appropriate method.
- [1] 42 nickels or 21 dimes, but appropriate work is shown.
- or [1] Appropriate work is shown, but no answer or an incorrect answer is found.
- or [1] 42 nickels and 21 dimes, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.

[22]

- [3] Four 20-cent and eight 32-cent stamps, and appropriate work is shown, such as a system of equations, or a linear equation such as 2x(.32) + .20x = 3.36, or trial and error with at least three trials and appropriate checks.
- [2] Appropriate work is shown, but one computational error is made, but appropriate quantities are found for each stamp.
- or [2] Appropriate work is shown, but the quantity for only one of the stamps is found.
- or [2] Appropriate work is shown, but the solutions are not labeled or the labels are reversed.
- or [2] The trial-and-error method is used to find correct solutions, but only two trials and appropriate checks are shown.
- [1] Appropriate work is shown, but two or more computational errors are made, but appropriate quantities are found for each stamp.
- or [1] The trial-and-error method is attempted, and at least six systematic trials and appropriate checks are shown, but no solution is found.
- or [1] An incorrect equation or system of equations of equal difficulty is solved appropriately for both solutions.
- or [1] A correct equation or system of equations is written, but no further correct work is shown.
- or [1] Four 20-cent and eight 32-cent stamps, but no work or only one trial with an appropriate check is shown.
- [0] Four and eight, but no work is shown, and the solutions are not labeled.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.

[23]

- [4] 65 adult tickets and 85 student tickets and an appropriate equation is shown, such as 7.50x + 4.75(150 x) = 891.25, or any other acceptable method is used.
- [3] Either 65 or 85 and appropriate work is shown.
- or [3] Appropriate work is shown, but one computational error is made that leads to two appropriate answers.
- [2] An incorrect equation is shown, but it is solved appropriately for two answers.
- or [2] The correct equation is shown, but two computational errors are made.
- [1] Appropriate work is shown, but no answer is found.
- or [1] 65 and 85 but no work is shown.
- [0] Either 65 or 85 and no work is shown. or [0] A zero response is completely
- incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure
- [24] obviously incorrect procedure.
 - a [2] The student writes an appropriate system of equations, such as b = f + 100 and 4b + 12f = 3,056, and defines the variables.
 - or [2] The student writes an appropriate equation, such as 4(100 + x) + 12x = 3,056, and defines the variable.
 - [1] A correct equation or correct equations are shown, but the variables are not defined. or [1] One error is made in the setup, such as b + f = 100.
 - [0] The student only defines the variables. b [2] 266, and appropriate work is shown, using an algebraic solution or a correct trialand-error method.
 - or [2] Appropriate work is shown for an incorrect part a equation or system of equations.
 - [1] Work is shown, but the answer is inappropriate, such as \$1,064.
 - or [1] 266, but no work is shown. a and b [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an
- [25] obviously incorrect procedure.

- [4] 36 T-shirts and 12 caps, and appropriate work is shown, such as an appropriate system of equations or a correct trial-and-error method with at least two trials and appropriate checks.
- [3] Appropriate work is shown, but only the correct number of T-shirts or the correct number of caps is determined.
- or [3] One error is made, resulting in an incorrect number of T-shirts or caps, but the corresponding number of the other item is determined appropriately.
- [2] An appropriate method is shown, but no answer is found.
- or [2] The variables are represented correctly, and a correct equation or system of equations is written, but the process is not completed. or [2] 36 T-shirts and 12 caps, but only one trial and appropriate checks are shown. or [2] The variables are represented correctly, but an incorrect equation is written, but the solution is completed appropriately.
- [1] 36 T-shirts and 12 caps, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously [26] incorrect procedure.

- [4] 210, and appropriate work is shown, such as a system of equations or the linear equation 5x + 2(295 - x) = 1,220.
- [3] Appropriate work is shown, but one computational error is made.
- or [3] Appropriate work is shown, but the number of children's tickets is found as the answer.
- [2] Appropriate work is shown, but two or more computational errors are made.
- or [2] Appropriate work is shown, but one conceptual error is made.
- or [2] An incorrect equation of equal difficulty is solved appropriately.
- or [2] 210, but a method other than an algebraic solution is used.
- [1] Appropriate work is shown, but one conceptual error and one computational error are made.
- or [1] The correct system of equations or linear equation is written, but no further correct work is shown.
- or [1] 210, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [27] incorrect procedure.
 - [2] 14 and 42, and appropriate work is shown, such as x + 3x = 56, a table, or trial and error with at least three trials and appropriate checks.
 - [1] Appropriate work is shown, but one computational error is made.
 - or [1] Appropriate work is shown, but only one of the two lengths is found.
 - or [1] A correct equation is written and solved, but the lengths are not stated.
 - or [1] An incorrect equation of equal difficulty is solved appropriately.
 - or [1] 14 and 42, but no work or fewer than three trials with appropriate checks are shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [28] incorrect procedure.

- [4] 17 nickels and 15 dimes, and appropriate work is shown, such as the equation 0.05x + 0.10(32 x) = 2.35 or trial and error with at least three trials and appropriate checks.
- [3] Appropriate work is shown, but one computational error is made.
- or [3] Appropriate work is shown, and the correct answers are found, but they are not labeled or are labeled incorrectly.
- or [3] Appropriate work is shown, but only the correct number of nickels or the correct number of dimes is found and labeled.
- [2] Appropriate work is shown, but two or more computational errors are made.
- or [2] Appropriate work is shown, but one conceptual error is made.
- or [2] The trial-and-error method is used to find the correct solution, but only two trials and appropriate checks are shown.
- or [2] The trial-and-error method is attempted and at least six systematic trials and appropriate checks are shown, but no solution is found.
- or [2] An incorrect system of equations of equal difficulty is solved appropriately for both the number of nickels and dimes.
- or [2] A correct equation is solved for *x*, but no further correct work is shown.
- [1] Appropriate work is shown, but one conceptual error and one computational error are made.
- or [1] A correct equation is written, but no further correct work is shown.
- or [1] 17 nickels and 15 dimes, but no work or only one trial with an appropriate check is shown.
- [0] 17 nickels or 15 dimes, but no work or only one trial with an appropriate check is shown.
- or [0] 17 and 15, but no work is shown, and the answers are not labeled or are labeled incorrectly.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an [29] obviously incorrect procedure.

- [31] C
- [32] D
 - [3] \$16,400, and appropriate work is shown, such as
 - 200 tickets sold at the door \$32 = \$6,400 400 tickets sold in advance \$25 = \$10,000 \$16,400
 - [2] The correct number of tickets is shown, but one computational error is made in computing the total amount of money collected.
 - or [2] \$6,400 and \$10,000 are calculated correctly, but they are not added to obtain the total.
 - [1] The numbers of tickets, 200 and 400, are calculated correctly.
 - or [1] An appropriate solution is found, but it is based on incorrect numbers of tickets.
 - or [1] \$16,400, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [33] incorrect procedure.
 - [4] \$6.15, and appropriate work is shown, such as solving simultaneous equations or using a trial-and-error method.
 - [3] \$2.95 (movie) and \$3.20 (game) are found, but they are not added.
 - or [3] Appropriate work is shown, but one computational error is made.
 - [2] The system of equations is set up correctly, but one conceptual error leads to an appropriate solution.
 - or [2] \$2.95 (movie) or \$3.20 (game), and appropriate work is shown.
 - [1] \$6.15, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [34] incorrect procedure.

- [2] \$6.85, and appropriate work is shown.
- [1] The correct rate for the first 5 minutes and the correct rate for each additional minute is shown, but the cost of a 30-minute call is not found.
- or [1] Appropriate work is shown, but one computational error is made.
- or [1] \$6.85, but no work is shown.
- [0] The student calculates either the rate for the first 5 minutes or the rate for each additional minute, but no further work is shown.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an
- obviously incorrect procedure. [35]

a [1] \$50

b(1)[1]5

(2) [1] \$125

c [1] \$10

a, b, and c [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an [36] obviously incorrect procedure.

a [1] c = 10m + 100 for Ron's Rental and c =20m + 20 for Josie's Rental.

- b [2] Two lines, rays, or segments are graphed and labeled correctly, using values arrived at by using a table or by using the slope and yintercept.
- [1] Two lines, rays, or segments are graphed correctly, but they are not labeled.
- or [1] One line, ray, or segment is graphed and labeled correctly, using values arrived at by using a table or by using the slope and yintercept.

c [1] 8

[37]

or [1] An appropriate number of months is found, based on an incorrect graph in part b. a, b, and c [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.

[4] y = 2x - 40, a correctly drawn graph with a slope of 2 and a y-intercept of -40, and 20, and appropriate work is shown.

[3] Appropriate work is shown, but one computational or graphing error is made.

- or [3] The equation and graph are correct, but the breakeven point is missing or is incorrect.
- [2] Appropriate work is shown, but more than one computational or graphing error is made.
- or [2] An incorrect equation is written, but an appropriate graph is drawn, and an appropriate breakeven point is identified.
- [1] An incorrect equation is written, but an appropriate graph is drawn, but the breakeven point is missing or is incorrect.
- or [1] A correct equation is written, but the graph is incorrect, and the breakeven point is not identified.
- or [1] y = 2x 40 and 20, but no work is shown and no graph is drawn.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [38] incorrect procedure.
 - [3] 5, and appropriate work is shown, such as the equation 60 + 5x = 135 - 10x, or trial and error with at least three trials and appropriate checks, or a graph.
 - [2] Appropriate work is shown, but one computational or graphing error is made. or [2] The trial-and-error method is used to find a correct solution, but only two trials and appropriate checks are shown.
 - [1] Appropriate work is shown, but more than one computational or graphing error is made. or [1] 5, but no work or only one trial with an appropriate check is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously [39] incorrect procedure.

- [3] 20, and appropriate work is shown, such as an equation, trial and error, or a graph. [2] Appropriate work is shown, such as 12.95 +0.25x = 14.95 + 0.15x, but one computational error is made. or [2] Appropriate work is shown, but an answer of \$17.95 is found. or [2] 20, and only a check is shown. [1] The student starts appropriate work to find when the prices are equal but does not complete it, such as starting to solve the correct equation, showing one incorrect trial, or drawing an incomplete graph. or [1] 20, but no work is shown. [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct
- response that was obtained by an obviously [40] incorrect procedure. [3] 345, and appropriate work is shown, such
 - as solving the inequality 1450x > 500,000, solving an equation, or trial and error with at least three trials and appropriate checks.
 - [2] Appropriate work is shown, but one computational or rounding error is made. or [2] The trial-and-error method is used to find a correct solution, but only two trials and appropriate checks are shown.
 - [1] Appropriate work is shown, but two or more computational or rounding errors are made.
 - or [1] Appropriate work is shown, but one conceptual error is made.
 - or [1] A correct inequality or equation is written, but no further correct work is shown. or [1] The trial-and-error method is attempted and at least six systematic trials and appropriate checks are shown, but no solution is found.
 - or [1] 345, but no work or only one trial with an appropriate check is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [41] incorrect procedure.
- [42] B

- a [2] 3 and an appropriate method is shown, such as trial and error or the equation 32 + 8x = 26 + 10x.
- [1] 3 and no work is shown.
- or [1] An appropriate method is shown, but an incorrect answer is found.
- b (1) [1] Best Cable Company and an appropriate explanation is given.
- b (2) [1] \$24 and an appropriate explanation is given.
- b (1) and b (2) [1] Best Cable Company and \$24 and no work is shown.
- a and b [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an
- [43] obviously incorrect procedure.
- [44] A
 - [2] More than 6 hours, and appropriate work is shown, using a graphic or algebraic solution.
 - [1] Appropriate work is shown, but one computational error or an error in analyzing the results is made.
 - or [1] More than 6 hours, but no work is
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [45] incorrect procedure.
- [46] B
- [47] B
- [48] B

- [4] Both inequalities are graphed correctly and at least one is labeled, and the solution set is labeled S.
- [3] Appropriate work is shown, but one graphing error is made, such as drawing a solid line for y > x - 4 or shading incorrectly, but the solution set is labeled *S*.
- or [3] Both inequalities are graphed correctly and at least one is labeled, but the solution set is not labeled or is labeled incorrectly.
- or [3] Both inequalities are graphed correctly, the solution set is labeled, but neither inequality is labeled.
- [2] Appropriate work is shown, but two or more graphing errors are made, but an appropriate solution set is labeled.
- or [2] Appropriate work is shown, but one conceptual error is made, such as graphing the lines y = -x + 2 and y = x - 4 and labeling the point of intersection S.
- [1] One inequality is graphed and shaded correctly, but no further correct work is shown.
- or [1] The lines y = -x + 2 and y = x 4 are graphed correctly, but no solution is indicated.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously [49] incorrect procedure.

- [4] The inequalities $x \le 10$, $y \le 12$, and x + y≤ 16 are graphed and shaded correctly on the given set of axes.
- [3] All inequalities are graphed and shaded correctly, but one incorrect type of line (dashed or broken) is used.
- or [3] All three inequalities are graphed correctly, but one inequality is not shaded or is shaded incorrectly.
- or [3] The inequality $x + y \le 16$ is graphed correctly, but an error is made in graphing either the horizontal or vertical line, but they are shaded appropriately.
- or [3] Only two of the three inequalities are graphed correctly, but all three are shaded appropriately.
- [2] All three inequalities are graphed correctly, but two are shaded incorrectly. or [2] Only two of the three inequalities are graphed and shaded correctly.
- [1] Only one of the three inequalities is graphed and shaded correctly.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously [50] incorrect procedure.

[1]	<u>C</u>
[2]	<u>A</u>
[3]	<u>B</u>
[4]	<u>B</u>
[5]	<u>B</u>
[6]	<u>B</u>
[7]	<u>C</u>
[8]	<u>C</u>
[9]	<u>A</u>
[10]	<u>C</u>
[11]	<u>A</u>
[12]	<u>D</u>
[13]	[2] $3(x + 7)(x - 2)$, and appropriate work is shown. [1] Appropriate work is shown, but one computational error is made. or [1] A conceptual error is made, such as incomplete factoring. or [1] $3(x + 7)(x - 2)$, but no work is shown. [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.
[14]	<u>C</u>
[15]	<u>B</u>
[16]	[2] $5(n + 4)(n - 4)$, and appropriate work is shown. [1] Appropriate work is shown, but one factoring error is made or the expression is not simplified completely. [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.

- [2] 3a(x-3)(x+3), and appropriate work is shown. [1] Appropriate work is shown, but one factoring error is made, or the expression is not factored completely. or [1] 3a(x-3)(x+3), but no work is shown. [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [17] incorrect procedure.

[1]	<u>D</u>		[2] $5rs^2\sqrt{2}$, and appropriate work is shown.
[2]	С		[1] A partially correct answer is found, such
[3]	C		as $5r\sqrt{2}s^4$ or $5s^2\sqrt{2}r^2$, and appropriate work is shown.
[4]	<u>A</u>		or [1] $7.07rs^2$, but appropriate work is shown. or [1] $5rs^2\sqrt{2}$, but no work is shown.
[5]	B [2] $\sqrt{20}$ and an appropriate explanation is		[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct
	given, such as the number cannot be written	[11]	response that was obtained by an obviously incorrect procedure.
	as a repeating or terminating decimal or it cannot be written as a fraction or it is not a	[12]	<u>B</u>
	perfect square. [1] $\sqrt{20}$ and an inappropriate explanation or	[13]	<u>C</u>
	no explanation is given.	[14]	A
	or [1] $\sqrt{20}$ and a correct explanation is given but one other number is also identified as	, [15]	<u>C</u>
	irrational. [0] All three numbers are identified as	[16]	<u>B</u>
	irrational.	[17]	<u>B</u>
	or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a	[18]	<u>B</u>
[6]	correct response that was obtained by an obviously incorrect procedure.	[19]	<u>D</u>
[7]	В	[20]	<u>A</u>
[8]	C	[21]	<u>C</u>
[9]	<u>-</u> В	[22]	<u>B</u>
[~]		[23]	<u>C</u>
	[2] 30√2, and appropriate work is shown.[1] Appropriate work is shown, but one	[24]	<u>A</u>
	computational error is made. or [1] Appropriate work is shown, but one	[25]	D
	conceptual error is made. or [1] Appropriate work is shown, but the answer is not in simplest radical form.	[26]	<u>D</u>
	or [1] $30\sqrt{2}$, but no work is shown.		
	[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously		
[10]	incorrect procedure.		

[1]	<u>D</u>
[2]	<u>C</u>
	[2] 3, and appropriate work is shown, such as factoring or trial and error with at least three trials and appropriate checks. [1] Appropriate work is shown, but one computational error is made. or [1] Appropriate work is shown, but one conceptual error is made, such as not rejecting the negative root. or [1] The trial-and-error method is attempted and at least six systematic trials and appropriate checks are shown, but no solution is found. or [1] 3, but no work or fewer than three trials and appropriate checks are shown. [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
[3]	response that was obtained by an obviously incorrect procedure.
[4]	<u>A</u>
[5]	<u>B</u>
[6]	<u>A</u>
[7]	<u>D</u>
[8]	<u>D</u>
[9]	<u>C</u>
10]	<u>A</u>
11]	<u>D</u>

- [3] -8 and 5 and appropriate work is shown, such as factoring or trial and error.
- [2] The student shows correct factoring into (x + 8)(x 5) or correct use of the quadratic formula but finds only one correct value for x.
- [1] Correct factoring is shown, but no values are found.

or

[1] Incorrect factoring is shown, but two appropriate values are found.

OI

[1] Either –8 or 5 is arrived at by trial and error.

or

- [1] –8 and 5 and no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [12] incorrect procedure.
 - [3] –7 and 4, and appropriate work is shown, such as factoring.
 - [2] Correct factoring (x + 7)(x 4) is shown, but only one correct value of x is found. or [2] Correct factoring is shown, but the negative value of x is rejected.
 - [1] Correct factoring is shown, but the values of *x* are not found.
 - or [1] Incorrect factoring is shown, but appropriate values are found.
 - or [1] Only one value is found by trial and error.
 - or [1] –7 and 4, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [13] incorrect procedure.

- [3] -6 and 4, and appropriate work is shown, such as factoring or trial and error with at least three trials and appropriate checks.

 [2] Appropriate work is shown, but one
- [2] Appropriate work is shown, but one computational error is made.
- or [2] Appropriate work is shown, but only one correct value for x is found.
- or [2] The trial-and-error method is used to find the correct solutions, but only two trials and appropriate checks are shown.
- [1] Appropriate work is shown, but two or more computational errors are made.
- or [1] Appropriate work is shown, but one conceptual error is made.
- or [1] The equation is factored correctly, but no values are found.
- or [1] The equation is factored incorrectly, but two appropriate values are found.
- or [1] -6 and 4, but no work or only one trial with an appropriate check is shown.
- [0] -6 or 4, but no work or only one trial with an appropriate check is shown.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an
- [14] obviously incorrect procedure.
- [15] A
- [16] <u>D</u>
- [17] C
- [18] C
- [19] D
- [20] A
- [21] <u>C</u>
- [22] D

- [3] 5, and appropriate work is shown, such as the quadratic equation (x + 7)(x 3) = 24 or trial and error with at least three trials and appropriate checks.
- [2] A correct quadratic equation is written, but one computational error is made in finding Tamara's age.
- or [2] 12 and 2 are found as the sisters' ages, but Tamara's age is not found.
- or [2] The trial-and-error method is used to find the correct solution, but only two trials and appropriate checks are shown.
- [1] Appropriate work is shown, but two or more computational errors are made.
- or [1] Appropriate work is shown, but one conceptual error is made.
- or [1] A correct quadratic equation is written, but no further correct work is shown.
- or [1] An incorrect equation of equal difficulty is solved appropriately for Tamara's age.
- or [1] The trial-and-error method is attempted and at least six systematic trials and appropriate checks are shown, but no solution is found.
- or [1] 5, but no work or only one trial with an appropriate check is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [23] incorrect procedure.

- [3] 4, 6, and 8, and appropriate work is shown, such as the correct quadratic equation or trial and error with at least three trials and appropriate checks.
- [2] The correct quadratic equation is solved, but one computational error is made, but three appropriate ages are listed.
- or [2] The correct quadratic equation is solved, but the negative root is not rejected, but three appropriate ages are listed.
- or [2] The correct quadratic equation is solved, but only one age is found.
- or [2] The trial-and-error method is used to find a correct solution, but only two trials and appropriate checks are shown.
- [1] An incorrect equation of lesser difficulty is solved appropriately, and the three ages are listed.
- or [1] An incorrect quadratic equation of equal difficulty is solved appropriately, and the three ages are listed.
- or [1] The correct quadratic equation is shown, but more than one computational error is made.
- or [1] The correct quadratic equation is shown, but no further correct work is shown. or [1] 4, 6, and 8, but no work or only one trial with an appropriate check is shown. [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct
- response that was obtained by an obviously [24] incorrect procedure.

- [4] 3, 5, and 7, and appropriate work is shown, such as an appropriate quadratic equation or trial-and-error method.
- [3] An appropriate equation is written and solved, but one computational error is made. or [3] An appropriate equation is written and solved, but the even solutions are also listed.
- [2] An incorrect quadratic equation is shown, but it is solved appropriately.
- or [2] Integers are misrepresented, but the subsequent quadratic equation is solved appropriately.
- or [2] An appropriate equation is written and solved, but more than one computational error is made.
- or [2] The correct solution is given, but only one trial is shown with appropriate checks when a trial-and-error method is used.
- [1] A linear equation is solved appropriately. or [1] 3, 5, and 7, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- incorrect procedure. [25]

[26] A

[27] C

- [4] 16 and appropriate work is shown, such as W(W+2) = 15.
- [3] Appropriate work is shown, but one computational error is made.
- or [3] L=5, W=3, and the perimeter = 16, but no work is shown.
- [2] Appropriate work is shown, but more than one computational error is made.
- or [2] L = 5, W = 3, and appropriate work is shown, but the perimeter is not found.
- or [2] The length and width are incorrect, but the perimeter is computed appropriately.
- [1] Length and width are appropriately defined in terms of a single variable.
- or [1] 16 but no work is shown.
- [0] L=5 and W=3 but no work is shown, and the perimeter is not found.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.
- [28]

- [4] Width = 20 and length = 25 and an appropriate algebraic equation is shown, such as $x^2 + 5x - 500 = 0$.
- [3] A correct quadratic equation is shown, but one error is made.
- or [3] A correct quadratic equation is shown, but solved for only one dimension.
- [2] An appropriate solution is shown, but the student fails to reject the negative root and finds two sets of dimensions.
- or [2] The quadratic equation (5x)(x) = 500 is solved appropriately for both dimensions, x = 10 and 5x = 50.
- [1] The student writes only the correct quadratic equation or only the equation x(x+5) = 500 or fails to solve the equation correctly.
- or [1] The student writes a linear equation from x(x+5) = 500, such as 2x + 5x = 500, but solves that equation appropriately. or [1] A correct equation is shown for the perimeter and solved appropriately.
- or [1] (5x)(x) = 500 is solved correctly for only one dimension.
- or [1] 20 and 25 but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [29] incorrect procedure.

- [4] x(x+10) = 144 or an equivalent equation and 8 = width and 18 = length, and appropriate work is shown.
- [3] Appropriate work is shown, but one computational error is made.
- or [3] A correct equation is used and a correct solution is found, but only one dimension is identified.
- [2] An appropriate solution is found to an incorrect equation of equal difficulty. or [2] A correct equation set equal to zero is shown, with no further work or incorrect work
- [1] A conceptual error is made, such as writing the equation 2x+2(x+10) = 144, but the dimensions are found appropriately. or [1] x(x+10) = 144 and 8 = width and 18 = length, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously [30] incorrect procedure.

- [4] w(w+3) = 40, width = 5, and length = 8, and appropriate work is shown.
- [3] w(w+3) = 40 and appropriate work is shown, but one computational error is made in finding the length and width.
- or [3] w(w+3) = 40 and appropriate work is shown, but only the width is found.
- [2] w(w+3) = 40 and appropriate work is shown, but the length and width are not identified.
- or [2] w(w+3) = 40 and appropriate work is shown, but more than one computational error is made in finding the length and width. or [2] An incorrect equation of equal difficulty is solved appropriately for the length and width.
- [1] w(w+3) = 40, but no further correct work is shown.
- or [1] Appropriate work is shown, but one conceptual error is made, such as solving the equation 2w+2w+6=40.
- or [1] w(w+3) = 40, width = 5, and length = 8, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [31] incorrect procedure.
- [32] <u>C</u>
- [33] <u>A</u>
- [34] C
- [35] C

- *a* [3] A parabola is correctly graphed through (0,0), (1,5), (2,8), (3,9), (4,8), (5,5), and (6,0).
- [2] The correct table of values is shown but is not graphed through the entire domain.
- or [2] The correct points are graphed but as a broken line graph not a curve.
- or [2] At least three values are correctly calculated and graphed.
- [1] At least two of the values are correctly calculated, and the student tried to graph all points.

b [1] 3

- or [1] The correct time, *x*, for an incorrect graph in part a is found.
- a and b [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an
 [36] obviously incorrect procedure.

- [4] 100 and a correct parabolic arch is drawn, and appropriate work is shown, such as a table of values for the parabola or correctly labeled points.
- [3] 100 and a correct parabolic arch is drawn, but no table of values or labeled points are shown
- or [3] 100 and a correct parabolic arch is drawn, and appropriate work is shown, but no scale or an incorrect scale is shown.
- or [3] A correct parabolic arch is drawn, but the maximum height is missing or is incorrect.
- [2] An incorrect parabolic arch is drawn, but an appropriate maximum height is found. or [2] A correct height is determined algebraically, but a parabolic arch is not drawn.
- or [2] 100 and an appropriate parabolic arch is drawn, but it is not drawn between $0 \le x \le 20$.
- [1] A correct parabolic arch is drawn, but no work is shown, such as a table of values or correctly labeled points, and the maximum height is missing or is incorrect.
- or [1] 100, but no work is shown and no parabolic arch is drawn.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [37] incorrect procedure.

- a [3] A parabola with points graphed at (0,0), (1,32), (2,48), (3,48), (4,32), and (5,0) is shown. [Points do not have to be labeled on the graph for full credit.]
- [2] Appropriate work is shown, such as a table of values, but one graphing error is made.
- or [2] The correct points are graphed, but the parabola is drawn incorrectly, such as connecting (2,48) and (3,48) as a line segment or not connecting the points at all.
- or [2] At least four correct values are found, and the parabola is graphed appropriately. or [2] A correct table of values is shown for
- all values from 0 to 5, but no graph is drawn. [1] Two or three correct values are found, and the parabola is graphed appropriately.
- or [1] A correct table of values is shown for an incorrectly transcribed equation, such as
- $h = 8t^2 + 40t$, but no graph is drawn.
- b [1] 2.5 is found algebraically or identified from a table or from the graph of the parabola.
- or [1] An appropriate value of t is found, based on an incorrect graph.
- or [1] 2 < t < 3 is given as the range of values based on the line segment drawn in part a. a and b [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an [38] obviously incorrect procedure.

- a [3] A parabola is correctly graphed through (0,0), (1,10), (2,16), (3,18), (4,16), (5,10), and (6,0).
- [2] A correct table of values is shown, but not all the points are graphed correctly.
- or [2] The correct points are graphed but as a broken-line graph, not a curve.
- or [2] At least four values are calculated correctly and graphed.
- [1] The student has at least two of the values calculated correctly and has tried to graph all the points.
- [0] Fewer than two values are calculated correctly.
- b [1] A maximum height of 18 is found. or [1] Correct y is found for an incorrect graph in part a.
- a and b [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an [39] obviously incorrect procedure.
 - [4] A correct graph is drawn, and 3.
 - [3] 3, and appropriate work is shown, but one graphing error is made.
 - or [3] A correct graph is drawn and the points 0.5 and 3.5 are identified, but the difference is not calculated.
 - [2] Appropriate work is shown, but two or more graphing errors are made.
 - or [2] Appropriate work is shown, but one conceptual error is made.
 - or [2] 3, but a method other than a graphic solution is used.
 - [1] Appropriate work is shown, but one conceptual error and one graphing error are made.
 - or [1] A correct graph is sketched with t = 0 to t = 4, but no further correct work is shown. or [1] 3, but no work is shown and no graph is drawn.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure
- [40] incorrect procedure.
- [41] B

- [3] $y = x^2 + 3x 18$, and appropriate work leading from the roots to the equation is shown.
- [2] Appropriate work is shown, but one computational error is made.
- or [2] $x^2 + 3x 18 = 0$, but appropriate work is shown.
- or [2] Only the correct factors (x + 6) and (x 3) are shown.
- [1] Appropriate work is shown, but more than one computational error is made.
- or [1] Only the roots -6 and 3 are shown, such as x = -6, x = 3.
- or [1] $y = x^2 + 3x 18$, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [42] incorrect procedure.
- [43] D
- [44] C
 - [4] 3, and an appropriate algebraic or graphic solution is shown.
 - [3] The equation is graphed correctly, but the time to reach the ground is not identified.
 - or [3] Appropriate work is shown for an algebraic solution, but either no solution is found or the negative root is not rejected.
 - or [3] An appropriate algebraic solution is shown, but one computational error is made.
 - [2] The equation is graphed incorrectly, but an appropriate time to reach the ground is identified.
 - or [2] The equation is factored incorrectly, but an appropriate solution is found.
 - [1] 3, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [45] incorrect procedure.

- [4] A correct graph is sketched and 1.25, and appropriate work is shown.
- [3] A correct graph is sketched, but one computational or rounding error is made in determining the time.
- or [3] Appropriate work is shown, but one error is made in sketching the graph, such as the axes are not labeled or are labeled incorrectly, but the time is determined correctly.
- or [3] A correct graph is sketched and appropriate work is shown to calculate the time, but the negative root is not rejected.
- [2] Appropriate work is shown, but two or more computational, rounding, or graphing errors are made.
- or [2] Appropriate work is shown, but one conceptual error is made.
- or [2] A correct graph is sketched, but no further correct work is shown.
- or [2] Appropriate work is shown to calculate the time, but no graph or an incorrect graph is sketched.
- [1] Appropriate work is shown to calculate the time, but one computational or rounding error is made, and no graph or an incorrect graph is sketched.
- or [1] Appropriate work is shown to calculate the time, but the negative root is not rejected, and no graph or an incorrect graph is sketched.
- or [1] 1.25, but no graph or an incorrect graph is sketched, and no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [46] incorrect procedure.
- [47] D

- [4] Correct graphs are drawn, and (0,5) and (4,-3) are stated.
- [3] Both equations are graphed, but one graphing error is made, but appropriate solutions are stated.
- or [3] Both graphs are drawn correctly, but only one solution is stated.
- [2] Both graphs are drawn correctly, but no solutions are stated.
- or [2] Both equations are graphed, but two or more graphing errors are made, but appropriate solutions are stated.
- or [2] Appropriate work is shown to find (0,5) and (4,-3), but a method other than graphing is used.
- or [2] Both equations are graphed, but one conceptual error is made.
- [1] Both equations are graphed, but one conceptual error and one graphing error are made.
- or [1] (0,5) and (4,-3) are stated, but no work is shown.
- [0] (0,5) or (4,-3) is stated, but no work is shown.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.

[48]

- [4] (10,0) and (1,9), and both graphs are drawn correctly.
- [3] Both graphs are drawn correctly, but only one solution is stated correctly.
- or [3] One graph of equal difficulty is drawn incorrectly, but the solutions are appropriate, based on the graphs.
- [2] (10,0) and (1,9), but the problem is solved algebraically instead of graphically.
- or [2] One graph of equal difficulty is drawn incorrectly, and only one solution is appropriate, based on the graphs.
- [1] Both the parabola and the line are graphed incorrectly, but the solutions are appropriate, based on the graphs.
- or [1] Incorrect solutions result from an algebraic method.
- or [1] (10,0) and (1,9), but no work is shown. [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [49] incorrect procedure.

- [4] 9 and 26, and appropriate work is shown, such as graphing and labeling the equations and identifying the points of intersection.
- [3] Both functions are graphed correctly, and the points of intersection are indicated, but the prices are not stated.
- or [3] The parabola is graphed correctly, but the line is graphed incorrectly, but appropriate prices are stated.
- [2] The line and the parabola are graphed and labeled, but a conceptual error is made, such as only one price is found because the graph of the parabola is incomplete.
- or [2] The line is graphed correctly, but the parabola is graphed incorrectly, but appropriate prices are stated.
- or [2] 9 and 26, but only an algebraic solution is shown.
- [1] Both the line and the parabola are graphed incorrectly, but appropriate prices are stated. or [1] 9 and 26, but no work is shown.
- [0] 9 or 26, but no work is shown. or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.
- [50] Obviously incorrect

[51] D

- [4] (-3,-5) and (1,3), and appropriate algebraic work is shown.
- [3] Appropriate algebraic work is shown, but x = -3 and x = 1 are given as the solution. or [3] Appropriate algebraic work is shown, but only one correct solution is given, such as (1,3).
- [2] (-3,-5) and (1,3), but a graphic solution is shown.
- or [2] Correct substitution and an algebraic equation set equal to zero are shown, but the result is not factored, such as $x^2 + 2x 3 = 0$. [1] Any correct substitution is shown, such as $2x + 1 = x^2 + 3x 2$.
- or [1] (-3,-5) and (1,3), but no algebraic work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [52] incorrect procedure.

- [4] (-1,-2) and (2,13), and appropriate work is shown, such as an algebraic or graphic solution or trial and error with at least three trials and appropriate checks.
- [3] Appropriate work is shown, but one computational or graphing error is made. or [3] Appropriate work is shown, but only one solution is found or only the x- or the y-values are found.
- [2] Appropriate work is shown, but two or more computational or graphing errors are made.
- or [2] Appropriate work is shown, but one conceptual error is made.
- or [2] The trial-and-error method is used to find the correct solutions, but only two trials and appropriate checks are shown.
- or [2] The trial-and-error method is attempted and at least six systematic trials and appropriate checks are shown, but no solution is found.
- or [2] Both equations are graphed correctly, but neither ordered pair is identified.
- or [2] Only one equation is graphed correctly, but an appropriate solution is found.
- or [2] An incorrect quadratic equation of equal difficulty is solved appropriately, and appropriate solutions are found.
- [1] Appropriate work is shown, but one conceptual error and one computational or graphing error are made.
- or [1] One equation is graphed correctly, but no further correct work is shown.
- or [1] An incorrect equation of a lesser degree of difficulty, such as a linear equation, is solved appropriately.
- or [1] A correct substitution is made and the system of equations is simplified to a single quadratic equation set equal to zero, but no further correct work is shown.
- or [1] (-1,-2) and (2,13), but no work or only one trial with an appropriate check is shown. [0] (-1,-2) or (2,13), but no work or only one trial with an appropriate check is shown. or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.

- [4] (3,14) and (-2,-1) and either an algebraic or a graphic solution is shown.
- [3] An appropriate method is shown, but only one correct ordered pair is identified.
- or [3] An appropriate method is shown, but one computational mistake is made.
- or [3] An appropriate method is shown, but values are given only for x.
- [2] The substitution is correct, but the quadratic produced is not factored correctly. or [2] Both equations are graphed correctly, but neither ordered pair is identified.
- [1] Only one equation is graphed correctly. or [1] The substitution is incorrect, but it produces a linear equation that is solved correctly.
- or [1] Only the substitution is correct.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.
 - [4] Yes, and appropriate work is shown, and an appropriate justification is given.
 - [3] Appropriate work is shown, and an appropriate justification is given, but one computational error is made, or the negative value of t is not rejected.
 - [2] An appropriate graph or equation is shown, such as $16t^2 8t 15 = 0$.
 - [1] An incorrect graph or equation of equal difficulty is used, but an appropriate solution is found.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [55] incorrect procedure.

- [1] B
- [2] C
- [3] C
- [4] B
- [5] A
- [6] C
- [7] C
- [8] D
- [9] A
- [10] B

$$[2] \ \frac{3x}{3x+5y}$$

- [1] One correct factoring is shown, either 3x(3x 5y) or (3x 5y)(3x + 5y).
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [11] incorrect procedure.
 - [2] $\frac{x+1}{x-5}$, and appropriate work is shown.
 - [1] Only one expression is factored correctly, such as (x + 5)(x + 1) or (x + 5)(x 5), but an appropriate simplification is done.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [12] <u>incorrect procedure.</u>
- [13] C
- [14] <u>A</u>

- [2] $\frac{-x-3}{10x+2}$ or an equivalent answer in
- simplest form, and appropriate work is shown.
- [1] Either the numerator or the denominator is factored completely.
- or [1] Appropriate work is shown, but

$$\frac{3-x}{x-3} = -1$$
 is not recognized.

or [1]
$$\frac{-x-3}{10x+2}$$
 or an equivalent answer in

simplest form, but no work is shown.

- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [15] incorrect procedure.
- [16] B
- [17] D
- [18] D
- [19] C
- [20] B
 - [2] 4(x-2) or 4x-8, and appropriate work is shown
 - [1] The problem is factored correctly but not reduced to simplest form.
 - or [1] Only two of the expressions are factored correctly, but an appropriate answer is found.
 - or [1] 4(x-2) or 4x-8, but no work is shown.
 - [0] Only the formula for volume is shown. or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an
- [21] obviously incorrect procedure.

[2]
$$\frac{x-3}{3}$$
 and multiplication by the reciprocal,

correct factoring, and canceling are shown.

[1] The difference of two squares,

$$x^2 - 9 = (x+3)(x-3)$$
, is factored correctly.

or [1] Appropriate work is shown, but the final answer is incorrect.

or [1]
$$\frac{x-3}{3}$$
 but no work is shown.

[22]

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.

[4] $\frac{x+3}{2}$, and appropriate work is shown.

[3] Appropriate work is shown, but one computational, factoring, or simplification error is made.

[2] Appropriate work is shown, but two or more computational, factoring, or simplification errors are made.

or [2] Appropriate work is shown, but one conceptual error is made, such as failing to multiply by the reciprocal of g(x) or trying to solve for x.

[1] Appropriate work is shown, but one conceptual error and one computational, factoring, or simplification error are made.

or [1]
$$\frac{x+3}{2}$$
, but no work is shown.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.

[23] <u>incorrect procedure.</u>

[6]
$$-\frac{8}{3}$$
, and appropriate work is shown.

[5] Appropriate work is shown, but one computational error is made.

[4] Appropriate work is shown, but two or more computational errors are made.

[3] Appropriate work is shown, but one conceptual error is made, such as not factoring out -1 when canceling out 2 - x.

[2] Appropriate work is shown, but one conceptual error and one computational error are made.

 $[1] - \frac{8}{3}$, but no work is shown.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously [24] incorrect procedure.

[6] -(x-3), -x+3, or 3-x, and appropriate work is shown.

[5] Appropriate work is shown, but one computational, factoring, or simplification error is made.

[4] Appropriate work is shown, but two computational, factoring, or simplification errors are made.

or [4] x-3, and appropriate work is shown.

[3] Appropriate work is shown, but three or more computational, factoring, or simplification errors are made.

or [3] Appropriate work is shown, but one conceptual error is made, such as not multiplying by the multiplicative inverse.

[2] Appropriate work is shown, but one conceptual error and one computational, factoring, or simplification error are made.

[1] -(x-3), -x+3, or 3-x, but no work is shown.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously

[25] incorrect procedure.

[26] C

[27] <u>A</u>

- [28] A
- [29] C
- [30] A
- [31] B
- [32] D
 - [2] $\frac{2x+3}{x(x+3)}$ or $\frac{2x+3}{x^2+3x}$, and appropriate

work is shown.

- [1] Appropriate work is shown, but one computational error is made or the answer is not simplified completely.
- or [1] Appropriate work is shown, but one conceptual error is made.

or [1]
$$\frac{2x+3}{x(x+3)}$$
 or $\frac{2x+3}{x^2+3x}$, but no work is

shown.

- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [33] incorrect procedure.
- [34] D
- [35] D

- [6] $\frac{x-4}{2}$, and appropriate work is shown.
- [5] Appropriate work is shown, but one computational error is made.
- [4] Appropriate work is shown, but two or more computational errors are made. or [4] Appropriate work is shown, but −1 is not factored out.
- [3] Appropriate work is shown, but one conceptual error is made, such as not following the correct order of operations.
- [2] Appropriate work is shown, but one conceptual error and one computational error are made.
- [1] Appropriate work is shown, but one conceptual error and two or more computational errors are made. or [1] Appropriate work is shown, but two
- or [1] $\frac{x-4}{2}$, but no work is shown.

conceptual errors are made.

- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [36] incorrect procedure.
 - [3] 4, and appropriate work is shown.
 - [2] Appropriate work is shown, but one computational error is made.
 - [1] Appropriate work is shown, but two or more computational errors are made.
 - or [1] Appropriate work is shown, but one conceptual error is made.
 - or [1] 4, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [37] incorrect procedure.
- [38] C
- [39] D
- [40] <u>A</u>
- [41] A
- [42] A

[43]	D	

[44] B

[2] 2, and appropriate work is shown.

[1] Appropriate work is shown, but one computational error or one conceptual error is made.

or [1] 2, but no work is shown.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously

[45] incorrect procedure.

[2] 2.4 and appropriate work is shown.

[1] The student shows correct use of the distributive property to obtain 2x - 6 or other appropriate algebraic technique.

or [1] 2.4 and no work is shown.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously

[46] incorrect procedure.

[2] 2.1, and appropriate work is shown.

[1] Appropriate work is shown, but one computational error is made.

or [1] Appropriate work is shown, but one conceptual error is made.

or [1] 2.1, but no work is shown.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect precedure.

[47] incorrect procedure.

[2] 38 and an appropriate method is shown, such as 36.64 - (21 + 6.14) = 9.50 and $\frac{9.50}{25}$ =

38 or an equation such as 21 + .25c + 6.14 = 36.64.

[1] 38 and no work is shown.

or [1] An appropriate method or equation is shown, but one computational mistake is made.

or [1] The answer of \$9.50 for local calls is found but is not divided by .25.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously

[48] incorrect procedure.

[49] B

[50] D

[2] 5 and appropriate work is shown, such as substituting \$18.11 for *p* and solving the equation correctly, or trial and error with at least three trials and appropriate checks.

[1] Appropriate work is shown, but one computational error is made.

or [1] Appropriate work is shown, but one conceptual error is made.

or [1] 5, but no work or fewer than three trials with appropriate checks are shown.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously

[51] incorrect procedure.

[52] A

[53] C

[54] B

- [2] 1 and -2, and appropriate work is shown.
- [1] Appropriate work is shown, but one computational error is made.
- or [1] Appropriate work is shown, but one conceptual error is made.
- or [1] Appropriate work is shown, but only one value is found.
- or [1] 1 and -2, but no work is shown.
- [0] 1 or -2, but no work is shown.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an [55] obviously incorrect procedure.
 - [4] 3 and -5, and appropriate work is shown, such as x(x + 7) = 5(x + 3) or trial and error with at least three trials and appropriate checks for each solution.
 - [3] Appropriate work is shown, but one computational or factoring error is made.
 - or [3] Appropriate work is shown, but only one correct solution is found.
 - or [3] The trial-and-error method is used to find both correct solutions, but only two trials and appropriate checks are shown for each solution.
 - [2] Appropriate work is shown, but two or more computational or factoring errors are made.
 - or [2] A correct quadratic equation is written and factored, but no further correct work is shown.
 - or [2] The trial-and-error method is attempted and at least six systematic trials and appropriate checks are shown, but neither solution is found.
 - [1] A correct quadratic equation is written, but no further correct work is shown.
 - or [1] 3 and -5, but no work or only one trial with an appropriate check is shown.
 - [0] 3 or -5, but no work or only one trial with an appropriate check is shown.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an [56] obviously incorrect procedure.

- [4] 2 and -3, and a correct quadratic equation is shown, such as x(x + 1) = 6, and solved algebraically.
- [3] The student shows a correct quadratic equation but makes one algebraic error and carries it to solution or no solution for the equation generated.
- or [3] Correct work is shown, but only one root is found as the answer.
- [2] A correct quadratic equation is used, but two or more errors are made.
- or [2] An incorrect quadratic equation of equal difficulty is shown and solved appropriately.
- [1] The student cross multiplies but produces only a linear equation that is solved appropriately.
- or [1] 2 and -3, but no algebraic work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [57] incorrect procedure.

- [4] 6 and –2, and appropriate work is shown, such as an algebraic solution or trial and error with at least three trials and appropriate checks.
- [3] Appropriate work is shown, but one computational or factoring error is made. or [3] Appropriate work is shown, but only one solution is found.
- [2] The correct quadratic equation is written in standard form.
- or [2] Appropriate work is shown, but two or more computational or factoring errors are made.
- or [2] Appropriate work is shown, but one conceptual error is made.
- or [2] The trial-and-error method is used to find at least one solution, but only two trials and appropriate checks are shown.
- or [2] The trial-and-error method is attempted and at least six systematic trials and appropriate checks are shown, but no solution is found.
- or [2] An incorrect quadratic equation of equal difficulty is solved appropriately.
- [1] $x^2 11x 12 = -7x$, but no further correct work is shown.
- or [1] 6 and -2, but no work or only one trial with an appropriate check is shown.
- or [1] An incorrect equation of a lesser degree of difficulty is solved appropriately.
- or [1] Appropriate work is shown, but one conceptual error and one computational or factoring error are made.
- [0] 6 or -2, but no work is shown.
 or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an [58] obviously incorrect procedure.
- [59] B

[2] 1 and 2, 1 < h < 2, or 1 < 1.854 < 2, and appropriate work is shown.

[1]
$$\frac{3}{h} = \frac{h}{3-h}$$
 is shown, but one

computational error is made.

- or [1] The positive root, 1.854, is obtained from the quadratic, but the two correct consecutive integers are not stated.
- or [1] 1 and 2, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [60] incorrect procedure.
- [61] B

[4]
$$\frac{-3\pm\sqrt{37}}{7}$$
 or an equivalent answer, and

appropriate work is shown.

- [3] A correct quadratic equation is written and appropriate work is shown, but one computational or simplification error is made.
- or [3] An incorrect quadratic equation of equal difficulty is solved appropriately.
- [2] A correct quadratic equation is written and appropriate work is shown, but two or more computational or simplification errors are made.
- or [2] Appropriate work is shown, but one conceptual error is made.
- or [2] A correct quadratic equation is written in standard form, but no further correct work is shown.
- [1] An incorrect equation of a lesser degree of difficulty is solved appropriately.

or [1]
$$\frac{-3\pm\sqrt{37}}{7}$$
 or an equivalent answer, but

no work is shown.

- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.
- [62] incorrect procedure.

- [4] 3 and $\frac{1}{2}$, and appropriate work is shown.
- [3] Appropriate work is shown, but one computational error is made.
- or [3] Appropriate work is shown, but only one of the values is found.
- [2] Appropriate work is shown, but two or more computational errors are made.
- or [2] Appropriate work is shown, but one conceptual error is made.
- or [2] The correct quadratic equation is written in standard form, but no further correct work is shown.
- or [2] An incorrect quadratic equation of equal difficulty is solved appropriately.
- [1] Appropriate work is shown, but one conceptual error and one computational error are made.
- or [1] An incorrect equation of a lesser degree of difficulty is solved appropriately.
- or [1] 3 and $\frac{1}{2}$ but no work is shown.
- [0] 3 or $\frac{1}{2}$, but no work is shown.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.
- [2] 8 or an equivalent answer, and appropriate work is shown.
- [1] The denominators are cleared correctly, such as 6(t+16)+6t=t(t+16), but the factoring is incorrect, or one error is made using the quadratic formula.
- or [1] The denominators are not cleared correctly, but an equation of equal difficulty is solved.
- or [1] 8 or an equivalent answer, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [64] incorrect procedure.

[63]

- [6] 3.5, and appropriate work is shown.
- [5] Appropriate work is shown, but one computational or rounding error is made.
- [4] A substitution error is made, resulting in an incorrect quadratic equation of equal difficulty, but the incorrect equation is solved appropriately.
- [3] A correct substitution is made, resulting in the correct quadratic equation in standard form, but the equation is not solved.
- [2] A substitution error is made, resulting in an incorrect equation of equal difficulty, and one computational or rounding error is made.
- [1] 3.5, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect precedure.
- [65] incorrect procedure.
- [66] B

[1]	\boldsymbol{C}	
111		
LŤJ	_	

- [2] C
- [3] B
- [4] A
 - [2] $\frac{1}{2}$ or an equivalent answer, and an

appropriate explanation is written.

[1] A correct explanation is written, but the probability is not stated.

or [1] $\frac{1}{2}$ or an equivalent answer, but no

explanation is written.

- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [5] incorrect procedure.
- [6] A
- [7] C
- [8] A
- [9] D
- [10] A
- [11] D
- [12] C
- [13] D
- [14] C
- [15] A

- [2] 6, and a correct tree diagram is drawn or sample space is listed.
- [1] A correct tree diagram is drawn or sample space is listed, but no answer or an incorrect answer is found.
- or [1] An appropriate answer is found, based on an incorrect tree diagram or sample space.
- or [1] 6, but no tree diagram is drawn or sample space is listed.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [16] incorrect procedure.
 - [2] 12, and a correct tree diagram or a correct sample space is shown.
 - [1] An incomplete tree diagram or sample space is shown with at least 8 possible combinations shown, and an appropriate number of outfits is found.
 - or [1] A correct tree diagram or sample space is shown, but the number of possible outfits is missing or is incorrect.
 - or [1] 12, but 3×4 is used to find the number of outfits.
 - [0] 12, but no work is shown.
 - or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an
- [17] obviously incorrect procedure.
 - a [1] A correct tree diagram or listing of all 8 possibilities is shown.

b [1]
$$\frac{1}{8}$$

- or [1] An appropriate answer is given for an incorrect part a tree diagram or listing. a and b
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [18] incorrect procedure.

- [3] $\frac{4}{9}$, and a correct tree diagram or sample space is shown.
- [2] A correct tree diagram or sample space is shown, but no probability or an incorrect probability is given.
- or [2] An incorrect tree diagram or sample space is shown, but an appropriate probability is found.
- [1] Appropriate work is shown, but one conceptual error is made.
- or [1] $\frac{4}{9}$, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [19] incorrect procedure.
- [20] D
- [21] B
- [22] C
- [23] D
- [24] D
- [25] D
 - [2] 72 and an appropriate method, such as $3\times6\times4$, is shown.
 - [1] 72 and no explanation is given. or [1] An appropriate method is shown, but the student has one computational mistake or an incomplete listing, such as 2 of the 3 clothing categories.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [26] incorrect procedure.
- [27] A
- [28] C
- [29] C
- [30] <u>C</u>

- [31] <u>B</u>
- [32] B
 - [3] 1,344, and appropriate work is shown, such as $8 \cdot 7 \cdot 6 \cdot 4$.
 - [2] Appropriate work is shown, but one computational error is made.
 - [1] Appropriate work is shown, but two or more computational errors are made.
 - or [1] Appropriate work is shown, but one conceptual error is made, such as basing the answer on ordering an appetizer and a soup, using $5 \cdot 3 \cdot 7 \cdot 6 \cdot 4$.
 - or [1] 1,344, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [33] incorrect procedure.
- [34] A
- [35] D
- [36] C
- [37] A
- [38] D

[3]
$$\frac{2}{24}$$
 or an equivalent answer, and an

appropriate explanation is given or appropriate work is shown, such as a tree diagram, sample space, or permutations.

- [2] Appropriate work is shown, but one computational error is made.
- or [2] Appropriate work is shown, but only a numerator or a denominator is determined correctly.

or [2]
$$\frac{2}{24}$$
 or an equivalent answer, but only

work for either the numerator or the denominator is shown.

- [1] The probability of the tallest or the probability of the shortest student being in the proper position is correct, such as .
- or [1] Only a tree diagram, sample space, or permutations are shown.

or [1]
$$\frac{2}{24}$$
 or an equivalent answer, but no

work is shown.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.

[3]
$$\frac{7}{20}$$
 or an equivalent answer, and

appropriate work is shown, such as $\frac{15}{25} \cdot \frac{14}{24}$

or
$$\frac{_{15}C_2}{_{25}C_2}$$
.

[2]
$$\frac{15}{25} \bullet \frac{14}{24}$$
 or $\frac{_{15}C_2}{_{25}C_2}$ is shown, but one

computational error is made or no further work is shown.

- or [2] $_{15}C_2$ and $_{25}C_2$ are computed correctly, but no further work is shown.
- or [2] Appropriate work is shown, but one computational error is made.
- [1] The correct probabilities are found, but they are added instead of multiplied.
- or [1] Only one of the two parts of the probability is correct.
- or [1] Appropriate work is shown, but more than one error is made.

or [1]
$$\frac{7}{20}$$
 or an equivalent answer, but no

work is shown.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously

- [40] incorrect procedure.
- [41] <u>A</u>
- [42] B
- [43] D
- [44] A
- [45] <u>A</u>
- [46] <u>C</u>
- [47] D

- [2] 120, and appropriate work is shown, such as $1 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1$.
- [1] Appropriate work is shown, but one computational error is made.
- or [1] 720 and $_6P_6$ or 6! is shown.
- or [1] 120, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [48] incorrect procedure.
- [49] B
 - [2] 210, and appropriate work is shown, such as $7 \cdot 6 \cdot 5$ or ${}_{7}P_{3}$.
 - [1] Appropriate work is shown, but no answer or an incorrect answer is found.
 - or [1] 210, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [50] incorrect procedure.
- [51] D
- [52] A
 - [2] 8,100 and appropriate work is shown, such as $9 \times 10 \times 10 \times 9$.
 - [1] 10,000 but appropriate work is shown. or [1] Appropriate work is shown, but the student multiplies incorrectly.
 - or [1] An appropriate pattern is shown, such as $9 \times 10 \times 10 \times 9$.
 - or [1] 8,100 but no work is shown.
 - [0] 38 is shown.
 - or [0] The student attempts to use the counting principle, but adds.
 - or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an
- [53] obviously incorrect procedure.

- [4] 7.98×10^6 or 7,980,000 and appropriate work is shown, such as $8 \times 10^6 2 \times 10^4$.
- [3] Appropriate work is shown, but one computational error is made.
- or [3] The student uses 1–9 instead of 0–9 as the number of digits in $8 \times 9^6 2 \times 9^4$.
- [2] The student correctly produces only one part, 8×10^6 or 2×10^4 , but carries the process to an appropriate result.
- or [2] Appropriate work is shown, but more than one error is made.
- [1] The student produces only one part, 8×9^6 or 2×9^4 .
- or [1] 7,980,000 but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [54] incorrect procedure.
 - [2] 37,440 and appropriate work is shown, such as 2 x 26 x 10 x 9 x 8 or $_{2}P_{1}\times_{26}P_{1}\times_{10}P_{3}$.
 - [1] Appropriate work is shown, but one computational or conceptual error is made. or [1] Appropriate work is shown for at least
 - one restriction, such as 2 x 26 or 10 x 9 x 8. or [1] 37,440 but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [55] incorrect procedure.

- [3] Option 2 will yield 82,576,000 more possibilities, and appropriate work is shown, such as $26^3 \cdot 10^4$ and $_{26}P_4 \cdot _{10}P_3$.
- [2] Appropriate work is shown, but one computational error is made, but the appropriate option is identified.
- or [2] The correct numbers of arrangements are found for both Option 1 and Option 2, but the question of which option will yield more arrangements is not answered or is answered incorrectly.
- [1] Appropriate work is shown, but more than one computational error is made, but the appropriate option is identified.
- or [1] Appropriate work is shown, but one conceptual error is made, but the appropriate option is identified.
- or [1] Either Option 1 or Option 2 is found correctly, but no further correct work is
- or [1] Option 2 will yield 82,576,000 more possibilities, but no work is shown.
- [0] Option 2, but no work or inappropriate work is shown.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an
- [56] obviously incorrect procedure.
- [57] B
- [58] B
- [59] D
- [60] A
- [61] A
- [62] A
- [63] C
- [64] C
- [65] B
- [66] B
- [67] B

- [2] 6 and appropriate work is shown, such as using the combination ${}_{4}C_{2}$, listing all six possible outcomes, or drawing a correct tree diagram.
- [1] A correct setup of combinations is shown, but an incorrect solution, such as leaving ${}_{4}C_{2}$, or no integral solution is found.
- or [1] An appropriate list or tree diagram is shown, but an incorrect solution is found, such as 5, by omitting one of the possible combinations.
- or [1] 12 but a complete list or tree diagram is shown.
- or [1] 6 but no work is shown.
- [0] The answer is completely incorrect, such as $_4P_2$ or 4×3 .
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.
- [2] 20,349, and appropriate work is shown, such as $_{21}C_5 = 20,349$.
- [1] Appropriate work is shown, but one computational error is made.
- or [1] Appropriate work is shown, but one conceptual error is made, such as determining the value of $_{21}P_5$.
- or [1] 20,349, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [69] incorrect procedure.

[68]

- [3] 490, and appropriate work is shown, such as 7•7•10.
- [2] Appropriate work is shown, but one computational error is made.
- or [2] Appropriate work is shown, but an incorrect answer is found, based on an incorrect

number of possible dessert combinations or an incorrect number of soup or appetizer choices.

or [2] Appropriate work is shown, but an incorrect answer is found, based on one error in

the tree diagram.

or [2] $\frac{1}{490}$, but appropriate work is shown.

[1] 7, 7, and 10 are added instead of multiplied.

or [1] The counting principle is used correctly, but incorrect substitutions are made, but

an appropriate answer is shown.

or [1] 490, but no work is shown.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure

[70] incorrect procedure.

- [3] 150, and appropriate work is shown, such as ${}_5C_2 {\bullet}_6 C_2$.
- [2] Appropriate work is shown, but one computational error is made.
- or [2] All the possible combinations of two mystery books and all the possible combinations of two biographies are calculated, but the answers are not multiplied.
- [1] Appropriate work is shown, but more than one computational error is made.
- or [1] Appropriate work is shown, but one conceptual error is made, such as the computation $_{11}C_4 = 330$.

or [1] 150, but no work is shown.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously

[71] incorrect procedure.

[72] C

- [2] 10, and appropriate work is shown, such as ${}_{5}C_{2}$ or a diagram or a list.
- [1] Appropriate work is shown, but one computational error is made.
- or [1] Appropriate work is shown, but one conceptual error is made.

or [1] 10, but no work is shown.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously

[73] incorrect procedure.

[74] B

[75] D

a [2] 10 and an appropriate tree diagram, list, sample space, or ${}_{5}C_{3} = 10$ is shown.

[1] 10 and no work is shown.

or [1] An appropriate method is shown, but not all 10 possible combinations are listed

b [1]
$$\frac{1}{10}$$

or [1] An appropriate answer is found for an incorrect response in part a.

c [1]
$$\frac{4}{10}$$
 or $\frac{2}{5}$ or 0.4

or [1] An appropriate answer is found for an incorrect response in part a.

a and b and c [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by

[76] an obviously incorrect procedure.

[4]
$$\frac{4}{10}$$
 and appropriate work is shown, such as the following illustration or any other correct method:

| Salusage | mushrooms | m

mushroom
anchovies

sausage
mushroom
anchovies

mushroom
anchovies

mushroom
anchovies

[3] Appropriate work is shown, but one computational error is made.

or [3] Appropriate work and complement $\frac{6}{10}$ are shown.

or [3] Appropriate work is shown, but the answer is incomplete.

[2] ${}_5C_2$ and the work is appropriate but incomplete.

or [2] 10 but appropriate work is shown.

or [2] A correct sample space or tree diagram is shown.

[1] Incorrect work leading to $0 \le$ fraction ≤ 1 or $0 \le$ percent ≤ 100 is shown.

or [1] $\frac{4}{10}$ but no work is shown.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.

[3] $\frac{8}{20}$ or an equivalent answer, and

appropriate work is shown, such as using a tree diagram or writing the equation

$$\frac{6}{20} + \frac{2}{20} = \frac{8}{20}$$

[2] One computational error is made in

finding $\frac{6}{20}$ or $\frac{2}{20}$, but an appropriate sum is

found.

or [2] $\frac{2}{20}$ and $\frac{6}{20}$ are found, but no sum is shown.

[1] $\frac{6}{20}$ or $\frac{2}{20}$, and appropriate work is

or [1] An appropriate answer is found, using replacement with a tree diagram or an

equation such as
$$\frac{3}{5} \cdot \frac{3}{5} + \frac{2}{5} \cdot \frac{2}{5} = \frac{13}{25}$$
.

or [1]
$$\frac{8}{20}$$
, but no work is shown.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously

[78] incorrect procedure.

[4] No, and an appropriate explanation is given, such as $P(15) = \frac{6}{56} < P(2) = \frac{12}{56}$.

[3] One of the two probabilities is found correctly, but one computational error is made in finding the other, but an appropriate conclusion is drawn, based on the values found.

or [3] Replacement is used to conclude

$$P(15) = \frac{6}{64} < P(2) = \frac{12}{64}.$$

or [3] The two probabilities are found correctly, but no conclusion or the incorrect conclusion is drawn.

[2] One of the probabilities is found correctly, but one computational error is made in finding the other, and no conclusion or the incorrect conclusion is drawn.

[1] An appropriate method is used, such as a tree diagram or sample space, but the probabilities are not determined or are determined incorrectly.

or [1]
$$P(15) = \frac{6}{56} < P(2) = \frac{12}{56}$$
, but no work

is shown.

[0] No, but no work is shown. or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a

response that was obtained by an obviously [79] incorrect procedure.

a [2] $\frac{30}{72}$ or an equivalent value is found and

an appropriate explanation is given.

[1] An acceptable method is used correctly, such as a tree diagram, sample space, or combinations, but the correct answer is not given.

or [1] Replacement is used, and an answer of $\frac{36}{81}$ or an equivalent is found.

or [1] $\frac{30}{72}$ and no work is shown.

b [2] $\frac{36}{72}$ or an equivalent value is found and

an appropriate explanation is given. or [2] An appropriate probability for an incorrect denominator for part *a* is found.

[1] An appropriate method is shown, but one computational mistake is made.

or [1] Replacement is shown, and the answer $\frac{36}{81}$ or an equivalent is found.

or [1] The student does not take into account both orders and answers $\frac{18}{72}$ or an equivalent.

or [1] $\frac{36}{72}$ and no work is shown.

or a and b

[1] An error in method is made but the erroneous answer is interpreted correctly in either part *a* or *b* or both.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously [80] incorrect procedure.

- [1] D
- [2] D
 - [4] The student draws a histogram, a stemand-leaf plot, or any other acceptable statistical graph, with proper labels and a title.
 - [3] The student makes one or two minor errors, such as a lack of label, title, or connected dots.
 - [2] The student makes several minor errors or one major error, such as not accounting for all 20 scores.
 - [1] The student draws just the beginning of a
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [3] incorrect procedure.
 - [2] A correct stem-and-leaf plot is drawn, including a key.
 - [1] The data are arranged correctly, but incorrect labels are written on the stem-andleaf columns. [Columns do not need to be labeled for a full-credit response, but full credit may not be awarded if the columns are labeled incorrectly.]
 - or [1] The data are listed in the stem-and-leaf plot, but not in ascending order.
 - or [1] One or two of the scores are left out of the stem-and-leaf plot.
 - or [1] Duplicate values are left out of the stem-and-leaf plot.
 - [0] Incorrect labels are written on the stemand-leaf columns, and scores are left out of the plot.
 - or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an
- obviously incorrect procedure.

- [3] The frequency table is completed correctly, showing frequencies of 6, 2, 4, 5, and 3, and a frequency histogram is drawn and labeled correctly.
- [2] The frequency table is completed correctly, but one graphing error is made, such as not labeling the axes, having nonequal intervals, or starting the x-axis at 50. or [2] The frequency table is completed incorrectly, but an appropriate frequency histogram is drawn.
- or [2] The frequency histogram is drawn and labeled correctly, but the frequency table is not completed.
- [1] The frequency table is completed correctly, but two or more graphing errors are made.
- or [1] The frequency table is completed correctly, but no frequency histogram is drawn or a bar graph is drawn.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [5] incorrect procedure.

- [4] A correct table and histogram with appropriate labels and scales are shown, such as
- the table below.

SCORE	TALLY	FREQUENCY
40-49	/	1
50-59	/	1
60-69	///	3
70-79	///	3
80-89	///	3

- [3] An incorrect table is shown, but the histogram is appropriate, based on this table. or [3] A correct table is shown, but one error is made on the histogram, such as using incorrect labels or no labels.
- or [3] An incomplete table is shown, but the histogram is correct.
- [2] An incomplete table is shown, and the histogram is partially correct.
- or [2] A correct table is shown, and a correct bar graph is made.
- [1] A correct table is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [6] incorrect procedure.

- *a* [2] An appropriate histogram is drawn with both axes labeled with a correct numerical scale.
- [1] A correct bar graph is drawn.
- or [1] The parts of the histogram are not labeled.
- or [1] Equal interval scales are not shown.
- or [1] One error on frequency calculation is made.
- [0] Two or more mistakes on frequency calculation are made.
- b [2] 60% and an appropriate explanation is given.
- [1] An appropriate method to find percent is shown, but a mistake is made in reading the

chart, such as
$$\frac{6}{15} = 40\%$$
 or $\frac{9}{15}$ is shown but

not given as a percent answer.

- or [1] 60% and no explanation is given. a and b
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [7] incorrect procedure.
 - *a* [3] The frequency table is completed correctly, and a histogram is drawn with a correct scale and is labeled correctly.
 - [2] One or two errors are made in the frequency table, but an appropriate histogram is drawn.
 - or [2] The frequency table is completed correctly, but one error is made in drawing the histogram.
 - [1] A correct histogram is drawn, but the frequency table is not completed.
 - *b* [1] The interval 91-100 is identified as containing the 75th percentile.
 - or [1] The appropriate interval is identified, based on an incorrect frequency table in part *a*.

a and b

- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [8] incorrect procedure.
- [9] C

[10] B

- [2] 1,225, and appropriate work is shown, such as solving an equation or writing an explanation.
- [1] Appropriate work is shown, but one computational error is made.
- or [1] Appropriate work is shown, but one conceptual error is made.
- or [1] Appropriate work is shown, but the conversion from years to months is incorrect, but an appropriate solution is found.
- or [1] 1,225, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [11] incorrect procedure.

[12] A

- [2] 77, and appropriate work is shown, such as $(76 + 78) \div 2$.
- [1] 76 and 78 are identified.
- or [1] 77, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [13] incorrect procedure.

[14] A

[2] An appropriate explanation is given, such as:

One very high or very low score in either class would have a great effect on the range for that class, but might not affect the median at all. The range is the difference between the two most extreme values, the lowest and the highest. The median, being the middle value, is not very sensitive to outliers or to extreme values.

- or [2] Specific examples are shown to illustrate the situation.
- [1] An understanding of median and range is demonstrated, but the specific situation is not explained.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [15] incorrect procedure.

[16] A

- [3] Mean = 79, median = 79, and mode = 78, and appropriate work is shown.
- [2] Appropriate work is shown, but only two of the three measures of central tendency are determined and identified correctly.
- or [2] Appropriate work is shown and all three measures of central tendency are determined correctly, but the measures are not identified or are identified incorrectly.
- [1] Appropriate work is shown, but only one of the three measures of central tendency is determined and identified correctly.
- or [1] Mean = 79, median = 79, and mode = 78, but no work is shown.
- [0] 79, 79, and 78, but no work is shown, and the answers are not identified or are identified incorrectly.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect precedure.
- [17] obviously incorrect procedure.

[18]	C
------	---

[19] D

[20] B

$\Gamma \cap 1$	
12.11	1 I <i>)</i>

[22] B

- [3] 34 and an appropriate explanation is given, such as $38 = \frac{46 + 2x}{3}$.
- [2] An appropriate method or equation is shown, but one computational mistake is made.
- or [2] The student does not take into consideration two dogs of equal weight and gives an answer of 68.
- [1] The student understands weighted average in that three dogs averaging 38 pounds must have a total weight of 114 pounds but does not subtract the known weight.
- or [1] 34 and no explanation is given.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [23] incorrect procedure.

- [2] \$350, and appropriate work is shown, such as $\frac{1450 + x}{5} = 360$ or trial and error with
- at least three trials and appropriate checks.
- [1] Appropriate work is shown, but one computational error is made.
- or [1] The total of the five salaries is shown to be $5 \times 360 = 1800$, but no further correct work is shown.
- or [1] \$350, but no work is shown or fewer than three trials with appropriate checks are shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.
- [27] incorrect procedure.

[3] 59 or 59°, and appropriate work is shown,

such as
$$63 = \frac{256 + x}{5}$$
 or

$$56 + 72 + 67 + 61 = 256$$
, $63 \times 5 = 315$, and $315 - 256 = 59$.

- [2] Appropriate work is shown, but one computational error is made.
- or [2] A value is chosen for Friday's temperature that rounds to 63, such as 57 or 61,

but whose mean is not exactly 63, and appropriate work is shown.

- [1] A limited understanding of the concept of the mean is shown, such as the sum of the temperatures must be 315, but the given temperatures are not subtracted.
- or [1] The correct mean of the four given temperatures is calculated.
- or [1] 59 or 59°, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [28] incorrect procedure.

[29] D

- [3] 95 and an appropriate method is shown that obtains an answer, such as 344 249 or a similar equation or method.
- or [3] Four scores are tried that round off to an average of 86, such as 93 or 94. Round off to 86 must be shown.
- [2] An appropriate method is shown, but one computational mistake is made.
- [1] The student understands weighted average and shows that the average of 83 for 3 tests is a total of 249 points.
- or [1] 95 and no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.
- [30] incorrect procedure.
- [31] A

- [3] 63, and appropriate work is shown, such as 400 (81 + 88 + 88) and determining the highest and lowest possible scores remaining that total 143.
- [2] Appropriate work is shown, but one computational error is made.
- [1] A total of 400 is shown, but one conceptual error is made, such as 257 is subtracted, and then 143 is split into 72 and 71, resulting in an answer of 71.
- or [1] Appropriate work is shown, but more than one computational error is made.
- or [1] No answer or an incorrect answer is found, but a list such as _____, ____, 81, 88, 88 is shown.
- or [1] 63, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously incorrect procedure.

- [4] Angelo is 66, Brandon is 26, and Carl is 31, and appropriate work is shown, such as solving an equation or trial and error with at least three trials and appropriate checks.
- [3] Appropriate work is shown, but one computational error is made.
- or [3] 66, 26, and 31, and appropriate work is shown, but the solutions are not labeled or are labeled incorrectly.
- [2] Appropriate work is shown, but two or more computational errors are made.
- or [2] Appropriate work is shown, but one conceptual error is made.
- or [2] The trial-and-error method is used to find a correct solution, but only two trials and appropriate checks are shown.
- or [2] The trial-and-error method is attempted and at least six systematic trials and appropriate checks are shown, but no solution is found.
- or [2] Carl is 31, and appropriate work is shown, but the ages of the other men are not found.
- or [2] An incorrect equation of equal difficulty is solved appropriately.
- [1] Appropriate work is shown, but one conceptual error and one computational error are made.
- or [1] A correct equation is written, but no further correct work is shown.
- or [1] Angelo is 66, Brandon is 26, and Carl is 31, but no work or only one trial with an appropriate check is shown.
- [0] Angelo is 66 *or* Brandon is 26 *or* Carl is 31, but no work is shown.
- or [0] 66, 26, and 31, but no work is shown, and the answers are not labeled or are labeled incorrectly.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an
- [33] obviously incorrect procedure.

- [4] Median = 91.5, mode = 92, and seventh test score = 96, and appropriate work is shown.
- [3] Appropriate work is shown, but one computational error is made.
- or [3] Seventh test score = 96, but only the median or the mode is found correctly, but appropriate work is shown.
- or [3] 91.5, 92, and 96, and appropriate work is shown, but the median and mode are not labeled or are labeled incorrectly.
- [2] Appropriate work is shown, but two or more computational errors are made.
- or [2] Appropriate work is shown, but one conceptual error is made.
- or [2] Both the median and the mode are found and labeled correctly, and appropriate work is shown, but the seventh test score is not found or is found incorrectly.
- or [2] Seventh test score = 96, and appropriate work is shown, but the median and the mode are not found or are found incorrectly.
- [1] Either the median or the mode is found and labeled correctly, and appropriate work is shown, but no further correct work is shown. or [1] Median = 91.5, mode = 92, and seventh
- test score = 96, but no work is shown.
- [0] Median = 91.5 or mode = 92 or seventh test score = 96, but no work is shown. or [0] 91.5, 92, and 96, but no work is shown and the answers are not labeled.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an
- [34] obviously incorrect procedure.
- [35] <u>C</u>

- [4] Mean = 225,000, median = 175,000, and the median is stated to be the best measure of central tendency, an appropriate justification is given, and appropriate work is shown.
- [3] Appropriate work is shown, but one computational error is made, but an appropriate measure of central tendency is stated, and an appropriate justification is given.
- or [3] Mean = 225,000, median = 175,000, and the median is stated to be the best measure of central tendency, but no justification is given.
- [2] Appropriate work is shown, but two or more computational errors are made, but an appropriate measure of central tendency is stated, and an appropriate justification is given.
- or [2] Appropriate work is shown, but one conceptual error is made.
- or [2] Appropriate work is shown to find mean = 225,000 and median = 175,000, but no further correct work is shown.
- [1] Appropriate work is shown, but one computational error and one conceptual error are made.
- or [1] Mean = 225,000 and median = 175,000, but no further work is shown.
- [0] Mean = 225,000 or median = 175,000, but no further work is shown.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an
- [36] obviously incorrect procedure.

[37]	D
------	---

[38] B

[39] A

[40] D

[41] B

[42] B

- [4] Correct cumulative frequencies of 7, 14, 24, and 30 and a fully labeled correct histogram are shown.
- [3] Incorrect cumulative frequencies are shown, but the histogram is appropriate for the data.
- or [3] Correct cumulative frequencies are shown, but a partially incorrect histogram is shown, such as the axes not being labeled, having nonequal intervals, or the x-axis starting at 50.
- [2] Only a frequency histogram is completed correctly.
- or [2] Only a correct cumulative frequency table and a correct bar graph are shown.
- [1] An appropriate bar graph is shown, but it is based on frequencies, not the cumulative frequency.
- or [1] Only a correct cumulative frequency table is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [43] incorrect procedure.
 - [4] The table is completed correctly, and an appropriate cumulative frequency histogram is drawn and labeled.
 - [3] The table is completed correctly, but one error is made in drawing the cumulative frequency histogram or one or more labeling errors are made.
 - or [3] The table is not completed correctly, but an appropriate cumulative frequency histogram is drawn, based on the table.
 - [2] One error is made in completing the table, and one graphing error is made in drawing the cumulative frequency histogram.
 - or [2] The table is completed correctly, but one conceptual error is made, such as drawing a frequency histogram or a cumulative frequency bar graph.
 - [1] The table is completed correctly, but no histogram is drawn.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously [44] incorrect procedure.

- [45] B
- [46] A
- [47] B
 - [2] y = 1.08x 2125 or an equivalent equation is written.
 - [1] One conceptual error is made, such as writing a regression equation that is not
 - or [1] The expression 1.08x 2125 is written, but no equation is written.
 - or [1] The correct values are identified for a and b, but no equation is written.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [48] incorrect procedure.

- [6] $\overline{W} = 44.6$ and $\overline{L} = 43.2$, the line of best-fit equation (y = -1.007559x + 88.137149) is shown, and an appropriate justification of point $(\overline{W}, \overline{L})$ fitting or not fitting, depending on the rounding of the equation, is given.
- [5] W or L is incorrect, but the rest of the work is appropriate.
- or [5] All conditions of the problem are met, except it is not stated whether $(\overline{W}, \overline{L})$ lies or does not lie on the line of best fit.
- or [5] \overline{W} and \overline{L} and the equation of the line of best fit are correct, but one error results in an incorrect conclusion, such as the
- calculation or interchanging of \overline{W} and \overline{L} .
- [4] Both \overline{W} and \overline{L} are incorrect, but the rest of the work is appropriate.
- or [4] W and L are correct, but the equation of the line of best fit is incorrect, but the justification is appropriate, based on the incorrect equation.
- or [4] \overline{W} and \overline{L} are correct, a correct scatter plot is drawn, a correct line of best fit is drawn, (W, L) is plotted correctly, and a statement indicating that the point does or does not fit the line is given, with an appropriate explanation, but no equation is
- or [4] All conditions of the problem are met, except for the justification of whether $(\overline{W}, \overline{L})$ lies on the line.
- [3] W and L are correct, but the equation of the line of best fit is stated incorrectly, and no further work is shown.
- [2] Only \overline{W} and \overline{L} are found correctly.
- [1] Only one mean is found correctly.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [49] incorrect procedure.

- [4] y = -34739.71292x + 313309.0909 and 209,090, and appropriate work is shown.
- [3] Appropriate work is shown, but one computational or rounding error is made.
- or [3] An incorrect linear equation with a negative slope is written, but an appropriate price is found for three blocks from the beach.
- [2] Appropriate work is shown, but two or more computational or rounding errors are
- or [2] Appropriate work is shown, but one conceptual error is made.
- or [2] A correct linear function is written, but no further correct work is shown.
- or [2] An incorrect linear equation with a positive slope is written, but an appropriate price is found for three blocks from the beach.
- [1] Appropriate work is shown, but one conceptual error and one computational or rounding error are made.
- or [1] 209,090, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [50] incorrect procedure.
 - [4] y = -0.58x + 1185.09 and 19.9, and appropriate work is shown.
 - [3] Appropriate work is shown, but one computational or rounding error is made.
 - [2] Appropriate work is shown, but two or more computational or rounding errors are made.
 - or [2] Appropriate work is shown, but one conceptual error is made.
 - or [2] A correct linear equation is written, but no further correct work is shown.
 - or [2] An incorrect linear equation is written, but an appropriate percentage is found.
 - [1] Appropriate work is shown, but one conceptual error and one computational or rounding error are made.
 - or [1] 19.9, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [51] incorrect procedure.

- [6] y = -6.2x + 12,451.2; 20.2 thousand; and 2008; and appropriate work is shown.
- [5] The correct equation is shown, but only the number of gallons or the year is correct.
- [4] The slope and y-intercept are incorrect, but the slope is negative and the number of gallons and the year are appropriate, based on the incorrect equation.
- [3] The slope and y-intercept are incorrect, but the slope is negative, but only the number of gallons or the year is appropriate, based on the incorrect equation.
- [2] The correct equation is shown, but the number of gallons and the year are not determined or are determined incorrectly.
- or [2] The incorrect equation y = 6.2x +12,451.2 is shown, but appropriate work is
- shown for the number of gallons and the year. [1] An incorrect equation is shown with a
- negative slope, and the number of gallons and the year are not determined.
- or [1] 20.2 thousand and 2008, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously [52] incorrect procedure.

- [4] a y = 0.8344648562x + 14.64960064 or an equivalent answer expressed to three significant digits
- and b 80, and appropriate work is shown.
- [3] One computational error is made or one rounding error is made with one of the numbers in the equation, such as truncating or not giving at least three significant digits.
- [2] Only the correct answer for either part a or part b is found.
- or [2] Appropriate work is shown, but more than one computational or rounding error is
- [1] 78 is substituted into an incorrect linear equation, but it is evaluated appropriately. or [1] y = 0.8344648562x + 14.64960064 and 80, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [53] incorrect procedure.
 - [4] $\bar{x} = 80$, $\bar{y} = 20.8$, and y = 0.25x + 0.8, and appropriate work is shown to prove that (\bar{x} , \overline{y}) is a point on the line of regression.
 - [3] Appropriate work is shown, but one computational error is made.
 - [2] Appropriate work is shown, but two or more computational errors are made. or [2] Appropriate work is shown, but one conceptual error is made.
 - [1] $\bar{x} = 80$, $\bar{y} = 20.8$, and y = 0.25x + 0.8, but no work is shown.
 - [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [54] incorrect procedure.

- [6] A correct scatter plot, y = 0.62x + 29.18, r = 0.92, and 83; and appropriate work is shown.
- [5] Appropriate work is shown, but one computational or rounding error is made.
- or [5] A correct scatter plot, equation, and score are shown, but no *r*-value is found.
- [4] A correct scatter plot and equation are shown, but the r-value and score are missing or incorrect.
- or [4] An incorrect equation is shown, but all further work is appropriate.
- or [4] The scatter plot is missing or incorrect, but all further work is appropriate.
- [3] The scatter plot is incorrect, but a correct equation and either a correct r-value or score are found.
- or [3] The scatter plot is correct, but an incorrect equation and either an appropriate rvalue or score based on the incorrect equation are found.
- [2] Only a correct scatter plot is shown, and all further work is missing or incorrect. or [2] Only a correct equation is shown, and all further work is missing or incorrect.
- [1] An incorrect equation is shown, but an appropriate score is found.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [55] incorrect procedure.

- [6] y = 0.01021x 1.66787, 4.56, and 913, and appropriate work is shown.
- [5] Appropriate work is shown, but one computational or rounding error is made. or [5] The expression 0.01021x - 1.66787 is written and 4.56 and 913 are found, and appropriate work is shown.
- [4] Appropriate work is shown, but two or more computational or rounding errors are
- or [4] A correct equation is written, but either the gross earnings or the number of theaters is not found, but appropriate work is shown. or [4] An incorrect equation of equal difficulty is written, but appropriate answers
- are found, and appropriate work is shown. [3] Appropriate work is shown, but one conceptual error is made.
- or [3] y = 0.01021x 1.66787, 4.56, and 913, but no work is shown.
- or [3] The expression 0.01021x 1.66787 is written and either 4.56 or 913 is found, and appropriate work is shown.
- [2] Appropriate work is shown, but one conceptual error and one computational or rounding error are made.
- or [2] A correct equation is written, but no further correct work is shown.
- [1] 4.56 and 913, but no work is shown.
- or [1] The expression 0.01021x 1.66787 is written, but no further correct work is shown.
- [0] Either 4.56 or 913, but no work is shown.
- or [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an
- obviously incorrect procedure.

- [6] p = 8.1875t + 72.7860, 1993, and 220.2, and appropriate work is shown.
- [5] Appropriate work is shown, but one computational or rounding error is made. or [5] The expression 8.1875t + 72.7860 is
- written and 1993 and 220.2 are found, and appropriate work is shown.
- [4] Appropriate work is shown, but two or more computational or rounding errors are made.
- or [4] A correct equation is written, but either the year or the predicted value for 2008 is not found, but appropriate work is shown.
- or [4] An incorrect equation is solved appropriately.
- [3] Appropriate work is shown, but one conceptual error is made.
- or [3] p = 8.1875t + 72.7860, 1993, and 220.2, but no work is shown.
- or [3] The expression 8.1875t + 72.7860 is written and either 1993 or 220.2 is found, and appropriate work is shown.
- [2] Appropriate work is shown, but one conceptual error and one computational or rounding error are made.
- or [2] A correct equation is written, but no further correct work is shown.
- or [2] 1993 and 220.2, but no work is shown.
- [1] The expression 8.1875t + 72.7860 is written, but no further correct work is shown.
- or [1] 1993 or 220.2, but no work is shown. [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously [57] incorrect procedure.

- [4] f(x) = 98.8571x + 737.3333 or y =98.8571x + 737.3333 and day 14, and appropriate substitution is made, such as 2050 = 98.8571x + 737.3333.
- [3] Appropriate work is shown, but one computational or rounding error is made. or [3] A correct linear regression equation is written and day 14, but no substitution is made.
- or [3] The expression 98.8571x + 737.3333 is written and day 14, and appropriate substitution is made, but no equation is written.
- [2] Appropriate work is shown, but two or more computational or rounding errors are
- or [2] Appropriate work is shown, but one conceptual error is made.
- or [2] A correct linear regression equation is written, but no further correct work is shown. or [2] An incorrect equation of equal
- difficulty is solved appropriately.
- [1] Appropriate work is shown, but one conceptual error and one computational or rounding error are made.
- or [1] The expression 98.8571x + 737.3333 is written, but no further correct work is shown. or [1] Day 14, but no work is shown.
- [0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously
- [58] incorrect procedure.