

1. 010101a, P.I. A.A.6

There are 461 students and 20 teachers taking buses on a trip to a museum. Each bus can seat a maximum of 52. What is the *least* number of buses needed for the trip?

[A] 8      [B] 11      [C] 10      [D] 9

2. 089914a, P.I. A.A.6

In a hockey league, 87 players play on seven different teams. Each team has at least 12 players. What is the largest possible number of players on any one team?

[A] 14      [B] 21      [C] 13      [D] 15

3. 069928a, P.I. A.A.6

A swimmer plans to swim at least 100 laps during a 6-day period. During this period, the swimmer will increase the number of laps completed each day by one lap. What is the *least* number of laps the swimmer must complete on the first day?

4. 080732a, P.I. A.A.6

Thelma and Laura start a lawn-mowing business and buy a lawnmower for \$225. They plan to charge \$15 to mow one lawn. What is the *minimum* number of lawns they need to mow if they wish to earn a profit of *at least* \$750?

5. 010737a, P.I. A.A.6

The Eye Surgery Institute just purchased a new laser machine for \$500,000 to use during eye surgery. The Institute must pay the inventor \$550 each time the machine is used. If the Institute charges \$2,000 for each laser surgery, what is the *minimum* number of surgeries that must be performed in order for the Institute to make a profit?

6. 060406a, P.I. A.A.6

Parking charges at Superior Parking Garage are \$5.00 for the first hour and \$1.50 for each additional 30 minutes. If Margo has \$12.50, what is the maximum amount of time she will be able to park her car at the garage?

[A]  $3\frac{1}{2}$  hours      [B]  $6\frac{1}{2}$  hours

[C]  $2\frac{1}{2}$  hours      [D] 6 hours

7. 080224a, P.I. A.A.6

A doughnut shop charges \$0.70 for each doughnut and \$0.30 for a carryout box. Shirley has \$5.00 to spend. At most, how many doughnuts can she buy if she also wants them in one carryout box?

8. 010938a, P.I. A.A.6

Mr. Braun has \$75.00 to spend on pizzas and soda pop for a picnic. Pizzas cost \$9.00 each and the drinks cost \$0.75 each. Five times as many drinks as pizzas are needed. What is the maximum number of pizzas that Mr. Braun can buy?

[1] C \_\_\_\_\_

[2] D \_\_\_\_\_

[3] 15 and an appropriate method or explanation is shown, such as trial and error or the inequality  $6x + 15 \geq 100$ .

[2] An appropriate method is shown, but it stops at 14.

[1] An appropriate method is shown, but no answer is found.

or [1] 15 and no explanation is given.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously

[3] incorrect procedure.

---

[2] 65, and appropriate work is shown, such as solving the inequality  $15x \geq 225 + 750$  or trial and error with at least three trials and appropriate checks.

[1] Appropriate work is shown, but one computational error is made.

or [1] Appropriate work is shown, but one conceptual error is made.

or [1] The trial-and-error method is attempted and at least six systematic trials and appropriate checks are shown, but no solution is found.

or [1] 65, but no work or fewer than three trials and appropriate checks are shown.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously

[4] incorrect procedure.

---

[3] 345, and appropriate work is shown, such as solving the inequality  $1450x > 500,000$ , solving an equation, or trial and error with at least three trials and appropriate checks.

[2] Appropriate work is shown, but one computational or rounding error is made.

or [2] The trial-and-error method is used to find a correct solution, but only two trials and appropriate checks are shown.

[1] Appropriate work is shown, but two or more computational or rounding errors are made.

or [1] Appropriate work is shown, but one conceptual error is made.

or [1] A correct inequality or equation is written, but no further correct work is shown.

or [1] The trial-and-error method is attempted and at least six systematic trials and appropriate checks are shown, but no solution is found.

or [1] 345, but no work or only one trial with an appropriate check is shown.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously

[5] incorrect procedure.

---

[6] A \_\_\_\_\_

[2] 6, and appropriate work is shown, such as  $0.70x + 0.30 \leq 5.00$  or trial and error with three trials and appropriate checks.

[1] The inequality is solved correctly, but the number of doughnuts is not found.

or [1] The trial-and-error method is used to find a correct solution, but fewer than three trials are shown.

or [1] 6, but no work is shown.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously

[7] incorrect procedure.

---

[4] 5, and appropriate work is shown, such as an algebraic solution or trial and error with at least three trials and appropriate checks.

[3] Appropriate work is shown, but one computational or rounding error is made.

[2] Appropriate work is shown, but two or more computational or rounding errors are made.

or [2] Appropriate work is shown, but one conceptual error is made.

or [2] The trial-and-error method is used, but only two trials and appropriate checks are shown.

or [2] The trial-and-error method is attempted, and at least six systematic trials and appropriate checks are shown, but no solution is found.

or [2] A correct inequality or equation is written, but no further correct work is shown.

[1] Appropriate work is shown, but one conceptual error and one computational or rounding error are made.

or [1] 5, but no work or only one trial with an appropriate check is shown.

[0] A zero response is completely incorrect, irrelevant, or incoherent or is a correct response that was obtained by an obviously

[8] incorrect procedure.