

# JMAP REGENTS BY DATE

NY Algebra I Regents Exam Questions  
from Fall 2023 to August 2025 Sorted by Date

[www.jmap.org](http://www.jmap.org)

**2023 Algebra I Sample Items**

1 What is the sum of  $3x\sqrt{7}$  and  $2x\sqrt{7}$ ?

1)  $5x\sqrt{7}$   
2)  $5x^2\sqrt{7}$

3)  $5x\sqrt{14}$   
4)  $5x^2\sqrt{14}$

2 What is an equation of the line that passes through the points (2,7) and (-1,3)?

1)  $y - 2 = \frac{3}{4}(x - 7)$

3)  $y - 7 = \frac{3}{4}(x - 2)$

2)  $y - 2 = \frac{4}{3}(x - 7)$

4)  $y - 7 = \frac{4}{3}(x - 2)$

3 Rationalize:  $\frac{3}{2\sqrt{6}}$

4 Use the method of completing the square to determine the exact values of  $x$  for the equation  $x^2 + 6x - 41 = 0$ . Express your answer in simplest radical form.

5 Solve the following systems of equations algebraically for all values of  $x$  and  $y$ :

$$y = x^2 + 5x - 17$$

$$x - y = 5$$

**0624AI**

- 1 A ball was launched into the air, and its height above the ground was recorded each second, as shown in the table below.

<b>Time (sec)</b>	0	1	2	3	4
<b>Height (ft)</b>	11	59	75	59	11


Based on these data, which statement is a valid conclusion?

- 1) The ball lands on the ground at 4 seconds.
- 2) The ball reaches a maximum height of 11 feet.
- 3) The ball was launched from a height of 0 feet.
- 4) The ball reaches its maximum height at 2 seconds.
- 2 A tour bus can seat, at most, 48 passengers. An adult ticket costs \$18 and a child ticket costs \$12. The bus company must collect at least \$650 to make a profit. If  $a$  represents the number of adult tickets sold and  $c$  represents the number of child tickets sold, which system of inequalities models this situation if they make a profit?
- 1)  $a + c < 48$   
 $18a + 12c > 650$
- 2)  $a + c \leq 48$   
 $18a + 12c \geq 650$
- 3)  $a + c < 48$   
 $18a + 12c < 650$
- 4)  $a + c \leq 48$   
 $18a + 12c \leq 650$
- 3 Which equation is always true?
- 1)  $x^2 \cdot x^3 = x^5$
- 2)  $3^x \cdot 3^2 = 9^{2x}$
- 3)  $-z^2 = z^2$
- 4)  $7^a \cdot 7^b = 7^{ab}$
- 4 The expression  $-2(x^2 - 2x + 1) + (3x^2 + 3x - 5)$  is equivalent to
- 1)  $x^2 + x - 4$
- 2)  $x^2 - x - 7$
- 3)  $x^2 + 7x - 4$
- 4)  $x^2 + 7x - 7$
- 5 Which sum is irrational?
- 1)  $-2\sqrt{12} + \sqrt{100}$
- 2)  $-\sqrt{4} + \frac{1}{3}\sqrt{900}$
- 3)  $\frac{1}{2}\sqrt{25} + \sqrt{64}$
- 4)  $\sqrt{49} + 3\sqrt{121}$
- 6 The solution to  $\frac{4(x-5)}{3} + 2 = 14$  is
- 1) 15
- 2) 14
- 3) 6
- 4) 4
- 7 On an island, a rare breed of rabbit doubled its population each month for two years. Which type of function best models the increase in population at the end of two years?
- 1) linear growth
- 2) linear decay
- 3) exponential growth
- 4) exponential decay





18 One Saturday, Dave took a long bike ride. The graph below models his trip.




What was Dave's average rate of change, in miles per hour, on this trip?

- 1) 10
- 2) 11
- 3) 11.6
- 4) 14.5

19 Which expression is equivalent to  $(x - 5)(2x + 7) - (x + 5)$ ?

- 1)  $2x^2 - 2x - 30$
- 2)  $2x^2 - 2x - 40$
- 3)  $2x^2 - 4x - 30$
- 4)  $2x^2 - 4x - 40$

20 The functions  $f(x)$  and  $g(x)$  are graphed on the set of axes below.



What is the solution to the equation  $f(x) = g(x)$ ?

- 1) 1 and 5
- 2) -5 and 0
- 3) -3 and 5
- 4) 0 and 4

- 21 When babysitting, Nicole charges an hourly rate and an additional charge for gas. She uses the function  $C(h) = 6h + 5$  to determine how much to charge for babysitting. The constant term of this function represents
- 1) the additional charge for gas
  - 2) the hourly rate Nicole charges
  - 3) the number of hours Nicole babysits
  - 4) the total Nicole earns from babysitting
- 22 When solved for  $x$  in terms of  $a$ , the solution to the equation  $3x - 7 = ax + 5$  is
- 1)  $\frac{12}{3a}$
  - 2)  $\frac{12}{3-a}$
  - 3)  $\frac{3a}{12}$
  - 4)  $\frac{3-a}{12}$
- 23 Wayne van Niekerk, a runner from South Africa, ran 400 meters in 43.03 seconds to set a world record. Which calculation would determine his average speed, in miles per hour?
- 1)  $\frac{400 \text{ m}}{43.03 \text{ sec}} \cdot \frac{1000 \text{ m}}{0.62 \text{ mi}} \cdot \frac{1 \text{ hr}}{3600 \text{ sec}}$
  - 2)  $\frac{400 \text{ m}}{43.03 \text{ sec}} \cdot \frac{0.62 \text{ mi}}{1000 \text{ m}} \cdot \frac{1 \text{ hr}}{3600 \text{ sec}}$
  - 3)  $\frac{400 \text{ m}}{43.03 \text{ sec}} \cdot \frac{0.62 \text{ mi}}{1000 \text{ m}} \cdot \frac{3600 \text{ sec}}{1 \text{ hr}}$
  - 4)  $\frac{400 \text{ m}}{43.03 \text{ sec}} \cdot \frac{1000 \text{ m}}{0.62 \text{ mi}} \cdot \frac{3600 \text{ sec}}{1 \text{ hr}}$
- 24 Which function has a domain of all real numbers and a range greater than or equal to three?
- 1)  $f(x) = -x + 3$
  - 2)  $g(x) = x^2 + 3$
  - 3)  $h(x) = 3^x$
  - 4)  $m(x) = |x + 3|$
- 25 Solve  $5(x - 2) \leq 3x + 20$  algebraically.
- 26 Given  $g(x) = x^3 + 2x^2 - x$ , evaluate  $g(-3)$ .
- 27 Given the relation  $R = \{(-1, 1), (0, 3), (-2, -4), (x, 5)\}$ . State a value for  $x$  that will make this relation a function. Explain why your answer makes this a function.
- 28 A survey of 150 students was taken. It was determined that  $\frac{2}{3}$  of the students play video games. Of the students that play video games, 85 also use social media. Of the students that do not play video games, 20% do not use social media. Complete the two-way frequency table.


	Play Video Games	Do Not Play Video Games	Total
Social Media			
No Social Media			
Total			

- 29 Use the method of completing the square to determine the exact values of  $x$  for the equation  $x^2 + 10x - 30 = 0$ .
- 30 Factor  $20x^3 - 45x$  completely.

- 31 Graph the following system of equations on the set of axes below.

$$y = x^2 - 3x - 6$$

$$y = x - 1$$



State the coordinates of all solutions.

- 32 The table below shows the amount of money a popular movie earned, in millions of dollars, during its first six weeks in theaters.

<b>Week (x)</b>	1	2	3	4	5	6
<b>Dollars Earned, in Millions (y)</b>	185	150	90	50	25	5


Write the linear regression equation for this data set, rounding all values to the *nearest hundredth*. State the correlation coefficient to the *nearest hundredth*. State what this correlation coefficient indicates about the linear fit of the data.

- 33 Use the quadratic formula to solve the equation  $3x^2 - 10x + 5 = 0$ . Express the answer in simplest radical form.

- 34 Graph the system of inequalities on the set of axes below.

$$3y + 2x \leq 15$$

$$y - x > 1$$



State the coordinates of a point in the solution to this system. Justify your answer.

- 35 Courtney went to a coffee shop to purchase lattes and donuts for her friends. One day she spent a total of \$15.50 on four lattes and two donuts. The next day she spent a total of \$18.10 on three lattes and five donuts. All prices included tax. If  $x$  represents the cost of one latte and  $y$  represents the cost of one donut, write a system of equations that can be used to model this situation. Courtney thinks that one latte costs \$2.75 and one donut costs \$2.25. Is Courtney correct? Justify your answer. Use your equations to determine algebraically the exact cost of one latte and the exact cost of one donut.





- 14 A survey of students at West High School was taken to determine a theme for the prom. The results of the survey are summarized in the table below.

	Beach Party	Hollywood	Broadway
Girls	86	112	68
Boys	123	77	79

Approximately what percentage of the students who chose the Broadway theme were girls?

- 1) 26  
2) 27  
3) 46  
4) 68
- 15 The sum of  $2\sqrt{54}$  and  $2\sqrt{6}$  is  
1)  $4\sqrt{60}$   
2)  $8\sqrt{15}$   
3)  $7\sqrt{6}$   
4)  $8\sqrt{6}$
- 16 The functions  $f(x) = x^2 - 5x - 14$  and  $g(x) = x + 2$  are graphed on the same set of axes. What are the solutions to the equation  $f(x) = g(x)$ ?  
1) -14 and 0  
2) 0 and 2  
3) -2 and 8  
4) -2 and 7
- 17 If  $x = 4a^2 - a + 3$  and  $y = a - 5$ , then which polynomial is equivalent to the product of  $x$  and  $y$ ?  
1)  $-17a^2 - 2a - 15$   
2)  $-17a^2 + 8a - 15$   
3)  $4a^3 - 21a^2 - 2a - 15$   
4)  $4a^3 - 21a^2 + 8a - 15$
- 18 What is an equation of the line that passes through (3,7) and has a slope of 2?  
1)  $y - 7 = 2(x - 3)$   
2)  $y - 3 = 2(x - 7)$   
3)  $y + 7 = 2(x + 3)$   
4)  $y + 3 = 2(x + 7)$
- 19 A geometric sequence with a common ratio of -3 is  
1) -10, -7, -4, -1, ...  
2) 14, 11, 8, 5, ...  
3) -2, -6, -18, -54, ...  
4) 4, -12, 36, -108, ...
- 20 When the equation  $6 - ax = ax - 2$  is solved for  $x$  in terms of  $a$ , and  $a \neq 0$ , the result is  
1)  $4a$   
2)  $\frac{4}{a}$   
3)  $2a$   
4)  $\frac{2}{a}$

21 Which function has the zeros  $-1$ ,  $3$ , and  $-4$ ?

1)  $f(x) = (x + 1)(x - 3)(x - 4)$

2)  $g(x) = (x - 1)(x + 3)(x - 4)$

3)  $h(x) = (x + 1)(x - 3)(x + 4)$

4)  $k(x) = (x - 1)(x + 3)(x + 4)$

22 The expression  $5^{a+2b}$  is equivalent to

1)  $5^a \cdot 5^2 \cdot 5^b$

2)  $5^a \cdot 25^b$

3)  $25^{2ab}$

4)  $25^{a+2b}$

23 In an arithmetic sequence, the first term is  $4$  and the third term is  $-2$ . What is the common difference?

1)  $-1$

2)  $-2$

3)  $-3$

4)  $-6$

24 Joe is ordering water for his swimming pool. He determines the volume of his pool to be about  $3240$  cubic feet. There are approximately  $7.5$  gallons of water in  $1$  cubic foot. A truck load holds  $6000$  gallons of water. Which expression would allow Joe to correctly calculate the number of truck loads of water he needs to fill his pool?


1)  $\frac{3240 \text{ ft}^3}{1 \text{ pool}} \cdot \frac{1 \text{ ft}^3}{7.5 \text{ gal}} \cdot \frac{6000 \text{ gal}}{1 \text{ truck load}}$

2)  $\frac{3240 \text{ ft}^3}{1 \text{ pool}} \cdot \frac{1 \text{ ft}^3}{7.5 \text{ gal}} \cdot \frac{1 \text{ truck load}}{6000 \text{ gal}}$

3)  $\frac{3240 \text{ ft}^3}{1 \text{ pool}} \cdot \frac{7.5 \text{ gal}}{1 \text{ ft}^3} \cdot \frac{6000 \text{ gal}}{1 \text{ truck load}}$

4)  $\frac{3240 \text{ ft}^3}{1 \text{ pool}} \cdot \frac{7.5 \text{ gal}}{1 \text{ ft}^3} \cdot \frac{1 \text{ truck load}}{6000 \text{ gal}}$

25 On the set of axes below, graph  $f(x) = x^2 + 4x + 1$ .



State the coordinates of the minimum.

26 If  $f(x) = \frac{30x^2}{x+2}$ , determine the value of  $f\left(\frac{1}{2}\right)$ .

27 Explain why the relation shown in the table below is a function.

<b>x</b>	-1	0	1	2
<b>y</b>	2	4	4	5

Complete the table below with values for both  $x$  and  $y$  so that this new relation is *not* a function.

<b>x</b>	-1	0	1	2	
<b>y</b>	2	4	4	5	

28 Solve algebraically for  $x$ :  $0.05(x - 3) = 0.35x - 7.5$

29 Use the quadratic formula to determine the exact roots of the equation  $x^2 + 3x - 6 = 0$ .

30 Factor  $5x^3 - 80x$  completely.

31 The owner of an ice cream stand kept track of the number of ice cream cones that were sold each day of the first week in June. She compared the ice cream sales to the average daily temperature. The data are shown in the table below.

<b>Average Daily Temp. (x)</b>	72	75	81	78	77	76	80
<b>Daily Ice Cream Cone Sales (y)</b>	126	183	263	229	200	185	249


State the linear regression equation for these data, rounding all values to the *nearest hundredth*. State the correlation coefficient, to the *nearest hundredth*, for the line of best fit for these data. State what this correlation coefficient indicates about the linear fit of the data.

- 32 Graph the system of inequalities on the set of axes below:

$$y > 3x - 4$$

$$x + 2y \leq 6$$

Label the solution set  $S$ .



Is the point  $(2, 2)$  a solution to the system? Justify your answer.

- 33 An object is launched upward at 64 feet per second from a platform 80 feet above the ground. The function  $s(t)$  models the height of the object  $t$  seconds after launch. If  $s(t) = -16t^2 + 64t + 80$ , state the vertex of  $s(t)$ , and explain in detail what each coordinate means in the context of the problem. After the object is launched, how many seconds does it take for the object to hit the ground? Justify your answer.

- 34 Solve the systems of equations algebraically for all values of  $x$  and  $y$ :

$$y = x^2 + 4x - 1$$


$$y = 2x + 7$$

- 35 Jen joined the Fan Favorite Movie Club at the local movie theater. At this theater, the cost of admission in May and June remains the same. In May, she saw 2 matinees and 3 regular-priced shows and spent \$38.50. In June, she went to 6 matinees and one regular-priced show and spent \$47.50. Write a system of equations to represent the cost,  $m$ , of a matinee ticket and the cost,  $r$ , of a regular-priced ticket. Jen said she spent \$5.75 on each matinee and \$9 on each regular show. Is Jen correct? Justify your answer. Use your system of equations to algebraically determine both the actual cost of each matinee ticket and the actual cost of each regular ticket.

**0125AI**

- 1 When factored, the expression  $x^3 - 36x$  is equivalent to
- |                       |                        |
|-----------------------|------------------------|
| 1) $(x + 6)(x - 6)$   | 3) $x(x + 6)(x - 6)$   |
| 2) $(x + 18)(x - 18)$ | 4) $x(x + 18)(x - 18)$ |
- 2 Which equation represents the line that passes through the points  $(-1, 8)$  and  $(4, -2)$ ?
- |                   |                      |
|-------------------|----------------------|
| 1) $y = -2x + 6$  | 3) $y = -0.5x + 7.5$ |
| 2) $y = -2x + 10$ | 4) $y = -0.5x + 8.5$ |
- 3 A geometric sequence is shown below.
- $$\frac{1}{2}, 2, 8, 32, \dots$$
- What is the common ratio?
- |                  |                  |
|------------------|------------------|
| 1) $\frac{1}{4}$ | 3) $\frac{1}{2}$ |
| 2) 2             | 4) 4             |
- 4 What is the constant term of the polynomial  $2x^3 - x + 5 + 4x^2$ ?
- |      |      |
|------|------|
| 1) 5 | 3) 3 |
| 2) 2 | 4) 4 |
- 5 A landscaping company charges a set fee for a spring cleanup, plus an hourly labor rate. The total cost is modeled by the function  $C(x) = 55x + 80$ . In this function, what does the 55 represent?
- |  |  |
|--|--|
| 1) the set fee for the cleanup         | 3) the profit earned by the company for one cleanup      |
| 2) the hourly labor rate for a cleanup | 4) the number of hours of labor required for one cleanup |
- 6 Which expression is equivalent to  $(5x^2 - 2x + 4) - (3x^2 + 3x - 1)$ ?
- |                    |                      |
|--------------------|----------------------|
| 1) $2x^2 + x + 3$  | 3) $2x^4 + x^2 + 3$  |
| 2) $2x^2 - 5x + 5$ | 4) $2x^4 - 5x^2 + 5$ |

7 A system of inequalities is graphed on the set of axes below.



Which point is a solution to this system?

- 1) (1,1)
- 2) (2,-2)
- 3) (1,8)
- 4) (4,2)

8 In an arithmetic sequence, the first term is 25 and the third term is 15. What is the tenth term in this sequence?

- 1) -20
- 2) -25
- 3) 70
- 4) 75

9 When the formula  $p = 2l + 2w$  is solved for  $w$ , the result is

- 1)  $w = \frac{2l+p}{2}$
- 2)  $w = \frac{p-2l}{2}$
- 3)  $w = \frac{p}{2} + l$
- 4)  $w = l - \frac{p}{2}$

10 Market Street Pizza kept a record of pizza sales for the month of February. The results are shown in the table below.


Type	Plain	Veggie	Meat Only	The Works
Thin Crust	300	80	120	100
Deep-dish	200	25	105	70

Of all the pizzas sold in February, what percent were plain, deep-dish pizzas?

- 1) 20%
- 2) 30%
- 3) 40%
- 4) 50%




16 The dot plots below represent test scores for 20 students on a math test.



The mode for this math test is 80 and the median is 85. Which dot plot correctly represents this data?

- 1) I
- 2) II
- 3) III
- 4) IV

17 A function is graphed on the set of axes below.



The domain of this function is

- 1)  $\{x|x > -2\}$
- 2)  $\{x|x \geq -2\}$
- 3)  $\{x|x > -4\}$
- 4)  $\{x|x \geq -4\}$

18 Which ordered pair is a solution to the equation  $y - 1 = 2\left(x + \frac{1}{4}\right)$ ?

- 1) (0.75, 0)
- 2) (1.25, 4)
- 3) (2.5, -6.5)
- 4) (4, -9.5)

19 Elena’s fastest time for the 50-meter dash is 7 seconds. She wants to know how fast this is in inches per minute. Which expression can Elena use for a correct conversion?

- 1)  $\frac{7 \text{ sec}}{50 \text{ meters}} \cdot \frac{60 \text{ sec}}{1 \text{ min}} \cdot \frac{1 \text{ meter}}{39.37 \text{ in}}$       3)  $\frac{50 \text{ meters}}{7 \text{ sec}} \cdot \frac{60 \text{ sec}}{1 \text{ min}} \cdot \frac{1 \text{ meter}}{39.37 \text{ in}}$   
 2)  $\frac{7 \text{ sec}}{50 \text{ meters}} \cdot \frac{1 \text{ min}}{60 \text{ sec}} \cdot \frac{39.37 \text{ in}}{1 \text{ meter}}$       4)  $\frac{50 \text{ meters}}{7 \text{ sec}} \cdot \frac{60 \text{ sec}}{1 \text{ min}} \cdot \frac{39.37 \text{ in}}{1 \text{ meter}}$





20 The table below shows the highest temperatures recorded in August for several years in one town.

Year	Temperature (°F)
1990	86
1991	78
1992	84
1993	95
1994	81
1995	77
1996	88
1997	93




The interquartile range of these data is

- 1) 7      3) 11  
 2) 10      4) 18

21 The function  $f(x) = x^2$  is multiplied by  $k$ , where  $k < -1$ . Which graph could represent  $g(x) = kf(x)$ ?

- 1) 
- 2) 
- 3) 
- 4) 

22 Which graph is the solution to the inequality  $6.4 - 4x \geq -2.8$ ?



23 The number of fish in a pond is eight more than the number of frogs. The total number of fish and frogs in the pond is at least 20. If  $x$  represents the number of frogs, which inequality can be used to represent this situation?





1)  $x + 8x \geq 20$

3)  $x + 8x \leq 20$


2)  $2x + 8 \geq 20$

4)  $2x + 8 \leq 20$

24 Which graph below represents a function that is always *decreasing* over the entire interval  $-3 < x < 3$ ?




25 The graph below models Sally's drive to the store.



State an interval when Sally is traveling at a constant speed. Explain your reasoning.

26 Graph the function  $f(x) = x^2 + 4x + 3$ .



State the equation of the axis of symmetry of  $f(x)$ .

27 The function  $f(x)$  is shown in the table below.

<b>x</b>	0	3	2	6	1	5	4	m
<b>f(x)</b>	6	2	7	5	8	4	3	9

State an appropriate value for  $m$  in the table, so that  $f(x)$  remains a function. Explain your reasoning.

28 Solve  $x^2 + 8x = 33$  for  $x$  by completing the square.

29 If  $f(x) = \frac{-3x-5}{2}$ , algebraically determine the value of  $x$  when  $f(x) = -22$ .

30 Rationalize the denominator of the fraction below. Express the solution in simplest form.

$$\frac{4}{\sqrt{2}}$$

31 Alex had \$1.70 in nickels and dimes on his desk. There were 25 coins in all. Write a system of equations that could be used to determine both the number of nickels,  $n$ , and the number of dimes,  $d$ , that Alex had. Use your system of equations to algebraically determine both the number of nickels and the number of dimes that he had.

32 The table below shows the average heart rate,  $x$ , and Calories burned,  $y$ , for seven men on an Olympic rowing team during a one-hour workout class.

<b>Average Heart Rate (x)</b>	135	147	150	144	146	153	143
<b>Calories Burned (y)</b>	725	812	866	761	825	863	737

Write the linear regression equation that models these data, rounding all values to the *nearest tenth*. State the correlation coefficient, rounded to the *nearest tenth*. State what the correlation coefficient suggests about the linear fit of these data.


33 Using the quadratic formula, solve  $x^2 + 4x - 3 = 0$ . Express your solution in simplest radical form.

34 Solve the following system of equations algebraically for all values of  $x$  and  $y$ :

$$y = x^2 - 7x + 12$$

$$y = 2x - 6$$

- 35 Anna plans to spend \$30 on balloons and party hats for her daughter's birthday party. Including tax, balloons cost \$2 each and party hats cost \$1.50 each. The number of party hats Anna needs is twice as many as the number of balloons. If  $x$  represents the number of balloons and  $y$  represents the number of party hats, write a system of equations that can be used to represent this situation. Graph your system of equations on the set of axes below.



State the coordinates of the point of intersection of your lines. Explain what each coordinate means in the context of the problem.

**0625AI**

1 The expression  $\frac{10}{\sqrt{2}}$  is equivalent to


1) 5

2) 20

3)  $5\sqrt{2}$

4)  $10\sqrt{2}$

2 A parabola is graphed on the set of axes below.



Over which interval is the parabola only increasing?

1)  $[1, 4]$

2)  $[3, \infty)$

3)  $(-\infty, 3]$

4)  $[-1, 1]$

3 Which scenario represents an exponential relationship?

1) Kirsten's New Year's resolution is to lose one pound each week.

2) Sarah wants to increase her grade by 5 points each quarter.

3) Tommy wants to reduce his spending by \$50 each month.

4) Dylan hopes to grow his business by 5% each month.

- 4 The geometry test scores for Andrea and Joe are shown in the table below.

Andrea	Joe
82	91
87	78
90	94
84	67

Which statement about their test scores is correct?

- 1) Both the mean and standard deviation of Andrea's test scores are higher than Joe's.
- 2) Both the mean and standard deviation of Joe's test scores are higher than Andrea's.
- 3) The mean of Andrea's test scores is higher than Joe's, but Joe's standard deviation is higher than Andrea's.
- 4) The mean of Joe's test scores is higher than Andrea's, but Andrea's standard deviation is higher than Joe's.
- 5 Which expression has a degree of 3 and a leading coefficient of 2?
- 1)  $2x^2 + 3x + 1$
- 2)  $6x^3 + 3x^2 - 2x$
- 3)  $3x^2 + 2x + 2$
- 4)  $2x^3 + x^2 + 4x$
- 6 The expression  $(-3x^2 + 9) - (7x^2 - 5x + 4)$  is equivalent to
- 1)  $-10x^2 + 5x + 5$
- 2)  $-10x^2 + 5x + 13$
- 3)  $-10x^2 - 5x + 5$
- 4)  $-10x^2 - 5x + 13$
- 7 The function  $h(x)$  is used to calculate the average height, in inches, of a tomato plant  $x$  weeks after it is transplanted. These data are represented in the table below.

$x$	$h(x)$
2	6
4	12
6	24
9	51
12	60
16	64

Between weeks 4 and 12, the average rate of change, in inches per week, is

- 1) 6
- 2) 8
- 3) 48
- 4) 58

8 Chloe is solving the equation  $x^2 + 5x = 3x + 3$ . Her first step is shown below.

$$\text{Given: } x^2 + 5x = 3x + 3$$

$$\text{Step 1: } x^2 + 2x - 3 = 0$$

Which property justifies this step?

- |                              |   |
|------------------------------|---|
| 1) the zero product property | 3) the distributive property            |
| 2) the commutative property  | 4) the subtraction property of equality |

9 Which function represents the graph of  $w(x) = |x|$  shifted 2 units to the right?

- |                     |                     |
|---------------------|---------------------|
| 1) $g(x) =  x + 2 $ | 3) $q(x) =  x  + 2$ |
| 2) $h(x) =  x - 2 $ | 4) $r(x) =  x  - 2$ |

10 Given the system of equations:

$$y + 4x = 5$$

$$2x - 3y = 10$$


A step in solving this system by using the substitution method would be

- |                         |                          |
|-------------------------|--------------------------|
| 1) $2(5 - 4x) + 4x = 5$ | 3) $2x - 3(5 - 4x) = 10$ |
| 2) $2(5 + 4x) + 4x = 5$ | 4) $2x - 3(5 + 4x) = 10$ |

11 Which equation is equivalent to  $x^2 - 6x = 27$ ?

- |                         |                          |
|-------------------------|--------------------------|
| 1) $(x - 3)^2 = 27 - 9$ | 3) $(x - 3)^2 = 27 + 36$ |
| 2) $(x - 3)^2 = 27 + 9$ | 4) $(x - 3)^2 = 27 - 36$ |

12 The box plots below summarize the ages of athletes on the swim team and the track team.




Based on the box plots, which statement must be true?

- |   |  |
|---|--|
| 1) The IQR of both teams is the same.                               | 3) The median age of the swim team is less than the median age of the track team.          |
| 2) There are more athletes on the swim team than on the track team. | 4) The range of ages of the swim team is smaller than the range of ages of the track team. |



19 Four graphs are shown below.



Which of the graphs represent(s) a function?

- 1) *A*, only  
 2) *A* and *B*, only  
 3) *A*, *B*, and *C*, only  
 4) *A*, *B*, *C*, and *D*

20 The formula to calculate kinetic energy is  $K = \frac{1}{2}mv^2$ , where  $K$  is kinetic energy,  $m$  is mass, and  $v$  is velocity.

When  $m$  is written in terms of  $K$  and  $v$ , the equation is

- 1)  $m = \frac{2K}{v^2}$   
 2)  $m = 2Kv^2$   
 3)  $m = \sqrt{2Kv^2}$   
 4)  $m = \frac{K}{2v^2}$


21 The solution to the equation  $\frac{2(3x-1)}{3} = x+2$  is

- 1)  $\frac{1}{3}$   
 2)  $\frac{2}{3}$   
 3)  $\frac{4}{3}$   
 4)  $\frac{8}{3}$

22 Which equation represents the sequence  $12, 6, 3, \frac{3}{2}, \dots$ , where  $a_1 = 12$ ?

- 1)  $a_n = 12 \cdot \left(\frac{1}{2}\right)^{n-1}$   
 2)  $a_n = 12 \cdot \left(\frac{1}{2}\right)^n$   
 3)  $a_n = 12 \cdot (2)^{n-1}$   
 4)  $a_n = 12 \cdot (2)^n$

23 The axis of symmetry is  $x = 2$  for which quadratic function?



$x$	$g(x)$
-2	6
-1	3
0	2
1	3
2	6

1)

2)  $j(x) = 2x^2 + 8x$

3)

4)  $h(x) = x^2 - 4x - 5$

24 Each day, a freight train passes by Anna's house. This freight train travels at 49 miles per hour. Each railroad car is 56 feet long. Which expression represents the number of railroad cars that pass by Anna's house per minute?

1)  $\frac{49 \text{ mi}}{1 \text{ hr}} \cdot \frac{1 \text{ mi}}{5280 \text{ ft}} \cdot \frac{1 \text{ hr}}{60 \text{ min}} \cdot \frac{1 \text{ car}}{56 \text{ ft}}$

3)  $\frac{49 \text{ mi}}{1 \text{ hr}} \cdot \frac{5280 \text{ ft}}{1 \text{ mi}} \cdot \frac{1 \text{ hr}}{60 \text{ min}} \cdot \frac{1 \text{ car}}{56 \text{ ft}}$

2)  $\frac{49 \text{ mi}}{1 \text{ hr}} \cdot \frac{1 \text{ mi}}{5280 \text{ ft}} \cdot \frac{60 \text{ min}}{1 \text{ hr}} \cdot \frac{1 \text{ car}}{56 \text{ ft}}$

4)  $\frac{49 \text{ mi}}{1 \text{ hr}} \cdot \frac{5280 \text{ ft}}{1 \text{ mi}} \cdot \frac{60 \text{ min}}{1 \text{ hr}} \cdot \frac{1 \text{ car}}{56 \text{ ft}}$

25 A survey was taken to determine whether students preferred to watch videos or listen to music. Of the 100 students surveyed, 44 were seniors. Of the 65 students who preferred to watch videos, 42 were juniors. Use this information to complete the frequency table below.

	Juniors	Seniors	Total
Watch Videos			
Listen to Music			
Total			

26 Solve the inequality for  $y$ :  $5(2 - y) > -11y - 8$


27 Express  $(5x - 3)(-2x + 7)$  as a trinomial in standard form.

28 The first and fourth terms in an arithmetic sequence are given below.

$$-20, \_, \_, -2$$


Determine the eighth term.

- 29 Write an equation in slope-intercept form for the line that passes through  $(-2, 5)$  and has a slope of  $-3$ . [Use of the set of axes below is optional.]



- 30 Factor the expression  $x^3 - 36x$  completely.

- 31 Graph  $f(x) = -3x$  and  $g(x) = x^2 + 2$  on the set of axes below.



State the values of  $x$  that satisfy the equation  $f(x) = g(x)$ .

- 32 Using the quadratic formula, solve  $6x^2 + 2x - 1 = 0$ . Express the answer in simplest radical form.
- 33 The table below shows the price of a new cell phone and the length of time, in months, since its release.

<b>Time Since Release, in Months (x)</b>	0	3	6	9	12
<b>Price, in Dollars (y)</b>	1200	1150	1100	1000	920


State the linear regression equation for this set of data. Round all values to the *nearest hundredth*. State the correlation coefficient for this data set, to the *nearest hundredth*. State what the correlation coefficient indicates about the linear fit of the data.

- 34 Solve the following system of equations algebraically for all values of  $x$  and  $y$ :

$$y = x^2 + 9x + 4$$

$$y - 2x = -6$$

- 35 Sarah earns \$6 per hour babysitting and \$12 per hour tutoring. Her goal is to earn at least \$120 per week. Sarah is allowed to work a maximum of 14 hours per week doing both jobs. If  $x$  represents the number of hours Sarah babysits and  $y$  represents the number of hours she tutors, write a system of inequalities that could model this situation. On the set of axes below, graph the system of inequalities that you wrote.



State a combination of hours babysitting and tutoring that would satisfy this situation. Justify your answer.

**0825AI**

- 1 Which expression is equivalent to  $100x^2 - 16$ ?
- 1)  $(50x - 8)(50x + 8)$                       3)  $(10x - 4)(10x + 4)$   
2)  $(50x - 8)(50x - 8)$                       4)  $(10x - 4)(10x - 4)$
- 2 Josie has \$2.30 in dimes and quarters. She has two more dimes than quarters. Which equation below can be used to determine  $x$ , the number of quarters she has?
- 1)  $0.35(2x + 2) = 2.30$                       3)  $0.25x + 0.10(x + 2) = 2.30$   
2)  $0.25(x + 2) + 0.10x = 2.30$                       4)  $0.25x + 0.10(x - 2) = 2.30$
- 3 If  $g(x) = -2x^2 + 16$  then  $g(-3)$  equals
- 1)  $-20$     3)  $34$   
2)  $-2$     4)  $52$
- 4 What are the zeros of  $f(x) = x^2 - 8x - 20$ ?
- 1)  $10$  and  $2$                                       3)  $-10$  and  $2$   
2)  $10$  and  $-2$                                       4)  $-10$  and  $-2$
- 5 Which point lies on the graph of  $y = 3x^2 - \frac{1}{4}x + 3$ ?
- 1)  $(-2, 15.5)$                                       3)  $(1, 6.25)$   
2)  $(-1, 5.75)$                                       4)  $(2, 15.5)$
- 6 Given  $f(x) = x^2$  and  $g(x) = 8x - 15$  graphed on the same set of axes, which value(s) of  $x$  will make  $f(x) = g(x)$ ?
- 1)  $3$ , only    3)  $3$  and  $5$   
2)  $9$ , only    4)  $9$  and  $25$
- 7 Which trinomial is written in standard form and has a constant term of five?
- 1)  $x^5 - 4x^2 + 10$                                       3)  $5x^4 - 3x^2 + 1$   
2)  $2x^2 + 6x^4 + 5$                                       4)  $4x^5 - 8x^2 + 5$
- 8 When solving  $x^2 + 6x = -8$  for  $x$ , a student wrote  $x^2 + 6x + 8 = 0$  as their first step. Which property justifies this step?
- 1) associative property                              3) zero property of addition  
2) commutative property                              4) addition property of equality





18 Three functions are given below.

$$f(x) = -|x + 2| + 7$$

$$g(x) = (x - 3)^2 - 4$$

x	h(x)
-4	5
-3	0
-2	-3
-1	-4
0	-3
1	0
2	5

Which functions have the same y-intercept?

- 1)  $f(x)$  and  $g(x)$
- 2)  $g(x)$  and  $h(x)$
- 3)  $f(x)$  and  $h(x)$
- 4) The functions all have different y-intercepts.

19 The sum of  $(x + 7)^2$  and  $(x - 3)^2$  is

- 1)  $2x^2 + 58$
- 2)  $2x^4 + 58$
- 3)  $2x^2 + 8x + 58$
- 4)  $2x^4 + 8x^2 + 58$

20 The product of  $2\sqrt{10}$  and  $3\sqrt{2}$  is



- 1)  $12\sqrt{5}$
- 2)  $5\sqrt{20}$
- 3)  $24\sqrt{5}$
- 4)  $5\sqrt{12}$

21 When  $6x^3 - 2x + 8$  is subtracted from  $5x^3 + 3x - 4$ , the result is

- 1)  $x^3 - 5x + 12$
- 2)  $x^3 + x + 4$
- 3)  $-x^3 + 5x - 12$
- 4)  $-x^3 + x + 4$

22 Three relations are shown below.

I.  $\{(0,1), (1,2), (2,3), (3,4)\}$



Which relations represent a function?


- 1) I and II, only
- 2) I and III, only
- 3) II and III, only
- 4) I, II, and III



- 28 A survey was given to 180 cell phone owners about the brand of phone they owned. The results showed that 59 adults owned Brand *B* and 32 teenagers owned Brand *A*. Of all the people surveyed, 40% owned Brand *A*. Complete the two-way frequency table below.

	Brand A	Brand B	Total
Adults			
Teenagers			
Total			

- 29 Determine the 8th term of a geometric sequence whose first term is 5 and whose common ratio is 3.
- 30 Using the method of completing the square, express  $x^2 + 14x - 28 = 0$  in the form  $(x - p)^2 = q$ .
- 31 Graph  $f(x) = -\frac{1}{3}x^2 + 4$  on the set of axes below.




State the vertex of this function. State the equation of the axis of symmetry of this function.

- 32 Vince wants to rent a canoe while he is on vacation. The canoe rental company charges \$18 for the first hour and \$7.50 for each additional hour,  $x$ . If Vince has \$78 to spend on renting a canoe, write an inequality in terms of  $x$  that models this situation. Algebraically determine the maximum number of hours that Vince could rent a canoe.

- 33 Graph the following system of inequalities on the set of axes below:

$$y \geq -\frac{1}{2}x - 3$$

$$y - 2x < 5$$



State the coordinates of a point that is in the solution to this system. Justify your answer.

- 34 Using the quadratic formula, solve  $x^2 - 6x + 3 = 0$ . Express the answer in simplest radical form.
- 35 Cameron sold hot dogs and sodas at a concession stand. He sold a total of 25 items for \$45.00. A hot dog sold for \$2.25 and a soda sold for \$1.50. All prices include tax. If  $x$  represents the number of hot dogs sold and  $y$  represents the number of sodas sold, write a system of equations that models this situation. Determine algebraically the number of hot dogs Cameron sold and the number of sodas he sold. A customer has \$20 to spend at the concession stand. Determine and state the maximum number of hot dogs he can purchase if he buys four sodas.

## 2023 Algebra I Sample Items Answer Section

- 1 ANS: 1                      PTS: 2                      REF: fall2301ai      NAT: N.RN.B.3  
TOP: Operations with Radicals                      KEY: addition
- 2 ANS: 4  
$$m = \frac{7-3}{2--1} = \frac{4}{3}$$
- PTS: 2                      REF: fall2302ai      NAT: A.REI.D.10      TOP: Writing Linear Equations  
KEY: other forms
- 3 ANS:  
$$\frac{3}{2\sqrt{6}} \cdot \frac{\sqrt{6}}{\sqrt{6}} = \frac{3\sqrt{6}}{12}$$
- PTS: 2                      REF: fall2303ai      NAT: N.RN.B.3      TOP: Operations with Radicals  
KEY: division
- 4 ANS:  
$$x^2 + 6x + 9 = 41 + 9$$
  
$$(x + 3)^2 = 50$$
  
$$x + 3 = \pm\sqrt{50}$$
  
$$x = -3 \pm 5\sqrt{2}$$
- PTS: 4                      REF: fall2304ai      NAT: A.REI.B.4      TOP: Solving Quadratics  
KEY: completing the square
- 5 ANS:  
$$x^2 + 5x - 17 = x - 5 \quad -6 - y = 5 \quad 2 - y = 5 \quad (-6, -11), (2, -3)$$
  
$$x^2 + 4x - 12 = 0 \quad y = -11 \quad y = -3$$
  
$$(x + 6)(x - 2) = 0$$
  
$$x = -6, 2$$
- PTS: 4                      REF: fall2305ai      NAT: A.REI.C.7      TOP: Quadratic-Linear Systems

## 0624AI

## Answer Section

- 1 ANS: 4                      PTS: 2                      REF: 062401ai                      NAT: F.IF.B.4  
TOP: Graphing Quadratic Functions                      KEY: key features
- 2 ANS: 2                      PTS: 2                      REF: 062402ai                      NAT: A.CED.A.3  
TOP: Modeling Systems of Linear Inequalities
- 3 ANS: 1                      PTS: 2                      REF: 062403ai                      NAT: A.APR.A.1  
TOP: Multiplication of Powers
- 4 ANS: 4  
 $-2x^2 + 4x - 2 + 3x^2 + 3x - 5 = x^2 + 7x - 7$
- PTS: 2                      REF: 062404ai                      NAT: A.APR.A.1                      TOP: Operations with Polynomials  
KEY: addition
- 5 ANS: 1                      PTS: 2                      REF: 062405ai                      NAT: N.RN.B.3  
TOP: Operations with Radicals                      KEY: classify
- 6 ANS: 2  
 $\frac{4(x-5)}{3} = 12$   
 $4x - 20 = 36$   
 $4x = 56$   
 $x = 14$
- PTS: 2                      REF: 062406ai                      NAT: A.REI.B.3                      TOP: Solving Linear Equations
- 7 ANS: 3                      PTS: 2                      REF: 062407ai                      NAT: F.LE.A.1  
TOP: Families of Functions
- 8 ANS: 3                      PTS: 2                      REF: 062408ai                      NAT: A.SSE.A.1  
TOP: Modeling Expressions
- 9 ANS: 2                      PTS: 2                      REF: 062409ai                      NAT: A.APR.B.3  
TOP: Zeros of Polynomials
- 10 ANS: 3  
 $\frac{5 - -1}{-1 - 2} = \frac{6}{-3} = -2$   $5 = -2(-1) + b$   
 $3 = b$
- PTS: 2                      REF: 062410ai                      NAT: F.IF.B.4                      TOP: Graphing Linear Functions
- 11 ANS: 1  
 $a(8) = 2^8 + 25 = 281$   $b(8) = 10(8) + 75 = 155$   $c(8) = \sqrt{400(8)} + 80 \approx 137$   $d(8) = 2(8 + 1)^2 - 10(8) + 50 = 132$
- PTS: 2                      REF: 062411ai                      NAT: F.LE.A.3                      TOP: Families of Functions

12 ANS: 2

$$25r^2 = 625$$

$$r^2 = 25$$

$$r = \pm 5$$

PTS: 2 REF: 062412ai NAT: F.IF.A.3 TOP: Sequences  
KEY: difference or ratio

13 ANS: 2

$$110 - 60 = 50$$

PTS: 2 REF: 062413ai NAT: S.ID.A.1 TOP: Box Plots  
KEY: interpret

14 ANS: 1

$$1) -7; 2) -4; 3) x = \frac{-6}{2(1)} = -3, c(-3) = (-3)^2 + 6(-3) + 3 = -6; 4) -5$$

PTS: 2 REF: 062414ai NAT: F.IF.C.9 TOP: Comparing Quadratic Functions  
15 ANS: 2 PTS: 2 REF: 062415ai NAT: F.BF.A.1  
TOP: Sequences KEY: explicit

16 ANS: 2

$$\text{mean: } \frac{3(0) + 3(1) + 4(2) + 5(3) + 2(4) + 2(5) + 1(6)}{3 + 3 + 4 + 5 + 2 + 2 + 1} = \frac{50}{20} = 2.5, \text{ mode: } 3, \text{ median: } \frac{2+3}{2} = 2.5$$

PTS: 2 REF: 062416ai NAT: S.ID.A.1 TOP: Dot Plots  
17 ANS: 4 PTS: 2 REF: 062417ai NAT: F.BF.B.3  
TOP: Transformations with Functions

18 ANS: 1

$$\frac{55 - 0}{5.5 - 0} = 10$$

PTS: 2 REF: 062418ai NAT: F.IF.B.6 TOP: Rate of Change  
19 ANS: 4

$$2x^2 + 7x - 10x - 35 - x - 5 = 2x^2 - 4x - 40$$

PTS: 2 REF: 062419ai NAT: A.APR.A.1 TOP: Operations with Polynomials  
KEY: multiplication  
20 ANS: 1 PTS: 2 REF: 062420ai NAT: A.REI.D.11  
TOP: Quadratic-Linear Systems  
21 ANS: 1 PTS: 2 REF: 062421ai NAT: F.LE.B.5  
TOP: Modeling Linear Functions

22 ANS: 2

$$3x - ax = 12$$

$$x(3 - a) = 12$$

$$x = \frac{12}{3 - a}$$

PTS: 2 REF: 062422ai NAT: A.REI.B.3 TOP: Solving Linear Equations

KEY: coefficients represented by letters

23 ANS: 3

PTS: 2

REF: 062423ai

NAT: N.Q.A.1

TOP: Conversions

24 ANS: 2

All four functions have a real domain.  $f$  has a real range.  $h$  has a positive real range.  $m$  has a nonnegative real range.

PTS: 2

REF: 062424ai

NAT: F.IF.A.2

TOP: Domain and Range

25 ANS:

$$5x - 10 \leq 3x + 20$$

$$2x \leq 30$$

$$x \leq 15$$

PTS: 2

REF: 062425ai

NAT: A.REI.B.3

TOP: Solving Linear Inequalities

26 ANS:

$$g(-3) = (-3)^3 + 2(-3)^2 - (-3) = -27 + 18 + 3 = -6$$

PTS: 2

REF: 062426ai

NAT: F.IF.A.2

TOP: Functional Notation

27 ANS:

$x$  may be any value other than  $-2, -1, 0$ , so that for any value of  $x$ , there is a unique  $y$ .

PTS: 2

REF: 062427ai

NAT: F.IF.A.1

TOP: Defining Functions

28 ANS:

	Play Video Games	Do Not Play Video Games	Total
Social Media	85	40	125
No Social Media	15	10	25
Total	100	50	150

PTS: 2

REF: 062428ai

NAT: S.ID.B.5

TOP: Frequency Tables

KEY: two-way

29 ANS:

$$x^2 + 10x = 30$$

$$x^2 + 10x + 25 = 30 + 25$$

$$(x + 5)^2 = 55$$

$$x + 5 = \pm\sqrt{55}$$

$$x = -5 \pm \sqrt{55}$$

PTS: 2 REF: 062429ai NAT: A.REI.B.4 TOP: Solving Quadratics

KEY: completing the square


30 ANS:

$$20x^3 - 45x = 5x(4x^2 - 9) = 5x(2x + 3)(2x - 3)$$

PTS: 2 REF: 062430ai NAT: A.SSE.A.2

TOP: Factoring the Difference of Perfect Squares

31 ANS:



PTS: 4 REF: 062431ai NAT: A.REI.C.7 TOP: Quadratic-Linear Systems

32 ANS:

$$y = -37.57x + 215.67, -0.98, \text{strong}$$

PTS: 4 REF: 062432ai NAT: S.ID.B.6 TOP: Regression

KEY: linear with correlation coefficient


33 ANS:

$$x = \frac{-(-10) \pm \sqrt{(-10)^2 - 4(3)(5)}}{2(3)} = \frac{10 \pm \sqrt{40}}{6} = \frac{10 \pm 2\sqrt{10}}{6} = \frac{5 \pm \sqrt{10}}{3}$$

PTS: 4 REF: 062433ai NAT: A.REI.B.4 TOP: Solving Quadratics

KEY: quadratic formula

34 ANS:



$(-1, 1)$  is a solution as it is in the overlap area.

PTS: 4

REF: 062434ai

NAT: A.REI.D.12

TOP: Graphing Systems of Linear Inequalities

35 ANS:

$4x + 2y = 15.5$   $5(4x + 2y = 15.5)$  Courtney is incorrect because of the following calculations:  $20x + 10y = 77.5$

$3x + 5y = 18.1$   $2(3x + 5y = 18.1)$

$6x + 10y = 36.2$

$14x = 41.3$

$x = 2.95$

$4(2.95) + 2y = 15.5$

$11.8 + 2y = 15.5$

$2y = 3.7$

$y = 1.85$

PTS: 6

REF: 062435ai

NAT: A.CED.A.3

TOP: Modeling Linear Systems

## 0824AI

## Answer Section

- 1 ANS: 4                      PTS: 2                      REF: 082401ai                      NAT: A.SSE.A.2  
TOP: Factoring Polynomials
- 2 ANS: 1                      PTS: 2                      REF: 082402ai                      NAT: F.LE.A.1  
TOP: Families of Functions
- 3 ANS: 4  
 $3(x^2 - 2x + 3) - (4x^2 + 3x - 1)$   
 $3x^2 - 6x + 9 - 4x^2 - 3x + 1$   
 $-x^2 - 9x + 10$
- PTS: 2                      REF: 082403ai                      NAT: A.APR.A.1                      TOP: Operations with Polynomials  
KEY: subtraction
- 4 ANS: 2                      PTS: 2                      REF: 082404ai                      NAT: A.CED.A.1  
TOP: Modeling Linear Equations
- 5 ANS: 1                      PTS: 2                      REF: 082405ai                      NAT: A.SSE.A.1  
TOP: Modeling Expressions
- 6 ANS: 4                      PTS: 2                      REF: 082406ai                      NAT: A.REI.A.1  
TOP: Identifying Properties
- 7 ANS: 4                      PTS: 2                      REF: 082407ai                      NAT: N.RN.B.3  
TOP: Operations with Radicals  
KEY: classify
- 8 ANS: 2  
 $x^2 + 6x = 18$   
 $x^2 + 6x + 9 = 18 + 9$   
 $(x + 3)^2 = 27$
- PTS: 2                      REF: 082408ai                      NAT: A.REI.B.4                      TOP: Solving Quadratics  
KEY: completing the square
- 9 ANS: 3  
69,70,70,71,72,74,76,78 ordered. median:  $\frac{71+72}{2} = 71.5$
- PTS: 2                      REF: 082409ai                      NAT: S.ID.A.1                      TOP: Box Plots  
KEY: represent
- 10 ANS: 3  
 $\frac{425-50}{350-100} = 1.5$
- PTS: 2                      REF: 082410ai                      NAT: F.IF.B.6                      TOP: Rate of Change
- 11 ANS: 3                      PTS: 2                      REF: 082411ai                      NAT: F.BF.B.3  
TOP: Transformations with Functions
- 12 ANS: 1                      PTS: 2                      REF: 082412ai                      NAT: F.LE.B.5  
TOP: Modeling Linear Functions

- 13 ANS: 4  
 $2m - 4 \leq 3(2m + 4)$   
 $2m - 4 \leq 6m + 12$   
 $-16 \leq 4m$   
 $-4 \leq m$
- PTS: 2 REF: 082413ai NAT: A.REI.B.3 TOP: Solving Linear Inequalities
- 14 ANS: 3  
 $\frac{68}{68 + 79} \approx 0.46$
- PTS: 2 REF: 082414ai NAT: S.ID.B.5 TOP: Frequency Tables  
 KEY: two-way
- 15 ANS: 4  
 $2\sqrt{54} + 2\sqrt{6} = 2\sqrt{9}\sqrt{6} + 2\sqrt{6} = 6\sqrt{6} + 2\sqrt{6} = 8\sqrt{6}$
- PTS: 2 REF: 082415ai NAT: N.RN.B.3 TOP: Operations with Radicals  
 KEY: addition
- 16 ANS: 3  
 $x^2 - 5x - 14 = x + 2$   
 $x^2 - 6x - 16 = 0$   
 $(x - 8)(x + 2) = 0$   
 $x = 8, -2$
- PTS: 2 REF: 082416ai NAT: A.REI.D.11 TOP: Quadratic-Linear Systems
- 17 ANS: 4  
 $(4a^2 - a + 3)(a - 5) = 4a^3 - 20a^2 - a^2 + 5a + 3a - 15 = 4a^3 - 21a^2 + 8a - 15$
- PTS: 2 REF: 082417ai NAT: A.APR.A.1 TOP: Operations with Polynomials  
 KEY: multiplication
- 18 ANS: 1 PTS: 2 REF: 082418ai NAT: A.REI.D.10  
 TOP: Writing Linear Equations KEY: other forms
- 19 ANS: 4 PTS: 2 REF: 082419ai NAT: F.IF.A.3  
 TOP: Sequences KEY: difference or ratio

20 ANS: 2

$$6 - ax = ax - 2$$

$$8 = 2ax$$

$$\frac{8}{2a} = x$$

$$\frac{4}{a} = x$$

PTS: 2

REF: 082420ai

NAT: A.REI.B.3

TOP: Solving Linear Equations

KEY: coefficients represented by letters

21 ANS: 3

PTS: 2

REF: 082421ai

NAT: A.APR.B.3

TOP: Zeros of Polynomials

22 ANS: 2

$$5^{a+2b} = 5^a \cdot 5^{2b} = 5^a \cdot 25^b$$

PTS: 2

REF: 082422ai

NAT: A.APR.A.1

TOP: Multiplication of Powers

23 ANS: 3

$$\frac{-2-4}{3-1} = \frac{-6}{2} = -3$$

PTS: 2

REF: 082423ai

NAT: F.IF.A.3

TOP: Sequences

KEY: difference or ratio

24 ANS: 4


PTS: 2

REF: 082424ai

NAT: N.Q.A.1

TOP: Conversions

25 ANS:



PTS: 2

REF: 082425ai

NAT: F.IF.C.7

TOP: Graphing Quadratic Functions

26 ANS:

$$f\left(\frac{1}{2}\right) = \frac{30\left(\frac{1}{2}\right)^2}{\frac{1}{2} + 2} = \frac{\frac{30}{4}}{\frac{5}{2}} = \frac{15}{2} \times \frac{2}{5} = 3$$

PTS: 2

REF: 082426ai

NAT: F.IF.A.2

TOP: Functional Notation

27 ANS:

x	-1	0	1	2	2
y	2	4	4	5	4

For every value of  $x$ , there is a unique value of  $y$ .

PTS: 2 REF: 082427ai NAT: F.IF.A.1 TOP: Defining Functions

28 ANS:

$$0.05(x - 3) = 0.35x - 7.5$$

$$x - 3 = 7x - 150$$

$$147 = 6x$$

$$24.5 = x$$

PTS: 2 REF: 082428ai NAT: A.REI.B.3 TOP: Solving Linear Equations

29 ANS:

$$x = \frac{-3 \pm \sqrt{(3)^2 - 4(1)(-6)}}{2(1)} = \frac{-3 \pm \sqrt{33}}{2}$$

PTS: 4 REF: 082429ai NAT: A.REI.B.4 TOP: Solving Quadratics

KEY: quadratic formula

30 ANS:

$$5x^3 - 80x = 5x(x^2 - 16) = 5x(x + 4)(x - 4)$$

PTS: 2 REF: 082430ai NAT: A.SSE.A.2

TOP: Factoring the Difference of Perfect Squares


31 ANS:

$$y = 15.13x - 959.63, 0.99, \text{strong}$$

PTS: 4 REF: 082431ai NAT: S.ID.B.6 TOP: Regression

KEY: linear with correlation coefficient

32 ANS:



; No, because  $2 > 3(2) - 4$  is false.

PTS: 4 REF: 082432ai NAT: A.REI.D.12 TOP: Graphing Systems of Linear Inequalities

33 ANS:

$$t = \frac{-64}{2(-16)} = 2 \quad h(2) = -16(2)^2 + 64(2) + 80 = -64 + 128 + 80 = 144 \quad (2, 144). \text{ At 2 seconds, the object is 144 feet}$$

above the ground.  $0 = -16t^2 + 64t + 80$

$$0 = t^2 - 4t - 5$$

$$0 = (t - 5)(t + 1)$$

$$t = 5$$

PTS: 4                      REF: 082433ai                      NAT: F.IF.B.4                      TOP: Graphing Quadratic Functions

KEY: key features

34 ANS:

$$x^2 + 4x - 1 = 2x + 7 \quad y = 2(-4) + 7 = -1 \quad (-4, -1), (2, 11)$$

$$x^2 + 2x - 8 = 0 \quad y = 2(2) + 7 = 11$$

$$(x + 4)(x - 2) = 0$$

$$x = -4, 2$$

PTS: 4                      REF: 082434ai                      NAT: A.REI.C.7                      TOP: Quadratic-Linear Systems

35 ANS:

$$2m + 3r = 38.5 \quad \text{Jen is not correct because the prices are } 6m + 9r = 115.5 \quad 2m + 3(8.5) = 38.5$$

$$6m + r = 47.5$$

$$6m + r = 47.5 \quad 2m + 25.5 = 38.5$$

$$8r = 68 \quad 2m = 13$$

$$r = 8.50 \quad m = 6.50$$

PTS: 6                      REF: 082435ai                      NAT: A.CED.A.3                      TOP: Modeling Linear Systems

## 0125AI

## Answer Section

1 ANS: 3

$$x^3 - 36x = x(x^2 - 36) = x(x + 6)(x - 6)$$

PTS: 2 REF: 012501ai NAT: A.SSE.A.2

TOP: Factoring the Difference of Perfect Squares

2 ANS: 1

$$m = \frac{8 - -2}{-1 - 4} = \frac{10}{-5} = -2 \quad y = mx + b$$

$$8 = -2(-1) + b$$

$$6 = b$$

PTS: 2 REF: 012502ai NAT: A.REI.D.10 TOP: Writing Linear Equations

KEY: slope-intercept form

3 ANS: 4

$$\frac{8}{2} = 4$$

PTS: 2 REF: 012503ai NAT: F.IF.A.3 TOP: Sequences

KEY: difference or ratio

4 ANS: 1 PTS: 2 REF: 012504ai NAT: A.SSE.A.1

TOP: Modeling Expressions

5 ANS: 2 PTS: 2 REF: 012505ai NAT: F.LE.B.5

TOP: Modeling Linear Functions

6 ANS: 2 PTS: 2 REF: 012506ai NAT: A.APR.A.1

TOP: Operations with Polynomials KEY: subtraction

7 ANS: 4 PTS: 2 REF: 012507ai NAT: A.REI.D.12

TOP: Graphing Systems of Linear Inequalities

8 ANS: 1

$$\frac{15 - 25}{3 - 1} = \frac{-10}{2} = -5 \quad a_{10} = 25 + (10 - 1)(-5) = 25 - 45 = -20$$

PTS: 2 REF: 012508ai NAT: F.BF.A.1 TOP: Sequences

KEY: explicit

9 ANS: 2

$$p = 2l + 2w$$

$$p - 2l = 2w$$

$$\frac{p - 2l}{2} = w$$

PTS: 2 REF: 012509ai NAT: A.CED.A.4 TOP: Transforming Formulas

10 ANS: 1

$$\frac{200}{300 + 200 + 80 + 25 + 120 + 105 + 100 + 70} = \frac{200}{1000} = 20\%$$

PTS: 2 REF: 012510ai NAT: S.ID.B.5 TOP: Frequency Tables

KEY: two-way

11 ANS: 1

$$-2(3x - 5) = \frac{9}{2}x - 2$$

$$-4(3x - 5) = 9x - 4$$

$$-12x + 20 = 9x - 4$$

$$24 = 21x$$

$$x = \frac{24}{21} = \frac{8}{7}$$

PTS: 2 REF: 012511ai NAT: A.REI.B.3 TOP: Solving Linear Equations

12 ANS: 3 PTS: 2 REF: 012512ai NAT: A.APR.A.1

TOP: Multiplication of Powers

13 ANS: 3 PTS: 2 REF: 012513ai NAT: F.LE.A.1

TOP: Families of Functions

14 ANS: 4 PTS: 2 REF: 012514ai NAT: A.REI.A.1

TOP: Identifying Properties

15 ANS: 4 PTS: 2 REF: 012515ai NAT: N.RN.B.3

TOP: Operations with Radicals

KEY: addition

16 ANS: 1 PTS: 2 REF: 012516ai NAT: S.ID.A.1

TOP: Dot Plots

17 ANS: 1 PTS: 2 REF: 012517ai NAT: F.IF.B.5

TOP: Domain and Range

KEY: graph

18 ANS: 2

$$4 - 1 = 2 \left( \frac{5}{4} + \frac{1}{4} \right)$$

$$3 = 3$$

PTS: 2 REF: 012518ai NAT: A.REI.D.10 TOP: Identifying Solutions

19 ANS: 4 PTS: 2 REF: 012519ai NAT: N.Q.A.1

TOP: Conversions

20 ANS: 3

77 78 81 84 86 88 93 95

79.5 90.5

$$90.5 - 79.5 = 11$$

PTS: 2 REF: 012520ai NAT: S.ID.A.2 TOP: Dispersion

KEY: basic

21 ANS: 4 PTS: 2 REF: 012521ai NAT: F.BF.B.3

TOP: Transformations with Functions

KEY: bimodalgraph

22 ANS: 4  
 $6.4 - 4x \geq -2.8$   
 $9.2 \geq 4x$   
 $2.3 \geq x$

PTS: 2 REF: 012522ai NAT: A.REI.B.3 TOP: Solving Linear Inequalities

23 ANS: 2  
 $x + x + 8 \geq 20$


PTS: 2 REF: 012523ai NAT: A.CED.A.1 TOP: Modeling Linear Inequalities

24 ANS: 4 PTS: 2 REF: 012524ai NAT: F.IF.C.7  
 TOP: Graphing Piecewise-Defined Functions

25 ANS:  
 5-6 minutes, as the speed remains at 35 mph during this interval.

PTS: 2 REF: 012525ai NAT: F.IF.B.4 TOP: Relating Graphs to Events

26 ANS:



PTS: 2 REF: 012526ai NAT: F.IF.C.7 TOP: Graphing Quadratic Functions

27 ANS:  
 7, as for each value of  $x$ , there is a unique value of  $y$ .

PTS: 2 REF: 012527ai NAT: F.IF.A.1 TOP: Defining Functions

28 ANS:  
 $x^2 + 8x + 16 = 33 + 16$   
 $(x + 4)^2 = 49$   
 $x + 4 = \pm 7$   
 $x = -11, 3$

PTS: 2 REF: 012528ai NAT: A.REI.B.4 TOP: Solving Quadratics  
 KEY: completing the square

29 ANS:

$$-22 = \frac{-3x - 5}{2}$$

$$-44 = -3x - 5$$

$$-39 = -3x$$

$$13 = x$$

PTS: 2 REF: 012529ai NAT: F.IF.A.2 TOP: Functional Notation

30 ANS:

$$\frac{4}{\sqrt{2}} \frac{\sqrt{2}}{\sqrt{2}} = \frac{4\sqrt{2}}{2} = 2\sqrt{2}$$

PTS: 2 REF: 012530ai NAT: N.RN.B.3 TOP: Operations with Radicals

KEY: division

31 ANS:

$$n + d = 25 \quad n + 9 = 25$$

$$5n + 10d = 170 \quad n = 16$$

$$5(25 - d) + 10d = 170$$

$$125 - 5d + 10d = 170$$

$$5d = 45$$

$$d = 9$$

PTS: 4 REF: 012531ai NAT: A.CED.A.3 TOP: Modeling Linear Systems

32 ANS:

$$y = 9.1x - 527.6, 0.9, \text{ strong relationship}$$

PTS: 4 REF: 012532ai NAT: S.ID.B.6 TOP: Regression

KEY: linear with correlation coefficient

33 ANS:

$$x = \frac{-4 \pm \sqrt{4^2 - 4(1)(-3)}}{2(1)} = \frac{-4 \pm \sqrt{28}}{2} = \frac{-4 \pm 2\sqrt{7}}{2} = -2 \pm \sqrt{7}$$

PTS: 4 REF: 012533ai NAT: A.REI.B.4 TOP: Solving Quadratics

KEY: quadratic formula

34 ANS:

$$x^2 - 7x + 12 = 2x - 6 \quad y = 2(6) - 6 = 6 \quad (6,6), (3,0)$$


$$x^2 - 9x + 18 = 0 \quad y = 2(3) - 6 = 0$$

$$(x - 6)(x - 3) = 0$$

$$x = 6, 3$$

PTS: 4 REF: 012534ai NAT: A.REI.C.7 TOP: Quadratic-Linear Systems

35 ANS:



$$2x + 1.5y = 30$$

$$y = 2x$$

hats.

(6,12) is the intersection, meaning Anna bought 6 balloons and 12

PTS: 6

REF: 012535ai

NAT: A.REI.C.6

TOP: Graphing Linear Systems

## 0625AI

## Answer Section

1 ANS: 3

$$\frac{10}{\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}} = \frac{10\sqrt{2}}{2} = 5\sqrt{2}$$

PTS: 2 REF: 062501ai NAT: N.RN.B.3 TOP: Operations with Radicals  
KEY: division

2 ANS: 2 PTS: 2 REF: 062502ai NAT: F.IF.C.7  
TOP: Graphing Quadratic Functions

3 ANS: 4 PTS: 2 REF: 062503ai NAT: F.LE.A.1  
TOP: Families of Functions

4 ANS: 3

	Andrea	Joe
mean	85.8	82.5
standard deviation	3.5	12.4

PTS: 2 REF: 062504ai NAT: S.ID.A.2 TOP: Central Tendency and Dispersion  
KEY: multiple data sets

5 ANS: 4 PTS: 2 REF: 062505ai NAT: A.SSE.A.1  
TOP: Modeling Expressions

6 ANS: 1 PTS: 2 REF: 062506ai NAT: A.APR.A.1  
TOP: Operations with Polynomials  
KEY: subtraction

7 ANS: 1

$$\frac{60 - 12}{12 - 4} = \frac{48}{8} = 6$$

PTS: 2 REF: 062507ai NAT: F.IF.B.6 TOP: Rate of Change

8 ANS: 4 PTS: 2 REF: 062508ai NAT: A.REI.A.1  
TOP: Identifying Properties

9 ANS: 2 PTS: 2 REF: 062509ai NAT: F.BF.B.3  
TOP: Transformations with Functions

10 ANS: 3 PTS: 2 REF: 062510ai NAT: A.REI.C.6  
TOP: Solving Linear Systems

11 ANS: 2 PTS: 2 REF: 062511ai NAT: A.REI.B.4  
TOP: Solving Quadratics  
KEY: completing the square

12 ANS: 1

$$IRQ = 16 - 12 = 14 - 10$$

PTS: 2 REF: 062512ai NAT: S.ID.A.1 TOP: Box Plots  
KEY: interpret

13 ANS: 3 PTS: 2 REF: 062513ai NAT: F.IF.B.5  
TOP: Domain and Range  
KEY: graph

14 ANS: 4 PTS: 2 REF: 062514ai NAT: N.RN.B.3  
TOP: Operations with Radicals  
KEY: classify

15 ANS: 3                      PTS: 2                      REF: 062515ai                      NAT: A.SSE.A.2  
TOP: Factoring the Difference of Perfect Squares

16 ANS: 1  
 $2\sqrt{27} + 4\sqrt{12} = 2\sqrt{9}\sqrt{3} + 4\sqrt{4}\sqrt{3} = 6\sqrt{3} + 8\sqrt{3} = 14\sqrt{3}$

PTS: 2                      REF: 062516ai                      NAT: N.RN.B.3                      TOP: Operations with Radicals  
KEY: addition

17 ANS: 1                      PTS: 2                      REF: 062517ai                      NAT: A.CED.A.1  
TOP: Modeling Linear Equations

18 ANS: 1  
 $g(-2) = \frac{2^{(-2)+3}}{(-2)^2 - 2} = \frac{2^1}{4-2} = 1$

PTS: 2                      REF: 062518ai                      NAT: F.IF.A.2                      TOP: Functional Notation  
19 ANS: 2                      PTS: 2                      REF: 062519ai                      NAT: F.IF.A.1  
TOP: Defining Functions

20 ANS: 1  
 $K = \frac{1}{2}mv^2$

$$2K = mv^2$$

$$m = \frac{2K}{v^2}$$

PTS: 2                      REF: 062520ai                      NAT: A.CED.A.4                      TOP: Transforming Formulas  
21 ANS: 4  
 $\frac{2(3x-1)}{3} = x+2$

$$6x - 2 = 3x + 6$$

$$3x = 8$$

$$x = \frac{8}{3}$$

PTS: 2                      REF: 062521ai                      NAT: A.REI.B.3                      TOP: Solving Linear Equations  
22 ANS: 1                      PTS: 2                      REF: 062522ai                      NAT: F.BF.A.1  
TOP: Sequences

23 ANS: 4  
 $x = \frac{-(-4)}{2(1)} = 2$

PTS: 2                      REF: 062523ai                      NAT: F.IF.C.9                      TOP: Comparing Quadratic Functions  
24 ANS: 3                      PTS: 2                      REF: 062524ai                      NAT: N.Q.A.1  
TOP: Conversions

25 ANS:

	Juniors	Seniors	Total
Watch Videos	42	23	65
Listen to Music	14	21	35
Total	56	44	100

PTS: 2 REF: 062525ai NAT: S.ID.B.5 TOP: Frequency Tables  
KEY: two-way

26 ANS:

$$5(2 - y) > -11y - 8$$

$$10 - 5y > -11y - 8$$

$$6y > -18$$

$$y > -3$$

PTS: 2 REF: 062526ai NAT: A.REI.B.3 TOP: Solving Linear Inequalities

27 ANS:

$$(5x - 3)(-2x + 7) = -10x^2 + 35x + 6x - 21 = -10x^2 + 41x - 21$$

PTS: 2 REF: 062527ai NAT: A.APR.A.1 TOP: Operations with Polynomials  
KEY: multiplication

28 ANS:

$$d = \frac{-2 - (-20)}{4 - 1} = \frac{18}{3} = 6 \quad a_8 = -20 + (8 - 1)6 = 22$$

PTS: 2 REF: 062528ai NAT: F.BF.A.1 TOP: Sequences

29 ANS:

$$y - 5 = -3(x + 2)$$

$$y = -3x - 6 + 5$$

$$y = -3x - 1$$


PTS: 3 REF: 062529ai NAT: A.REI.D.10 TOP: Writing Linear Equations  
KEY: slope-intercept form

30 ANS:

$$x(x^2 - 36) = x(x + 6)(x - 6)$$

PTS: 2 REF: 062530ai NAT: A.SSE.A.2  
TOP: Factoring the Difference of Perfect Squares

31 ANS:



-2 and -1

PTS: 4 REF: 062531ai NAT: A.REI.D.11 TOP: Quadratic-Linear Systems

32 ANS:

$$x = \frac{-2 \pm \sqrt{2^2 - 4(6)(-1)}}{2(6)} = \frac{-2 \pm \sqrt{4 + 24}}{12} = \frac{-2 \pm \sqrt{28}}{12} = \frac{-2 \pm 2\sqrt{7}}{12} = \frac{-1 \pm \sqrt{7}}{6}$$

PTS: 4 REF: 062532ai NAT: A.REI.B.4 TOP: Solving Quadratics

KEY: quadratic formula

33 ANS:

$$y = -23.67x + 1216, -0.99, \text{strong}$$

PTS: 4 REF: 062533ai NAT: S.ID.B.6 TOP: Regression

KEY: linear with correlation coefficient

34 ANS:

$$x^2 + 9x + 4 = 2x - 6 \quad y = 2(-5) - 6 = -16 \quad (-5, -16), (-2, -10)$$


$$x^2 + 7x + 10 = 0 \quad y = 2(-2) - 6 = -10$$

$$(x + 5)(x + 2) = 0$$

$$x = -5, -2$$

PTS: 4 REF: 062534ai NAT: A.REI.C.7 TOP: Quadratic-Linear Systems

35 ANS:



$$6x + 12y \geq 120$$

$$x + y \leq 14$$

8 hours of babysitting and 6 hours of tutoring.  $6(8) + 12(6) \geq 120$ 

$$8 + 6 \leq 14$$

PTS: 6 REF: 062535ai NAT: A.CED.A.3 TOP: Modeling Systems of Linear Inequalities

## 0825AI

## Answer Section

- 1 ANS: 3                   PTS: 2                   REF: 082501ai           NAT: A.SSE.A.2  
TOP: Factoring the Difference of Perfect Squares
- 2 ANS: 3                   PTS: 2                   REF: 082502ai           NAT: A.CED.A.1  
TOP: Modeling Linear Equations
- 3 ANS: 2  
 $g(-3) = -2(-3)^2 + 16 = -18 + 16 = -2$
- PTS: 2                   REF: 082503ai           NAT: F.IF.A.2           TOP: Functional Notation
- 4 ANS: 2  
 $x^2 - 8x - 20 = 0$   
 $(x - 10)(x + 2) = 0$   
 $x = 10, -2$
- PTS: 2                   REF: 082504ai           NAT: A.APR.B.3           TOP: Zeros of Polynomials
- 5 ANS: 1  
 $3(-2)^2 - \frac{1}{4}(-2) + 3 = 12 + \frac{1}{2} + 3 = 15.5$
- PTS: 2                   REF: 082505ai           NAT: A.REI.D.10       TOP: Identifying Solutions
- 6 ANS: 3  
 $x^2 = 8x - 15$   
 $x^2 - 8x + 15 = 0$   
 $(x - 3)(x - 5) = 0$   
 $x = 3, 5$
- PTS: 2                   REF: 082506ai           NAT: A.REI.D.11       TOP: Quadratic-Linear Systems
- 7 ANS: 4                   PTS: 2                   REF: 082507ai           NAT: A.SSE.A.1  
TOP: Modeling Expressions
- 8 ANS: 4                   PTS: 2                   REF: 082508ai           NAT: A.REI.A.1  
TOP: Identifying Properties
- 9 ANS: 4  
 $f(x) = (x - 1)^2 - 3; g(x) = |x + 1|; h(x) = -4(2)^x; j(x) = 4x + 1$
- PTS: 2                   REF: 082509ai           NAT: F.LE.A.1           TOP: Families of Functions

10 ANS: 2

$$3x^2 - 24x = 0$$

$$3x(x - 8) = 0$$

$$x = 0, 8$$

PTS: 2 REF: 082510ai NAT: A.REI.B.4 TOP: Solving Quadratics

KEY: factoring

11 ANS: 2

$$\frac{2.5 - 10}{3 - 1} = \frac{-7.5}{2} = -3.75$$

PTS: 2 REF: 082511ai NAT: F.IF.B.6 TOP: Rate of Change

12 ANS: 1

PTS: 2

REF: 082512ai

NAT: S.ID.C.8

TOP: Correlation Coefficient

13 ANS: 4

PTS: 2

REF: 082513ai

NAT: F.BF.B.3

TOP: Transformations with Functions

14 ANS: 2

$$\text{mean: } \frac{0 + 4 + 12 + 12 + 12 + 10}{20} = \frac{50}{20} = 2.5, \text{ mode: } 2, \text{ median: } \frac{2 + 2}{2} = 2$$

PTS: 2 REF: 082514ai NAT: S.ID.A.1 TOP: Dot Plots

15 ANS: 1

PTS: 2

REF: 082515ai

NAT: F.IF.A.2

TOP: Domain and Range

16 ANS: 3

PTS: 2

REF: 082516ai

NAT: A.APR.A.1

TOP: Powers of Powers

17 ANS: 1

$$2A = h(b_1 + b_2)$$

$$\frac{2A}{b_1 + b_2} = h$$

PTS: 2 REF: 082517ai NAT: A.CED.A.4 TOP: Transforming Formulas

18 ANS: 1

$$f(0) = 5, g(0) = 5, h(0) = -3$$

PTS: 2 REF: 082518ai NAT: F.IF.C.9 TOP: Comparing Functions

19 ANS: 3

$$(x + 7)^2 + (x - 3)^2 = x^2 + 14x + 49 + x^2 - 6x + 9 = 2x^2 + 8x + 58$$

PTS: 2 REF: 082519ai NAT: A.APR.A.1 TOP: Operations with Polynomials

KEY: multiplication

20 ANS: 1

$$2\sqrt{10} \cdot 3\sqrt{2} = 6\sqrt{20} = 6\sqrt{4} \sqrt{5} = 12\sqrt{5}$$

PTS: 2 REF: 082520ai NAT: N.RN.B.3 TOP: Operations with Radicals

KEY: multiplication

- 21 ANS: 3                   PTS: 2                   REF: 082521ai           NAT: A.APR.A.1  
TOP: Operations with Polynomials           KEY: subtraction
- 22 ANS: 4                   PTS: 2                   REF: 082522ai           NAT: F.IF.A.1  
TOP: Defining Functions
- 23 ANS: 1                   PTS: 2                   REF: 082523ai           NAT: A.REI.C.6  
TOP: Solving Linear Systems
- 24 ANS: 4                   PTS: 2                   REF: 082524ai           NAT: N.Q.A.1  
TOP: Conversions

25 ANS:

$$\frac{1}{6}(4x + 12) = 9$$

$$4x + 12 = 54$$


$$4x = 42$$

$$x = 10.5$$

- PTS: 2                   REF: 082525ai           NAT: A.REI.B.3           TOP: Solving Linear Equations
- 26 ANS:  
Irrational because the sum can not be written as the ratio of two integers.

PTS: 2                   REF: 082526ai           NAT: N.RN.B.3           TOP: Operations with Radicals  
KEY: classify

27 ANS:



- PTS: 2                   REF: 082527ai           NAT: F.IF.C.7           TOP: Graphing Absolute Value Functions
- 28 ANS:

	Brand A	Brand B	Total
Adults	40	59	99
Teenagers	32	49	81
Total	72	108	180

- PTS: 2                   REF: 082528ai           NAT: S.ID.B.5           TOP: Frequency Tables  
KEY: two-way
- 29 ANS:  
 $a_8 = 5(3)^{8-1} = 10935$

PTS: 2                   REF: 082529AI           NAT: F.BF.A.1           TOP: Sequences

30 ANS:

$$x^2 + 14x = 28$$


$$x^2 + 14x + 49 = 28 + 49$$

$$(x + 7)^2 = 77$$

PTS: 2 REF: 082530ai NAT: A.REI.B.4 TOP: Solving Quadratics

KEY: completing the square

31 ANS:

(0,4),  $x = 0$ 

PTS: 2 REF: 082531sai NAT: F.IF.C.7 TOP: Graphing Quadratic Functions

32 ANS:


$$7.5x + 18 \leq 78 \quad 9 \text{ hours}$$

$$7.5x \leq 60$$

$$x \leq 8$$

PTS: 2 REF: 082532ai NAT: A.CED.A.1 TOP: Modeling Linear Inequalities

33 ANS:



$$(0,0) \quad 0 \geq -\frac{1}{2}(0) - 3 \quad y - 2x < 5$$

$$0 \geq -3 \quad 0 - 2(0) < 5$$

$$0 < 5$$

PTS: 4 REF: 082533ai NAT: A.REI.D.12 TOP: Graphing Systems of Linear Inequalities

34 ANS:

$$x = \frac{6 \pm \sqrt{(-6)^2 - 4(1)(3)}}{2(1)} = \frac{6 \pm \sqrt{24}}{2} = \frac{6 \pm 2\sqrt{6}}{2}$$

PTS: 4 REF: 082534ai NAT: A.REI.B.4 TOP: Solving Quadratics

KEY: quadratic formula

35 ANS:

$$\begin{array}{ll} x + y = 25 & y = 25 - x \quad y = 25 - 10 = 15 \quad 2.25x + 1.5(4) \leq 20 \quad 6 \text{ hot dogs} \\ 2.25x + 1.5y = 45 & 2.25x + 1.5(25 - x) = 45 \quad 2.25x + 6 \leq 20 \\ & 2.25x + 37.5 - 1.5x = 45 \quad 2.25x \leq 14 \\ & .75x = 7.5 \quad x \leq 6.\bar{2} \\ & x = 10 \end{array}$$

PTS: 6 REF: 082535ai NAT: A.CED.A.3 TOP: Modeling Linear Systems