

Algebra II Regents at Random Worksheets

- 1 The transportation methods used by the upperclassmen at Calhoun High School are summarized in the table below.

Upperclassmen Transportation Methods			
	Drive	Take the Bus	Walk
Junior	58	75	12
Senior	81	39	12

Are the events "being a junior" and "driving to school" independent? Using statistical evidence, justify your answer.

- 2 Which equation is equivalent to $P = 210x^{\frac{4}{3}}y^{\frac{7}{3}}$
- $P = \sqrt[3]{210x^4y^7}$
 - $P = 70xy^2\sqrt[3]{xy}$
 - $P = 210xy^2\sqrt[3]{xy}$
 - $P = 210xy^2\sqrt[3]{x^3y^5}$
- 3 A survey was given to 1250 randomly selected high school students at the end of their junior year. The survey offered four post-graduation options: two-year college, four-year college, military, or work. Of the 1250 responses, 475 chose a four-year college. State *one* possible conclusion that can be made about the population of high school juniors, based on this survey.
- 4 The heights of the members of a ski club are normally distributed. The average height is 64.7 inches with a standard deviation of 4.3 inches. Determine the percentage of club members, to the nearest percent, who are between 67 inches and 72 inches tall.
- 5 Over the set of integers, factor the expression $2x^4 - 10x^3 + 3x^2 - 15x$ completely.
- 6 Solve the equation $x^2 + 3x + 11 = 0$ algebraically. Express the answer in $a + bi$ form.
- 7 Solve algebraically for all values of x :

$$\frac{8}{x+5} - \frac{3}{x} = 5$$
- 8 Jay is training for a bike race over fifteen weeks. At the end of the first week, he has ridden ten miles, and he is planning to increase his weekly distance by nine percent each week. Approximately how many miles total will he have ridden from the beginning of his training to the end of the fifteenth week?
- 10.989
 - 33.417
 - 163.5
 - 293.609
- 9 Given the functions $f(x) = 2x + \frac{5}{2}$ and $g(x) = \frac{3}{x}$, what are the solutions to $f(x) = g(x)$?
- (0.75, 4) or (-2, -1.5)
 - $x = 0.75$ or $x = -2$
 - $y = -1.5$ or $y = 4$
 - (-2, 0.75)
- 10 The crew aboard a small fishing boat caught 350 pounds of fish on Monday. From that Monday through the end of the week on Friday, the weight of the fish caught increased 15% per day. The total weight, in pounds, of fish caught is approximately
- 411
 - 612
 - 1748
 - 2360

- 11 Which value, to the *nearest tenth*, is an approximate solution for the equation $f(x) = g(x)$,

if $f(x) = \frac{5}{x-3}$ and $g(x) = 2(1.3)^x$?

- 1) 3.2
- 2) 3.9
- 3) 4.0
- 4) 5.6

- 12 A brewed cup of coffee contains 130 mg of caffeine. The half-life of caffeine in the bloodstream is 5.5 hours. Write a function, $C(t)$ to represent the amount of caffeine in the bloodstream t hours after drinking one cup of coffee.

- 13 Beginning July 1, 2019, Michelle deposited \$250 into an account that yields 0.15% each month. She continued to make \$250 deposits into this account on the first of each month for 3 years. Which expression represents the amount of money that was in the account after her last deposit was made on June 1, 2022?

- 1) $250(1.0015)^3$
- 2) $250(1.0015)^{36}$
- 3) $\frac{250 - 250(1.0015)^3}{1 - 1.0015}$
- 4) $\frac{250 - 250(1.0015)^{36}}{1 - 1.0015}$

- 14 What is the value of y for the system shown below?

$$3x + 4y - 5z = -27$$

$$2x + 3y - z = -3$$


$$6x - y + 4z = 3$$

- 1) -27
- 2) 6
- 3) 3
- 4) -3

- 15 The solution set of the equation $x - 1 = \sqrt{2x + 6}$ is

- 1) $\{5, -1\}$
- 2) $\{5\}$
- 3) $\{-1\}$
- 4) $\{ \}$

- 16 On the graph below, draw at least one complete cycle of a sine graph passing through point $(0, 2)$ that has an amplitude of 3, a period of π , and a midline at $y = 2$.



Based on your graph, state an interval in which the graph is increasing.

- 17 Given $a > 1$, use the properties of rational exponents to determine the value of x for the equation below.

$$\frac{\sqrt[5]{a^{10}}}{\left(a^3\right)^{\frac{1}{2}}} = a^x$$

- 18 The expression $8^{\frac{x}{2}} \cdot 8^{\frac{x}{3}}$ is equivalent to


- 1) $\sqrt[6]{8^{5x}}$
- 2) $64^{\frac{5x}{6}}$
- 3) $\sqrt[5]{8^{2x}}$
- 4) $64^{\frac{x^2}{6}}$

- 19 A researcher wants to determine if nut allergies and milk allergies are related to each other. The researcher surveyed 1500 people and asked them if they are allergic to nuts or milk. The survey results are summarized in the table below.

	Allergic to Nuts	Not Allergic to Nuts
Allergic to Milk	3	42
Not Allergic to Milk	12	1443

Determine the probability that a randomly selected survey respondent is allergic to milk. Determine the probability that a randomly selected survey respondent is allergic to milk, given that the person is allergic to nuts. Based on the survey data, determine whether nut allergies and milk allergies are independent events. Justify your answer.

- 20 The explicit formula $a_n = 6 + 6n$ represents the number of seats in each row in a movie theater, where n represents the row number. Rewrite this formula in recursive form.
- 21 Graph $y = f(x)$, where $f(x) = \log_2(x - 1) + 3$ on the set of axes below.



State the equation of the asymptote of $f(x)$. When $f(x)$ is reflected over the line $y = x$, a new function is formed: $g(x) = 2^{x-3} + 1$. State the equation of the asymptote of $g(x)$.

- 22 Factor $x^3 + 4x^2 - 9x - 36$ completely.

- 23 Is $x + 3$ a factor of $7x^3 + 27x^2 + 9x - 27$? Justify your answer.
- 24 In a small city, there are 22 gas stations. The mean price for a gallon of regular gas was \$2.12 with a standard deviation of \$0.05. The distribution of the data was approximately normal. Given this information, the middle 95% of the gas stations in this small city likely charge
- 1) \$1.90 to \$2.34 for a gallon of gas
 - 2) \$1.97 to \$2.27 for a gallon of gas
 - 3) \$2.02 to \$2.22 for a gallon of gas
 - 4) \$2.07 to \$2.17 for a gallon of gas
- 25 Jin solved the equation $\sqrt{4-x} = x + 8$ by squaring both sides. What extraneous solution did he find?
- 1) -5
 - 2) -12
 - 3) 3
 - 4) 4
- 26 Which expression is equivalent to $x^8 - y^8$?
- 1) $(x - y)^8$
 - 2) $(x^2 + y^2)^2(x^2 - y^2)^2$
 - 3) $(x^4 + y^4)(x^2 + y^2)(x + y)(x - y)$
 - 4) $(x + y)^4(x - y)^4$
- 27 The expression $i^2(5x - 2i)^2$ is equivalent to
- 1) $-25x^2 + 20xi - 4$
 - 2) $-25x^2 + 20xi + 4$
 - 3) $25x^2 + 20xi + 4$
 - 4) $25x^2 + 4$

28 The exact value of $\sin\left(\frac{8\pi}{3}\right)$ is

- 1) $\frac{1}{2}$
- 2) $-\frac{1}{2}$
- 3) $-\frac{\sqrt{3}}{2}$
- 4) $\frac{\sqrt{3}}{2}$

29 To prepare for lacrosse tryouts, Kole is increasing the amount of time he spends at the gym. This week he is spending 150 minutes there and he plans to increase this amount by 2% each week. The amount of time, in minutes, that he plans to spend at the gym t weeks from now is given by the function $A(t) = 150(1.02)^t$. In terms of a daily growth rate, the amount of time Kole is planning to spend at the gym can best be modeled by the function

- 1) $A(t) = 150(1.14869)^{\frac{t}{7}}$
- 2) $A(t) = 150(1.14869)^{7t}$
- 3) $A(t) = 150(1.00283)^{\frac{t}{7}}$
- 4) $A(t) = 150(1.00283)^{7t}$

30 The equation of the parabola that has its focus at the point $(-3, 2)$ and directrix at $y = 0$ is

- 1) $y = \frac{1}{4}(x+3)^2 + 1$
- 2) $y = \frac{1}{4}(x-3)^2 + 1$
- 3) $y = \frac{1}{8}(x+3)^2 + 1$
- 4) $y = \frac{1}{8}(x-3)^2 + 1$

31 Describe the translations that map $f(x) = \log x$ to $g(x) = \log(x+3) - 5$.

32 What are the solutions to $4x^2 - 7x - 2 = -10$

- 1) $-\frac{1}{4}, 2$
- 2) $\frac{7}{8} \pm \frac{\sqrt{79}}{8}i$
- 3) $\frac{7}{8} \pm \frac{\sqrt{241}}{8}$
- 4) $\frac{7}{8} \pm \frac{\sqrt{143}}{8}i$

33 The cost of a brand-new electric-hybrid vehicle is listed at \$33,400, and the average annual depreciation for the vehicle is 15%. The car's value can be modeled by the function

$V(x) = 33,400(0.85)^x$, where x represents the years since purchase. Julia and Jacob have each written a function that is equivalent to the original.

Jacob's function: $V(x) = 33,400(0.1422)^{\frac{1}{12}x}$

Julia's function: $V(x) = 33,400(0.9865)^{12x}$

Whose function is correctly rewritten to reveal the approximate monthly depreciation rate? Justify your answer.

34 Given $\tan \theta = -\frac{4}{3}$ where $\frac{\pi}{2} < \theta < \pi$, what is the value of $\sec \theta$?

- 1) $-\frac{5}{3}$
- 2) $-\frac{3}{5}$
- 3) $\frac{4}{5}$
- 4) $\frac{5}{3}$

35 Solve the following system of equations algebraically for all values of x , y , and z :

$$3x - 8y + 2z = -60$$

$$2x - 7y - 5z = -31$$

$$-6x + 2y - 4z = 36$$


- 36 The number of bacteria in a sample, which can be modeled by an exponential regression, is shown in the table below.

Time Since Observation Began (hours)	0	1	2	3.5	4
Number of Bacteria	40	48	57	75	82

Assuming this trend continues, approximately how many bacteria would be present 8 hours after the observation began?

- | | |
|--------|--------|
| 1) 123 | 3) 168 |
| 2) 127 | 4) 180 |

- 37 A sporting goods manufacturer is trying to determine if they should continue to produce multiple types of hockey pucks. The company surveyed 50 randomly chosen customers and asked them if they purchased both game regulation pucks and lighter training pucks. Of those surveyed, 40 of them said that they purchase both types of pucks. A simulation that was run 100 times based on the survey results produced the approximately normal results below.



- Determine an interval containing the middle 95% of plausible values that estimates the proportion of all customers who would purchase both types of pucks from the company.
- The company will continue to manufacture both types of hockey pucks if it is reasonable to assume that the true proportion of customers who buy both types of hockey pucks is above 0.60. Using the interval from part *a*, explain whether or not the company should continue to produce both types of hockey pucks.

- 38 Which expression is equivalent to $2xy^2\sqrt[3]{x^2y}$?

- 1) $2x^{\frac{5}{3}}y^{\frac{7}{3}}$
- 2) $2xy$
- 3) $2x^{\frac{2}{3}}y^{\frac{2}{3}}$
- 4) $2x^7y^4$

- 39 The graph of which function has a period of 3?

- 1) $y = -7 \sin\left(\frac{2\pi}{3}x\right) - 5$
- 2) $y = -7 \sin\left(\frac{3\pi}{2}x\right) + 9$
- 3) $y = -7 \sin(3x) - 5$
- 4) $y = 3 \sin(\pi x) + 9$


- 40 What is the remainder when $4x^3 - 3x + 3$ is divided by $x - 2$?

- 1) -23
- 2) -7
- 3) 13
- 4) 29

- 41 The solution to the equation $6(2^{x+4}) = 36$ is

- 1) -1
- 2) $\frac{\ln 36}{\ln 12} - 4$
- 3) $\ln(3) - 4$
- 4) $\frac{\ln 6}{\ln 2} - 4$

- 42 A grocery store orders 50 bags of oranges from a company's distribution center. The bags have a mean weight of 3.85 pounds per bag. The company claims that their bags of oranges have a mean weight of 4 pounds. The grocery store ran a simulation of 50 bags, 2500 times, assuming a mean of 4 pounds. The results are shown below.



Is the mean weight of the grocery store's sample unusual? Explain using the results of the simulation.


- 43 What is the value of x in the solution of the system of equations below?
- $$5x + 2y - z = -14$$
- $$7y - z = 31$$
- $$5y + 4z - 5x = -23$$
- 1) -17
 - 2) 2
 - 3) $-\frac{1}{5}$
 - 4) -7
- 44 Given $f(x) = x^4 + x^3 - 3x^2 + 9x - 108$ and $f(3) = 0$, which values satisfy $f(x) = 0$?
- 1) -4, 3 only
 - 2) -3, 4 only
 - 3) $\pm 3i, -4, 3$
 - 4) $\pm 3i, -3, 4$
- 45 The equations $y = 3t + 6$ and $y = (1.82)^t$ approximately model the growth of two separate populations where $t > 0$. What is the best approximation of the time, t , at which the populations are the same?
- 1) -1.9
 - 2) 0.3
 - 3) 5.1
 - 4) 21.3
- 46 A family owned grocery store in New Hartford, NY employs 49 people whose ages are approximately normally distributed with a mean of 36 years and a standard deviation of 6.2 years. Ryan has been hired to work at this store. He is 30 years old. How many people who work at this store would you expect to be younger than Ryan?
- 1) 17
 - 2) 7
 - 3) 41
 - 4) 8
- 47 The solution set for the equation $x + 1 = \sqrt{4x + 25}$ is
- 1) $\{ \}$
 - 2) $\{6\}$
 - 3) $\{6, -4\}$
 - 4) $\{-4\}$
- 48 In an attempt to get the student body's opinion of a new dress code, members of the statistics class surveyed the students of the first period computer science class. Explain a statistical bias in the method of data collection.

- 49 Markus is a long-distance walker. In one race, he walked 55 miles in t hours and in another race walked 65 miles in $t + 3$ hours. His rates are shown in the equations below.

$$r = \frac{55}{t} \quad r = \frac{65}{t+3}$$

Markus walked at an equivalent rate, r , for each race. Determine the number of hours that each of the two races took.

- 50 For the function $d(x) = \sqrt[3]{x+2}$, the inverse function, $d^{-1}(x)$, equals
- $\sqrt[3]{x+2}$
 - $x^3 + 2$
 - $-\sqrt[3]{x+2}$
 - $x^3 - 2$
- 51 In the town of Skaneateles, New York, house prices since 2008 have changed based on the function $H(t) = 200,000(1.045)^t$, where t is the number of years since 2008 and $H(t)$ is the median house price. Determine the average rate of change for the median house price in Skaneateles, from 2010 to 2018 to the *nearest dollar per year*. Explain what this rate of change means as it relates to median house prices.
- 52 If $f(x)$ is represented by the graph below, which translation of $f(x)$ would have imaginary roots?



- $f(x+5)$
- $f(x-5)$
- $f(x)+5$
- $f(x)-5$

- 53 Given $f(x) = x^3 - 3$ and $f^{-1}(x) = \sqrt[3]{x-3b}$, the value of b is

- 1
- 1
- 3
- 3

- 54 The function $d(t) = 2\cos\left(\frac{\pi}{6}t\right) + 5$ models the

water depth, in feet, at a location in a bay, t hours since the last high tide. Determine the *minimum* water depth of the location, in feet, and justify your answer.

- 55 Given $x > 0$, the expression $\left(\frac{1}{x^{-2}}\right)^{-\frac{3}{4}}$ is equivalent

to

- $x\sqrt{x}$
- $\frac{1}{x\sqrt{x}}$
- $\sqrt[3]{x^2}$
- $\frac{1}{\sqrt[3]{x^2}}$

- 56 For which equations will the value $s = 4$ make the statement an identity?

I $(2x - 3)^2 = 4x^2 - 3sx + 9$

II $(x - 2)^3 = (x - 2)(x^2 + sx + s)$


- I, only
- II, only
- I and II
- neither I nor II

- 57 Solve $3.8e^{1.5t} = 16$ algebraically for t to the *nearest hundredth*.

- 58 Can $f(x) = x^3 + 7$ be classified as an odd function? Justify your answer.

- 59 Determine if $x + 4$ is a factor of $2x^3 + 10x^2 + 4x - 16$. Explain your answer.


- 60 In a packaging plant, a machine packs boxes with jars. The machine's manufacturer states that a box is packed, on average, every 42 seconds. To test that claim, the packaging plant randomly selects a sample of 10 boxes and finds the sample mean to be 49.8 seconds. The company ran a simulation of 1000 trials based on the manufacturer's claim. The approximately normal results are shown below.



Based on the simulation, determine an interval containing the middle 95% of plausible mean times. Round your answer to the *nearest hundredth*. Is the time 49.8 seconds unusual? Use statistical evidence to justify your answer.

- 61 Sketch a graph of polynomial $P(x)$, given the criteria below:

- $P(x)$ has zeros only at -5 , 1 , and 4
- As $x \rightarrow \infty, P(x) \rightarrow -\infty$
- As $x \rightarrow -\infty, P(x) \rightarrow -\infty$



- 62 The expression $\frac{4x^2 - 5}{x^2 - 1}$ is equivalent to

- 1) $4 - \frac{1}{x^2 - 1}$
- 2) $4 + \frac{1}{x^2 - 1}$
- 3) $4 - \frac{9}{x^2 - 1}$
- 4) $4 - \frac{4}{x^2 - 1}$

- 63 Given the equation $S(x) = 1.7 \sin(bx) + 12$, where the period of $S(x)$ is 12, what is the value of b ?

- 1) $\frac{\pi}{6}$
- 2) 24π
- 3) $\frac{\pi}{12}$
- 4) 6π

- 64 A random sample of 152 students was surveyed on a particular day about how they got to school. The survey results are summarized in the table below.

	Attendance Status	
	Late	On-Time
Method of Transportation		
Car	6	24
Bus	20	80
Walk	4	18


Which statement is best supported by the data?

- 1) The probability of being late given that a student walked is greater than the probability that a student walked given that the student was late.
- 2) The probability of being late given that a student walked is less than the probability that a student walked given that the student was late.
- 3) The probability of being late given that a student walked is equal to the probability that a student walked given that the student was late.
- 4) The probability of being late given that a student walked cannot be determined.
- 65 Mr. Zachary posts review assignments on the Betamath website for his students. On his last test, 49% of his students used Betamath and passed. Overall, 68% of his students used Betamath. Approximately what percentage of Mr. Zachary's students passed, given that they used Betamath?
- 1) 19%
- 2) 32%
- 3) 33%
- 4) 72%
- 66 If $f(x) = \sqrt[3]{x} + 4$, then $f^{-1}(x)$ equals
- 1) $\sqrt[3]{x-4}$
- 2) $(x-4)^3$
- 3) $x^3 + \frac{1}{4}$
- 4) $-\sqrt[3]{x} - 4$
- 67 Given i is the imaginary unit, which expression is equivalent to $5i(2x + 3i) - x\sqrt{-9}$?
- 1) $15 + 13xi$
- 2) $-15 + 13xi$
- 3) $15 + 7xi$
- 4) $-15 + 7xi$
- 68 Solve algebraically for x : $\frac{1}{2x} - \frac{5}{6} = \frac{3}{x}$
- 69 Given $y = -2x$ and $x^2 + y^2 = 5$, the point of intersection in Quadrant II is
- 1) $(1, -2)$
- 2) $(-2, 1)$
- 3) $(-1, 1)$
- 4) $(-1, 2)$
- 70 The average cost of a gallon of milk in the United States between the years of 1995 and 2018 can be modeled by the equation $P(t) = -0.0004t^3 + 0.0114t^2 - 0.0150t + 2.6602$, where $P(t)$ represents the cost, in dollars, and t is time in years since January 1995. During this time period, in what year did $P(t)$ reach its maximum?
- 1) 1995
- 2) 2013
- 3) 2014
- 4) 2018
- 71 Given $f(x) = 2x^2 + 7x - 15$ and $g(x) = 3 - 2x$, what is $\frac{f(x)}{g(x)}$ for all defined values?
- 1) $-x - 5$
- 2) $-x + 5$
- 3) $x - 5$
- 4) $x + 5$

- 72 The volume of a cardboard box can be modeled by $V(x)$, which is the product of the length, width, and height, x . If the length can be represented by $L(x) = 18 - 2x$ and the width can be represented by $W(x) = 18 - 2x$, then which function represents $V(x)$?

- 1) $V(x) = 4x^2 - 72x + 324$
- 2) $V(x) = 4x^3 - 72x^2 + 324x$
- 3) $V(x) = -3x + 36$
- 4) $V(x) = 4x^3 + 324x$

- 73 Which equation is graphed in the diagram below?



- 1) $y = -3 \sin\left(\frac{\pi}{3}x\right) + 5$
- 2) $y = -3 \cos\left(\frac{\pi}{3}x\right) + 5$
- 3) $y = -5 \sin\left(\frac{\pi}{3}x\right) + 3$
- 4) $y = -5 \cos\left(\frac{\pi}{3}x\right) + 3$

- 74 Express $(2xi^3 - 3y)^2$ in simplest form.

- 75 Given $\cos \theta = -\frac{2}{7}$ with θ in Quadrant II, find the exact value of $\sin \theta$.

- 76 Given $p \neq q$, $p = \left(\frac{1}{2}\right)^q$, expressed in logarithmic form, is equivalent to

- 1) $\log_p\left(\frac{1}{2}\right) = q$
- 2) $\log_q(p) = \frac{1}{2}$
- 3) $\log_{\frac{1}{2}}(p) = q$
- 4) $\log_{\frac{1}{2}}(q) = p$

- 77 Consider the function $f(x) = 2^x$. Is $f(x)$ an even function? Justify your answer. Write an equation for $g(x)$, the function that results after $f(x)$ is shifted up 5 units. Write an equation for $h(x)$, the inverse of $g(x)$.

- 78 Which expression is equivalent to

$$(x - 2)^2 + 27(x - 2) - 90?$$

- 1) $(x + 30)(x - 3)$
- 2) $(x + 28)(x - 5)$
- 3) $(x - 30)(x + 3)$
- 4) $(x - 2)(x + 25)(x - 90)$

- 79 A culture of 1000 bacteria triples every 10 hours. Which expression models the number of bacteria in the sample after t hours?

- 1) $1000e^{3t}$
- 2) $1000(3)^t$
- 3) $1000(3)^{10t}$
- 4) $1000(3)^{\frac{t}{10}}$

- 80 The graph of $y = 2^x - 4$ is positive on which interval?

- 1) $(-\infty, \infty)$
- 2) $(2, \infty)$
- 3) $(0, \infty)$
- 4) $(-4, \infty)$

81 The number of employees who work nights and weekends at a department store is summarized in the table below.

	Works Nights	Doesn't Work Nights
Works Weekends	8	40
Doesn't Work Weekends	12	60


Let N represent the event "works nights" and let W represent the event "works weekends." Based on the table, are N and W independent events?

- 1) Yes, because $P(N) \cdot P(W) = P(N \cap W)$. 3) No, because $P(N) \cdot P(W) = P(N \cap W)$.
 2) Yes, because $P(N) \cdot P(W) \neq P(N \cap W)$. 4) No, because $P(N) \cdot P(W) \neq P(N \cap W)$.

82 If $4(10^{5x-2}) = 12$ then x equals

- 1) $\frac{2.3}{5}$
 2) $\frac{1}{3} \left(\frac{\log 12}{\log 40} + 5 \right)$
 3) $\frac{\log(3) + 2}{5}$
 4) $\frac{1}{5} \left(\frac{\log 12}{\log 4} + 2 \right)$

83 On the axes below, graph $y = 3.2(1.8)^x$.



84 A research assistant receives a first year salary of \$90,000 and a 2% annual raise throughout the first ten years of employment. In total, how much money will be earned over the first ten years, to the nearest dollar?

- 1) \$91,837
 2) \$109,709
 3) \$877,917
 4) \$985,475

85 What is the focus of the parabola

$$8(y + 2) = (x + 5)^2?$$

- 1) $(-5, 0)$
 2) $(-5, -4)$
 3) $(5, 0)$
 4) $(5, 4)$


86 The parabola with equation $12(y + 1) = (x - 4)^2$ has

- 1) a vertex at $(4, 2)$
 2) a focus at $(4, -1)$
 3) a directrix $y = -4$
 4) four units between the focus and vertex

87 Abby is told that each day there is a 50% chance it will rain. Which simulation can Abby perform to determine the likelihood of it raining for the next seven days?

- 1) Flip a coin seven times, count how many heads, and repeat 50 times.
 2) Roll a die seven times, count how many twos, and repeat 50 times.
 3) Roll a pair of dice, count totals of seven, and repeat 50 times.
 4) Flip a coin 50 times and count how many heads.

88 Which graph has imaginary roots?



89 The seventh term of the geometric sequence $\sqrt{6}, -2\sqrt{3}, 2\sqrt{6}, -4\sqrt{3}, \dots$ is

- 1) $6\sqrt{6}$
- 2) $-6\sqrt{3}$
- 3) $8\sqrt{6}$
- 4) $-8\sqrt{3}$

90 Factored completely, $x^4 + 4x^3 - 9x^2 - 36x$ is equivalent to

- 1) $x(x+9)(x-9)(x+4)$
- 2) $x(x+3)(x-3)(x+4)$
- 3) $(x^3 - 9x)(x+4)$
- 4) $x(x^2 - 9)(x+4)(x+4)$

91 A manufacturer claims that the number of ounces of a beverage dispensed by one of its automatic dispensers is normally distributed with a mean of 8.0 ounces and a standard deviation of 0.04 ounces. To the *nearest tenth of a percent*, what percent of the cups filled by this company's dispenser will contain between 7.9 and 8.11 ounces?

- 1) 99.5
- 2) 99.4
- 3) 99.1
- 4) 97.6

92 For all positive values of x , which expression is equivalent to $\sqrt{x} \cdot \sqrt[4]{x^{11}}$?

- 1) $x^{\frac{19}{22}}$
- 2) $x^{\frac{11}{8}}$
- 3) $x^{\frac{13}{4}}$
- 4) $x^{\frac{2}{11}}$

93 The monthly unemployment rate of towns in the United States is approximately normally distributed with a mean rate of 5.2% and a standard deviation of 1.6%. Determine the percentage of towns, to the *nearest integer*, that have a monthly unemployment rate greater than 6%.

94 Given $q(x) = 2 \log(x)$ and $r(x) = (x-2)^3 - 4$, what is a solution of $q(x) = r(x)$ to the *nearest tenth*?

- 1) 1.1
- 2) 3.7
- 3) 3.9
- 4) 4.3

95 A grocery store owner wonders how many customers bring reusable bags to the store. An employee stands at the store entrance for two hours and counts the number of people bringing in reusable bags. This type of study is best classified as

- 1) a census
- 2) an experiment
- 3) an observational study
- 4) a survey

- 96 The population of China, in millions, can be modeled by the function $P(x) = 316.93e^{0.0133x}$, where x is the number of years since 1900. The population of India since 1900 is summarized in the table below:

Years since 1900	Population (millions)
0	243
10	254
20	268
30	285
40	324
50	376.3
60	450.6
70	555.1
80	699
90	873.3
100	1056.6
110	1234.3
120	1380

Which country's population had a greater average rate of change between 1950 and 2020? Justify your answer.

- 97 Tim deposits \$300 into a savings account. The annual interest rate is 2.7% and compounds monthly. He uses the equation
- $$A = 300 \left(1 + \frac{0.027}{12} \right)^{12t}$$
- to determine how much money he will have after t years. Which equation is equivalent to Tim's equation?

- 1) $A = 300 \left[(1.00225)^{12} \right]^t$
- 2) $A = 300(0.08558)^{12t}$
- 3) $A = 300 \left[1 + \left(\frac{0.027}{12} \right)^{12t} \right]$
- 4) $A = (300)^{12t} (1)^{12t} + \left(\frac{0.027}{12} \right)^{12t}$

- 98 Write a recursive formula for the sequence 8, 20, 50, 125, 312.5, ...

- 99 When $\left(\frac{1}{\sqrt[3]{y^2}} \right) y^4$ is written in the form y^n , what is the value of n ? Justify your answer.

- 100 The solution of $\frac{x}{x+3} + \frac{2}{x-4} = \frac{2x+27}{x^2-x-12}$ is

- 1) -3
- 2) -7
- 3) 3
- 4) 7

- 101 The probability of having math homework is $\frac{1}{3}$ and the probability of having English homework is $\frac{1}{7}$. The probability of having math homework or having English homework is $\frac{9}{21}$. What is the probability of having math homework and having English homework?

- 1) $\frac{19}{21}$
- 2) $\frac{1}{5}$
- 3) $\frac{1}{21}$
- 4) $\frac{10}{21}$

- 102 The solutions to the equation $3x^2 - 4x + 2 = 2x - 3$ are

- 1) $\frac{2}{3} \pm \frac{\sqrt{2}}{3}i$
- 2) $1 \pm \frac{\sqrt{6}}{3}i$
- 3) $1 \pm \frac{\sqrt{12}}{3}$
- 4) $1 \pm 2\sqrt{6}i$

- 103 If $x - 5$ is a factor of

$p(x) = ax^4 + bx^3 + cx^2 + dx + e$, then which statement must be true?

- 1) $p(-5) = 0$
- 2) $p(-5) \neq 0$
- 3) $p(5) = 0$
- 4) $p(5) \neq 0$

- 104 At the Lakeside Resort, the probability that a guest room has a view of the lake is 0.24. The probability that a guest room has a queen-size bed is 0.74. Let A be the event that the guest room has a view of the lake, and let B be the event that the guest room has a queen-size bed. Events A and B are found to be independent of each other. Determine the exact probability that a randomly selected guest room has a view of the lake and a queen-size bed. Determine the exact probability that a randomly selected guest room has a view of the lake or a queen-size bed.





- 105 A study of the red tailed hawk population in a given area shows the population, $H(t)$, can be represented by the function $H(t) = 50(1.19)^t$ where t represents the number of years since the study began. In terms of the monthly rate of growth, the population can be best approximated by the function

- 1) $H(t) = 50(1.015)^{12t}$
- 2) $H(t) = 50(1.15)^{\frac{t}{12}}$
- 3) $H(t) = 50(1.19)^{12t}$
- 4) $H(t) = 50(1.19)^{\frac{t}{12}}$


- 106 The roots of the equation $0 = x^2 + 6x + 10$ in simplest $a + bi$ form are

- 1) $-3 \pm 2i$
- 2) $-6 \pm i$
- 3) $-3 \pm i$
- 4) $-3 \pm i\sqrt{2}$

- 107 Which graph best represents the graph of $f(x) = (x + a)^2(x - b)$, where a and b are positive real numbers?



108 Two surveys were conducted to estimate the proportion of teens who use social media at least once per day.



Based on these results, it was determined that approximately 75% of teens use social media at least once per day. What is the best explanation of the difference in the results between the two surveys?

- | | |
|---|--|
| 1) The smaller sample size of five teens resulted in a smaller margin of error and should provide a more accurate estimate. | 3) The larger sample size of 50 teens resulted in a smaller margin of error and should provide a more accurate estimate. |
| 2) The smaller sample size of five teens resulted in a bigger margin of error and should provide a more accurate estimate. | 4) The larger sample size of 50 teens resulted in a bigger margin of error and should provide a more accurate estimate. |

109 A company wishes to determine the cooking time for one pound of spaghetti. The company's technicians cooked one pound of spaghetti and recorded the time needed for the spaghetti to be ready to eat. Repeating this process 35 times resulted in an approximately normal distribution, with a mean of 9.82 minutes and a standard deviation of 1.4 minutes. In which interval should the middle 95% of cooking times fall?

- 1) (8.42, 11.22)
- 2) (7.02, 12.62)
- 3) (9.35, 10.29)
- 4) (6.82, 11.32)

110 Which equation is true for all real values of x ?

- 1) $x^4 + x = (x + 1)(x^3 - x^2 + x)$
- 2) $x^4 + x = (x + 1)(x^3 + x)$
- 3) $x^4 + x = (x^2 + x)^2$
- 4) $x^4 + x = (x - 1)(x^3 + x^2 + x)$

111 A teacher randomly divides all of her students into two groups. She grades the homework for one group but does not grade the homework for the other group. All homework is returned to the students. She then compares test scores for each of the groups to see if grading homework has an effect on the tests cores. This method of data collection is best described as


- 1) an experiment
- 2) an unbiased survey
- 3) a simulation
- 4) an observational study

112 Which expression is equivalent to

$$(x + 3)^2 + 4(x + 3) - 5?$$

- 1) $(x + 5)(x - 1)$
- 2) $(x + 8)(x + 2)$
- 3) $(x - 2)(x + 4)$
- 4) $x^2 + 4x + 16$


- 113 The populations of honeybees in two different colonies are studied for four months. During this time, the colony population can be approximated by $P(t) = P_0 e^{rt}$, where $P(t)$ is the colony population of bees at t months, P_0 is the initial population, and r is the growth rate. Colony A has an initial population of 10,000 bees and a continuous growth rate of 0.25. Colony B has an initial population of 6000 bees and a continuous growth rate of 0.45. Write functions for both $A(t)$ and $B(t)$ that model the honeybee populations of the colonies after t months. Graph $A(t)$ and $B(t)$ for $0 \leq t \leq 4$.



State, to the *nearest tenth of a month*, when the colonies will have the same population. Determine algebraically how long it will take, to the *nearest tenth of a month*, for the population in Colony A to triple.

- 114 The asymptote of the graph of $f(x) = 5 \log(x + 4)$ is
- 1) $y = 6$
 - 2) $x = -4$
 - 3) $x = 4$
 - 4) $y = 5$

- 115 The profit function, $p(x)$, is found by subtracting the cost function, $c(x)$, from the revenue function, $r(x)$. Which function below represents the cost function given $p(x) = -15x^2 + 600x + 60$ and $r(x) = -0.4x^2 + 130x + 1200$?
- 1) $c(x) = -14.6x^2 + 470x - 1140$
 - 2) $c(x) = -14.6x^2 + 730x - 1260$
 - 3) $c(x) = 14.6x^2 - 470x + 1140$
 - 4) $c(x) = 14.6x^2 + 730x - 1260$
- 116 Algebraically find the zero of $c(x) = x^3 + 2x^2 - 16x - 32$. On the axes below, sketch $y = c(x)$.




- 117 During the summer, Adam saved \$4000 and Betty saved \$3500. Adam deposited his money in Bank A at an annual rate of 2.4% compounded monthly. Betty deposited her money in Bank B at an annual rate of 4% compounded quarterly. Write two functions that represent the value of each account after t years if no other deposits or withdrawals are made, where Adam's account value is represented by $A(t)$, and Betty's by $B(t)$. Using technology, determine, to the *nearest tenth of a year*, how long it will take for the two accounts to have the same amount of money in them. Justify your answer.
- 118 Solve the system of equations shown below algebraically:

$$(x - 4)^2 + (y - 1)^2 = 9$$

$$x - y = 6$$

- 119 The graph below shows the amount of a radioactive substance left over time.




The daily rate of decay over an 8-day interval is approximately

- 1) 23%
 - 2) 95%
 - 3) 5%
 - 4) 77%
- 120 Megan is performing an experiment in a lab where the air temperature is a constant 73°F and the liquid is 237°F . One and a half hours later, the temperature of the liquid is 112°F . Newton's law of cooling states $T(t) = T_a + (T_0 - T_a)e^{-kt}$ where:
- $T(t)$: temperature, $^{\circ}\text{F}$, of the liquid at t hours
- T_a : air temperature
- T_0 : initial temperature of the liquid
- k : constant
- Determine the value of k , to the *nearest thousandth*, for this liquid. Determine the temperature of the liquid using your value for k , to the *nearest degree*, after two and a half hours. Megan needs the temperature of the liquid to be 80°F to perform the next step in her experiment. Use your value for k to determine, to the *nearest tenth of an hour*, how much time she must wait since she first began the experiment.
- 121 Given x is a real number, write the expression in simplest $a + bi$ form: $(x + 2i)(3 - 2xi) + 2x^2i$

122 Algebraically solve for x : $2x = 6 + 2\sqrt{x - 1}$

- 123 Determine an equation for the parabola with focus $(-2, 4)$ and directrix $y = 10$. (The use of the grid below is optional.)



- 124 Taylor wants to open an investment account with the \$1200 she received for her birthday. She has narrowed her choices down to two banks. America's Bank offers 6.4% annual interest compounded quarterly. Barnyard Bank offers 6.35% annual interest compounded continuously. Write functions for $A(t)$ and $B(t)$ to represent the value of her investment with America's Bank and Barnyard Bank as a function of time, t , in years. Taylor would like to invest the \$1200 into one bank for ten years making no additional deposits and no withdrawals. With which bank will Taylor earn the most money? Justify your answer. Taylor chooses to invest her money in Barnyard Bank. Algebraically determine how long, to the *nearest tenth of a year*, it will take her initial investment to triple assuming she makes no deposits or withdrawals.
- 125 An initial investment of \$1000 reaches a value, $V(t)$, according to the model $V(t) = 1000(1.01)^{4t}$, where t is the time in years. Determine the average rate of change, to the *nearest dollar per year*, of this investment from year 2 to year 7.

- 126 The cost, in dollars, of a single-ride fare in the New York City subway in the years since 1904 is listed in the table below.

Years since 1904 (x)	0	49	72	91	99	111
Fare (y)	\$0.05	\$0.15	\$0.50	\$1.50	\$2.00	\$2.75

Which equation best models the cost of a single-ride fare based on these data?

- 1) $y = 0.0375(1.0392)^x$ 3) $y = 0.0234x - 0.487$
 2) $y = 1.0392(0.0375)^x$ 4) $y = -0.179 + 0.356\ln(x)$
- 127 Functions f and g are given below.

$$f(x) = \frac{7}{2}x^2 - 5x + 11$$

$$g(x) = 3x^2 - 7x + 25$$
 When $2f(x)$ is subtracted from $g(x)$, the result is
 1) $4x^2 - 3x - 3$
 2) $-4x^2 + 3x + 3$
 3) $4x^2 - 17x - 47$
 4) $-4x^2 - 17x + 47$
- 128 The temperature, F , in degrees Fahrenheit, after t hours of a roast put into an oven is given by the equation $F = 325 - 185e^{-0.4t}$. What was the temperature of the roast when it was put into the oven?
 1) 325
 2) 200
 3) 185
 4) 140
- 129 Which point is in Quadrant III and is a solution to the system below?

$$y = x^2 - 24$$


$$y = x - 12$$

 1) (4, -8)
 2) (-3, -15)
 3) (-4, -16)
 4) (-3, -33)
- 130 Which equation does *not* represent an identity?
 1) $x^2 - y^2 = (x + y)(x - y)$
 2) $(x - y)^2 = (x - y)(x - y)$
 3) $(x + y)^2 = x^2 + 2xy + y^2$
 4) $(x + y)^3 = x^3 + 3xy + y^3$
- 131 Which expression is equivalent to $\frac{6x^4 + 4x^3 + x + 200}{x + 2}$?
 1) $6x^2 - 8x + 17 + \frac{166}{x + 2}$
 2) $6x^2 + 16x + 33 + \frac{266}{x + 2}$
 3) $6x^3 + 16x^2 + 32x + 65 + \frac{330}{x + 2}$
 4) $6x^3 - 8x^2 + 16x - 31 + \frac{262}{x + 2}$
- 132 Which statements must be true about the polynomial function $k(x) = -2x^3 - 11x^2 - 12x + 9$?
 I. $(x - 3)$ is a factor of $k(x)$
 II. $k(0) = 9$
 III. $\frac{k(x)}{x + 2}$ has a remainder of 5
 1) II, only
 2) I and II
 3) II and III
 4) I, II, and III
- 133 Given $f(x) = 2x^3 - 3x^2 - 5x - 12$ and $g(x) = x - 3$, the quotient of $\frac{f(x)}{g(x)}$ is
 1) $2x^2 + 3x + 4$
 2) $2x^3 + 3x^2 + 4x$
 3) $2x^2 - 9x + 22 - \frac{78}{x - 3}$
 4) $2x^3 - 9x^2 + 22x - 78$

- 134 A manufacturer of sweatshirts finds that profits and costs fluctuate depending on the number of products created. Creating more products doesn't always increase profits because it requires additional costs, such as building a larger facility or hiring more workers. The manufacturer determines the profit, $p(x)$, in thousands of dollars, as a function of the number of sweatshirts sold, x , in thousands. This function, p , is given below.


$$p(x) = -x^3 + 11x^2 - 7x - 69$$

Graph $y = p(x)$, over the interval $0 \leq x \leq 9$, on the set of axes below.



Over the given interval, state the coordinates of the maximum of p and round all values to the *nearest integer*. Explain what this point represents in terms of the number of sweatshirts sold and profit. Determine how many sweatshirts, to the *nearest whole sweatshirt*, the manufacturer would need to produce in order to first make a positive profit. Justify your answer.

- 135 Given 3 is a root of $f(x) = x^4 - x^3 - 21x^2 + 45x$, what are the other unique roots of $f(x)$?
- 1) -5, only
 - 2) -5 and 0
 - 3) -3, 1 and 5
 - 4) -5, -3 and 0
- 136 Graph *at least one* cycle of $y = 5 \sin(4x) - 3$ on the set of axes below.




- 137 To solve the equation $\frac{7}{x+7} + \frac{4x}{x-7} = \frac{3x+7}{x-7}$, Joan's first step is to multiply both sides by the least common denominator. Which statement is true?
- 1) -14 is an extraneous solution.
 - 2) 7 and -7 are extraneous solutions.
 - 3) 7 is an extraneous solution.
 - 4) There are no extraneous solutions.
- 138 If θ is an angle in standard position whose terminal side passes through the point $(-3, -4)$, which statement is true?
- 1) $\sec \theta > 0$ and $\tan \theta > 0$
 - 2) $\sec \theta < 0$ and $\tan \theta < 0$
 - 3) $\sec \theta > 0$ and $\tan \theta < 0$
 - 4) $\sec \theta < 0$ and $\tan \theta > 0$

139 Which function has a greater average rate of change on the interval $[-1, 4]$? Justify your answer.

x	$m(x)$
-2	-3
-1	1
0	1
1	3
2	13
3	37
4	81
5	151

$$p(x) = 3^x + 1$$

140 The height of a running trail is modeled by the quartic function $y = f(x)$ shown below, where x is the distance in miles from the start of the trail and y is the height in feet relative to sea level.



If this trail has a minimum height of 16 feet below sea level, which function(s) could represent a running trail whose minimum height is half of the minimum height of the original trail?

I. $y = f\left(\frac{1}{2}x\right)$ II. $y = f(x) + 8$ III. $y = \frac{1}{2}f(x)$

- 1) I, only
- 2) II, only
- 3) I and III
- 4) II and III

141 An initial investment of \$5000 in an account earns 3.5% annual interest. Which function correctly represents a recursive model of the investment after n years?

- 1) $A = 5000(0.035)^n$
- 2) $a_0 = 5000$
 $a_n = a_{n-1}(0.035)$
- 3) $A = 5000(1.035)^n$
- 4) $a_0 = 5000$
 $a_n = a_{n-1}(1.035)$

142 Robert is buying a car that costs \$22,000. After a down payment of \$4000, he borrows the remainder from a bank, a six year loan at 6.24% annual interest rate. The following formula can be used to calculate his monthly loan payment.

$$R = \frac{(P)(i)}{1 - (1 + i)^{-t}}$$

R = monthly payment

P = loan amount





i = monthly interest rate

t = time, in months

Robert's monthly payment will be

- 1) \$298.31
- 2) \$300.36
- 3) \$307.35
- 4) \$367.10

- 143 Which graph could represent a 4th degree polynomial function with a positive leading coefficient, 2 real zeros, and 2 imaginary zeros?



- 144 The rational expression $\frac{2x^4 - 5x^2 + 3x - 2}{x - 3}$ is equivalent to

- 1) $2x^3 - 5x - 12 - \frac{38}{x - 3}$
- 2) $2x^3 + 6x^2 + 13x + 42 + \frac{124}{x - 3}$
- 3) $2x^3 - 5x + 18 - \frac{56}{x - 3}$
- 4) $2x^3 - 6x^2 + 13x - 36 + \frac{106}{x - 3}$

- 145 A cafeteria food manager studied the lunchtime eating habits of a group of employees in their office building. The purpose of the study was to determine the proportion of employees who purchased lunch in the cafeteria, brought their lunch from home, or purchased lunch from an outside vendor. This collection of data would best be classified as

- 1) a census
- 2) an experiment
- 3) an observational study
- 4) a simulation

- 146 Given $\sin \theta = \frac{7}{25}$ and θ terminates in quadrant II, what is the value of $\tan \theta$?

- 1) $-\frac{7}{24}$
- 2) $-\frac{24}{7}$
- 3) $\frac{7}{24}$
- 4) $\frac{24}{7}$

- 147 The height, above ground, of a Ferris wheel car can be modeled by the function

$$h(t) = -103.5 \cos\left(\frac{2\pi t}{5}\right) + 108.5 \text{ where } h \text{ is}$$

measured in feet and t is measured in minutes. State the period of the function and describe what the period represents in this context.

Algebra II Regents at Random Worksheets

148 Consider the data in the table below.

x	1	2	3	4	5	6
y	3.9	6	11	18.1	28	40.3


State an exponential regression equation to model these data, rounding all values to the *nearest thousandth*.

149 The value of an automobile t years after it was purchased is given by the function

$$V = 38,000(0.84)^t.$$
 Which statement is true?

- 1) The value of the car increases 84% each year.
- 2) The value of the car decreases 84% each year.
- 3) The value of the car increases 16% each year.
- 4) The value of the car decreases 16% each year.

150 The graph of a cubic polynomial function $p(x)$ is shown below.



If $p(x)$ is written as a product of linear factors, which factor would appear twice?

- 1) $x - 2$
- 2) $x + 2$
- 3) $x - 3$
- 4) $x + 3$

151 For $x \neq 0$ and $y \neq 0$, $\sqrt[3]{81x^{15}y^9} = 3^a x^5 y^3$. Determine the value of a .

152 If the focus of a parabola is $(0,6)$ and the directrix is $y = 4$, what is an equation for the parabola?

- 1) $y^2 = 4(x - 5)$
- 2) $x^2 = 4(y - 5)$
- 3) $y^2 = 8(x - 5)$
- 4) $x^2 = 8(y - 6)$

153 Determine the solution of $\sqrt{3x+7} = x-1$ algebraically.

154 The monthly high temperature ($^{\circ}\text{F}$) in Buffalo, New York can be modeled by $B(m) = 24.9 \sin(0.5m - 2.05) + 55.25$, where m is the number of the month and January = 1. Find the average rate of change in the monthly high temperature between June and October, to the *nearest hundredth*. Explain what this value represents in the given context.

155 What is the solution set of the equation

$$\frac{x+2}{x} + \frac{x}{3} = \frac{2x^2+6}{3x}?$$

- 1) $\{-3\}$
- 2) $\{-3, 0\}$
- 3) $\{3\}$
- 4) $\{0, 3\}$

156 Algebraically determine the solution set for the system of equations below.

$$y = 2x^2 - 7x + 4$$

$$y = 11 - 2x$$

- 157 The table below shows the food preferences of sports fans whose favorite sport is football or baseball.


Favorite Food to Eat While Watching Sports

	Wings	Pizza	Hot Dogs
Football	14	20	6
Baseball	6	12	42

The probability that a fan prefers pizza given that the fan prefers football is

- $$\begin{array}{ll} 1) & \frac{1}{2} \\ 2) & \frac{1}{5} \end{array} \qquad \begin{array}{ll} 3) & \frac{5}{8} \\ 4) & \frac{13}{25} \end{array}$$

- 158 The graph of a quadratic function is shown below.



When the graph of $x + y = 4$ is drawn on the same axes, one solution to this system is

- 1) $(4,0)$
- 2) $(1,5)$
- 3) $(2,2)$
- 4) $(3,1)$

- 159 According to a study done at a hospital, the average weight of a newborn baby is 3.39 kg, with a standard deviation of 0.55 kg. The weights of all the newborns in this hospital closely follow a normal distribution. Last year, 9256 babies were born at this hospital. Determine, to the *nearest integer*, approximately how many babies weighed more than 4 kg.

- 160 Given x and y are positive, which expressions are equivalent to $\frac{x^3}{y}$?

$$\text{I. } \left(\frac{y}{x^3}\right)^{-1} \quad \text{II. } \sqrt[3]{x^9}(y^{-1}) \quad \text{III. } \frac{x^6\sqrt[4]{y^8}}{x^3y^3}$$

- 1) I and II, only
- 2) I and III, only
- 3) II and III, only
- 4) I, II, and III

- 161 Which statement regarding polynomials and their zeros is true?

- 1) $f(x) = (x^2 - 1)(x + a)$ has zeros of 1 and $-a$, only.
- 2) $f(x) = x^3 - ax^2 + 16x - 16a$ has zeros of 4 and a , only.
- 3) $f(x) = (x^2 + 25)(x + a)$ has zeros of ± 5 and $-a$.
- 4) $f(x) = x^3 - ax^2 - 9x + 9a$ has zeros of ± 3 and a .

- 162 The heights of the students at Central High School can be modeled by a normal distribution with a mean of 68.1 and a standard deviation of 3.4 inches. According to this model, approximately what percent of the students would have a height less than 60 inches or greater than 75 inches?

- 1) 0.86%
- 2) 1.26%
- 3) 2.12%
- 4) 2.98%


163 Consider the following patterns:

- I. $16, -12, 9, -6.75, \dots$
- II. $1, 4, 9, 16, \dots$
- III. $6, 18, 30, 42, \dots$
- IV. $\frac{1}{2}, \frac{2}{3}, \frac{3}{4}, \frac{4}{5}, \dots$

Which pattern is geometric?

- 1) I
- 2) II
- 3) III
- 4) IV

164 Marissa and Sydney are trying to determine if there is enough interest in their school to put on a senior musical. They randomly surveyed 100 members of the senior class and 43% of them said they would be interested in being in a senior musical. Marissa and Sydney then conducted a simulation of 500 more surveys, each of 100 seniors, assuming that 43% of the senior class would be interested in being in the musical. The output of the simulation is shown below.



The standard deviation of the simulation is closest to

- 1) 0.02
- 2) 0.05
- 3) 0.09
- 4) 0.43

165 Given $f(x) = x^4 - x^3 - 6x^2$, for what values of x will $f(x) > 0$?

- 1) $x < -2$, only
- 2) $x < -2$ or $x > 3$
- 3) $x < -2$ or $0 \leq x \leq 3$
- 4) $x > 3$, only

166 Audra is interested in studying the number of students entering kindergarten in the Ahlville Central School District over the next several years. Using data dating back to 2015, she determines that the number of kindergarteners is decreasing at an exponential rate. She creates a formula to model this situation $y = a(b)^x$, where x is the number of years since 2015 and y is the number of students entering kindergarten. If there were 105 students entering kindergarten in Ahlville in 2015, which statement about Audra's formula is true?

- 1) a is positive and b is negative.
- 2) a is negative and b is positive.
- 3) Both a and b are positive.
- 4) Both a and b are negative.

167 Show why $x - 3$ is a factor of $m(x) = x^3 - x^2 - 5x - 3$. Justify your answer.

168 Consider the function $f(x) = 2x^3 + x^2 - 18x - 9$. Which statement is true?

- 1) $2x - 1$ is a factor of $f(x)$.
- 2) $x - 3$ is a factor of $f(x)$.
- 3) $f(3) \neq f\left(-\frac{1}{2}\right)$
- 4) $f\left(\frac{1}{2}\right) = 0$

169 Chet has \$1200 invested in a bank account modeled by the function $P(n) = 1200(1.002)^n$, where $P(n)$ is the value of his account, in dollars, after n months. Chet's debt is modeled by the function $Q(n) = 100n$, where $Q(n)$ is the value of debt, in dollars, after n months. After n months, which function represents Chet's net worth, $R(n)$?

- 1) $R(n) = 1200(1.002)^n + 100n$
- 2) $R(n) = 1200(1.002)^{12n} + 100n$
- 3) $R(n) = 1200(1.002)^n - 100n$
- 4) $R(n) = 1200(1.002)^{12n} - 100n$

170 Write $\frac{x\sqrt{x^3}}{\sqrt[3]{x^5}}$ as a single term in simplest form, with a rational exponent.

Algebra II Regents Exam Questions at Random Worksheet # 25

- 171 A cup of coffee is left out on a countertop to cool. The table below represents the temperature, $F(t)$, in degrees Fahrenheit, of the coffee after it is left out for t minutes.

t	0	5	10	15	20	25
F(t)	180	144	120	104	93.3	86.2

Based on these data, write an exponential regression equation, $F(t)$, to model the temperature of the coffee. Round all values to the *nearest thousandth*.

- 172 Write the expression $A(x) \bullet B(x) - 3C(x)$ as a polynomial in standard form.

$$A(x) = x^3 + 2x - 1$$

$$B(x) = x^2 + 7$$

$$C(x) = x^4 - 5x$$

- 173 A population is normally distributed with a mean of 23 and a standard deviation of 1.2. The percentage of the population that falls below 21, to the *nearest hundredth*, is

- 1) 0.05
- 2) 4.78
- 3) 8.29
- 4) 91.30

- 174 The depth of the water, $d(t)$, in feet, on a given day at Thunder Bay, t hours after midnight is modeled

$$\text{by } d(t) = 5 \sin\left(\frac{\pi}{6}(t-5)\right) + 7. \text{ Which statement}$$

about the Thunder Bay tide is *false*?

- 1) A low tide occurred at 2 a.m.
- 2) The maximum depth of the water was 12 feet.
- 3) The water depth at 9 a.m. was approximately 11 feet.
- 4) The difference in water depth between high tide and low tide is 14 feet.

- 175 The sum of the first 20 terms of the series $-2 + 6 - 18 + 54 - \dots$ is

- 1) -610
- 2) -59
- 3) 1,743,392,200
- 4) 2,324,522,934

- 176 The George family would like to borrow \$45,000 to purchase a new boat. They qualified for a loan with an annual interest rate of 6.75%. The monthly loan payment can be found using the formula below.

$$M = \frac{P\left(\frac{r}{12}\right)\left(1 + \frac{r}{12}\right)^n}{\left(1 + \frac{r}{12}\right)^n - 1}$$

M = monthly payment

P = amount borrowed

r = annual interest rate

n = number of monthly payments

What is the monthly payment if they would like to pay off the loan in five years?

- 1) \$262.99
- 2) \$252.13
- 3) \$915.24
- 4) \$885.76

- 177 Which equation represents a polynomial identity?

- 1) $x^3 + y^3 = (x + y)^3$
- 2) $x^3 + y^3 = (x + y)(x^2 - xy + y^2)$
- 3) $x^3 + y^3 = (x + y)(x^2 - xy - y^2)$
- 4) $x^3 + y^3 = (x - y)(x^2 + xy + y^2)$

- 178 Given that i is the imaginary unit, the expression $(x - 2i)^2$ is equivalent to


- 1) $x^2 + 4$
- 2) $x^2 - 4$
- 3) $x^2 - 2xi - 4$
- 4) $x^2 - 4xi - 4$

Algebra II Regents Exam Questions at Random Worksheet # 26

179 Which expression is equivalent to $\frac{x^3 - 2}{x - 2}$?

- 1) x^2
- 2) $x^2 + 2x + 4 + \frac{6}{x - 2}$
- 3) $x^2 - 2$
- 4) $x^2 - 2x + 4 - \frac{10}{x - 2}$

180 The function $v(x) = x(3 - x)(x + 4)$ models the volume, in cubic inches, of a rectangular solid for $0 \leq x \leq 3$. Graph $y = v(x)$ over the domain $0 \leq x \leq 3$.



To the *nearest tenth of a cubic inch*, what is the maximum volume of the rectangular solid?

181 Sonja is cutting wire to construct a mobile. She cuts 100 inches for the first piece, 80 inches for the second piece, and 64 inches for the third piece. Assuming this pattern continues, write an explicit equation for a_n , the length in inches of the n th piece. Sonja only has 40 feet of wire to use for the project and wants to cut 20 pieces total for the mobile using her pattern. Will she have enough wire? Justify your answer.

182 Written in simplest form, the fraction $\frac{x^3 - 9x}{9 - x^2}$,

where $x \neq \pm 3$, is equivalent to

- 1) $-x$
- 2) x
- 3) $\frac{-x(x + 3)}{(3 + x)}$
- 4) $\frac{x(x - 3)}{(3 - x)}$


183 Mrs. Favata's statistics class wants to conduct a survey to see how students feel about changing the school mascot's name. Which plan is the best process for gathering an appropriate sample?

- 1) Survey students in a random sample of senior homerooms.
- 2) Survey every tenth student entering art classes in the school.
- 3) Survey every fourth student entering the cafeteria during each lunch period.
- 4) Survey all members of the school's varsity sports teams.


184 For $f(x) = \cos x$, which statement is true?

- 1) $2f(x)$ and $f(2x)$ are even functions.
- 2) $f(2x)$ and $f(x) + 2$ are odd functions.
- 3) $2f(x)$ and $f\left(x + \frac{\pi}{2}\right)$ are odd functions.
- 4) $f(x) + 2$ is an odd function and $f\left(x + \frac{\pi}{2}\right)$ is an even function.

185 Determine the average rate of change, in mph, from 2 to 4 hours on the graph shown below.



- 186 The volume of air in an average lung during breathing can be modeled by the graph below.




Using the graph, write an equation for $N(t)$, in the form $N(t) = A \sin(Bt) + C$. That same lung, when engaged in exercise, has a volume that can be modeled by $E(t) = 2000 \sin(\pi t) + 3200$, where $E(t)$ is volume in mL and t is time in seconds. Graph *at least one* cycle of $E(t)$ on the same grid as $N(t)$. How many times during the 5-second interval will $N(t) = E(t)$?

- 187 For all values of x for which the expression is

defined, $\frac{x^2 + 3x}{x^2 + 5x + 6}$ is equivalent to

- 1) $1 - \frac{x}{x+2}$
- 2) $\frac{x}{x+2}$
- 3) $\frac{3x}{5x+6}$
- 4) $1 + \frac{1}{2x+6}$

- 188 An angle, θ , is rotated counterclockwise on the unit circle, with its terminal side in the second quadrant, as shown in the diagram below.



Which value represents the radian measure of angle θ ?

- 1) 1
 - 2) 2
 - 3) 65.4
 - 4) 114.6
- 189 How many real solutions exist for the system of equations below?

$$y = \frac{1}{4}x - 8$$

$$y = \frac{1}{2}x^2 + 2x$$

- 1) 1
 - 2) 2
 - 3) 3
 - 4) 0
- 190 A public radio station held a fund-raiser. The table below summarizes the donor category and method of donation.

		Donor Category	
		Supporter	Patron
Method of Donation	Phone calls	400	672
	Online	1200	2016

To the *nearest thousandth*, find the probability that a randomly selected donor was categorized as a supporter, given that the donation was made online. Do these data indicate that being a supporter is independent of donating online? Justify your answer.

Algebra II Regents Exam Questions at Random Worksheet # 28

- 191 If $f(x) = 2x^4 - x^3 - 16x + 8$, then $f\left(\frac{1}{2}\right)$
- 1) equals 0 and $2x + 1$ is a factor of $f(x)$
 - 2) equals 0 and $2x - 1$ is a factor of $f(x)$
 - 3) does not equal 0 and $2x + 1$ is not a factor of $f(x)$
 - 4) does not equal 0 and $2x - 1$ is a factor of $f(x)$

- 192 Which expression is equivalent to $(x + yi)(x^2 - xyi - y^2)$, where i is the imaginary unit?

- 1) $x^3 + y^3i$
- 2) $x^3 - xy^2 - (xy^2 + y^3)i$
- 3) $x^3 - 2xy^2 - y^3i$
- 4) $x^3 - y^3i$

- 193 If θ is an angle in standard position whose terminal side passes through the point $(-2, -3)$, what is the numerical value of $\tan \theta$?


- 1) $\frac{2}{3}$
- 2) $\frac{3}{2}$
- 3) $-\frac{2}{\sqrt{13}}$
- 4) $-\frac{3}{\sqrt{13}}$

- 194 Which statement about data collection is most accurate?


- 1) A survey about parenting styles given to every tenth student entering the library will provide unbiased results.
- 2) An observational study allows a researcher to determine the cause of an outcome.
- 3) Margin of error increases as sample size increases.
- 4) A survey collected from a random sample of students in a school can be used to represent the opinions of the school population.

- 195 Given i is the imaginary unit, simplify $(5xi^3 - 4i)^2$ as a polynomial in standard form.

- 196 Which function has the characteristic as $x \rightarrow -\infty, f(x) \rightarrow -\infty$?



1)



2)

3) $f(x) = 5(4)^{-x}$

4) $f(x) = -\log_5(-x)$

- 197 Given that $\left(\frac{\frac{17}{8}}{\frac{y}{y^{\frac{5}{4}}}}\right)^{-4} = y^n$, where $y > 0$, determine the value of n .

- 198 Algebraically determine the values of x that satisfy the system of equations below:

$$y = x^2 + 8x - 5$$

$$y = 8x - 4$$

- 199 Describe the transformation applied to the graph of $p(x) = 2^x$ that forms the new function $q(x) = 2^{x-3} + 4$.

- 200 Solve algebraically for n : $\frac{2}{n^2} + \frac{3}{n} = \frac{4}{n^2}$

Algebra II Regents Exam Questions at Random Worksheet # 29

201 Given the polynomial identity $x^6 + y^6 = (x^2 + y^2)(x^4 - x^2y^2 + y^4)$, which equation must also be true for all values of x and y ?

- 1) $x^6 + y^6 = x^2(x^4 - x^2y^2 + y^4) + y^2(x^4 - x^2y^2 + y^4)$
- 2) $x^6 + y^6 = (x^2 + y^2)(x^2 - y^2)(x^2 - y^2)$
- 3) $(x^3 + y^3)^2 = (x^2 + y^2)(x^4 - x^2y^2 + y^4)$
- 4) $(x^6 + y^6) - (x^2 + y^2) = x^4 - x^2y^2 + y^4$

202 A rush-hour commuter train has arrived on time 64 of its first 80 days. As arrivals continue, which equation can be used to find x , the number of consecutive days that the train must arrive on schedule to raise its on-time performance rate to 90%?

- 1) $\frac{64}{80+x} = \frac{90}{100}$
- 2) $\frac{64+x}{80+x} = \frac{90}{100}$
- 3) $\frac{64+x}{80} = \frac{90}{100}$
- 4) $\frac{x}{80+x} = \frac{90}{100}$

203 Given $f(x) = 3^{x-1} + 2$, as $x \rightarrow -\infty$

- 1) $f(x) \rightarrow -1$
- 2) $f(x) \rightarrow 0$
- 3) $f(x) \rightarrow 2$
- 4) $f(x) \rightarrow -\infty$

204 Which function is even?

- 1) $f(x) = x^3 + 2$
- 2) $f(x) = x^2 + 1$
- 3) $f(x) = |x + 2|$
- 4) $f(x) = \sin(2x)$

205 The expression $\frac{x^4 - 5x^2 + 4x + 14}{x + 2}$ is equivalent to

- 1) $x^3 - 2x^2 - x + 6 + \frac{2}{x+2}$
- 2) $x^3 - 5x + 4 - \frac{14}{x+2}$
- 3) $x^3 + 2x^2 - x + 2 + \frac{18}{x+2}$
- 4) $x^3 + 2x^2 - 9x + 22 - \frac{30}{x+2}$

206 Factor completely over the set of integers:

$$-2x^4 + x^3 + 18x^2 - 9x$$

207 A Foucault pendulum can be used to demonstrate that the Earth rotates. The time, t , in seconds, that it takes for one swing or period of the pendulum

can be modeled by the equation $t = 2\pi\sqrt{\frac{L}{g}}$ where

L is the length of the pendulum in meters and g is a constant of 9.81 m/s^2 . The first Foucault pendulum was constructed in 1851 and has a pendulum length of 67 m. Determine, to the *nearest tenth of a second*, the time it takes this pendulum to complete one swing. Another Foucault pendulum at the United Nations building takes 9.6 seconds to complete one swing. Determine, to the *nearest tenth of a meter*, the length of this pendulum.

208 The amount of a substance, $A(t)$, in grams, remaining after t days is modeled by

$$A(t) = 50(0.5)^{\frac{t}{3}}. \text{ Which statement is false?}$$

- 1) In 20 days, there is no substance remaining.
- 2) After two half-lives, there is 25% of the substance remaining.
- 3) The amount of the substance remaining can also be modeled by

$$A(t) = 50(2)^{\frac{-t}{3}}.$$

- 4) After one week, there is less than 10g of the substance remaining.

209 Solve the system of equations algebraically.

$$x^2 + y^2 = 25$$


$$y + 5 = 2x$$

Algebra II Regents Exam Questions at Random Worksheet # 30

210 The expression $\frac{x^2 + 12}{x^2 + 3}$ can be rewritten as

- 1) $\frac{10}{x^2 + 3}$
- 2) $1 + \frac{9}{x^2 + 3}$
- 3) $x + 9$
- 4) 4

211 The parabola $y = -\frac{1}{20}(x - 3)^2 + 6$ has its focus at (3,1). Determine and state the equation of the directrix. (The use of the grid below is optional.)



212 The equation below can be used to model the height of a tide in feet, $H(t)$, on a beach at t hours.


$$H(t) = 4.8 \sin\left(\frac{\pi}{6}(t + 3)\right) + 5.1$$

Using this function, the amplitude of the tide is


- 1) $\frac{\pi}{6}$
- 2) 4.8
- 3) 3
- 4) 5.1

213 Solve the equation $\sqrt{49 - 10x} + 5 = 2x$ algebraically.

214 Consider the function $y = h(x)$, defined by the graph below.



Which equation could be used to represent the graph shown below?




- 1) $y = h(x) - 2$
- 2) $y = h(x - 2)$
- 3) $y = -h(x)$
- 4) $y = h(-x)$

215 Given $x \neq -3$, which expression is equivalent to $\frac{2x^3 + 3x^2 - 4x + 5}{x + 3}$?


- 1) $2x^3 + 9x^2 + 23x + 74$
- 2) $2x^2 - 3x + 5 - \frac{10}{x + 3}$
- 3) $2x^3 - 3x^2 + 5x - 10$
- 4) $2x^2 + 9x + 23 + \frac{74}{x + 3}$

- 216 An app design company believes that the proportion of high school students who have purchased apps on their smartphones in the past 3 months is 0.85. A simulation of 500 samples of 150 students was run based on this proportion and the results are shown below.



Suppose a sample of 150 students from your high school showed that 88% of students had purchased apps on their smartphones in the past 3 months. Based on the simulation, would the results from your high school give the app design company reason to believe their assumption is *incorrect*? Explain.

- 217 In the quadratic formula, $b^2 - 4ac$ is called the discriminant. The function $f(x)$ has a discriminant value of 8, and $g(x)$ has a discriminant value of -16 . The quadratic graphs, $h(x)$ and $j(x)$, are shown below.



Which quadratic functions have imaginary roots?

- 1) $g(x)$ and $h(x)$
- 2) $g(x)$ and $j(x)$
- 3) $f(x)$ and $h(x)$
- 4) $f(x)$ and $j(x)$

- 218 A parabola that has a vertex at $(2, 1)$ and a focus of $(2, -3)$ has an equation of

- 1) $y = \frac{1}{16}(x - 2)^2 + 1$
- 2) $y = -\frac{1}{16}(x + 2)^2 - 1$
- 3) $y = -\frac{1}{16}(x - 2)^2 + 1$
- 4) $y = -\frac{1}{16}(x - 2)^2 - 3$

- 219 A fruit fly population can be modeled by the equation $P = 10(1.27)^t$, where P represents the number of fruit flies after t days. What is the average rate of change of the population, rounded to the *nearest hundredth*, over the interval $[0, 10.5]$? Include appropriate units in your answer.

- 220 The table below shows the results of gender and music preference. Based on these data, determine if the events "the person is female" and "the person prefers classic rock" are independent of each other. Justify your answer.

	Rap	Techno	Classic Rock	Classical
Male	39	17	42	12
Female	17	37	36	15

- 221 In watching auditions for lead singer in a band, Liem became curious as to whether there is an association between how animated the lead singer is and the amount of applause from the audience. He decided to watch each singer and rate the singer on a scale of 1 to 5, where 1 is the least animated and 5 is the most animated. He did this for all 5 nights of auditions and found that the more animated singers did receive louder applause. The study Liem conducted would be best described as

- 1) experimental
- 2) observational
- 3) a sample survey
- 4) a random assignment

- 222 Which function has the greatest y-intercept?

- 1) $f(x) = 4\sin(2x)$
- 2) $g(x) = 3x^4 + 2x^3 + 7$
- 3) $h(x) = 5e^{2x} + 3$
- 4) $j(x) = 6\log_2(3x + 4)$


- 223 Solve algebraically for x to the nearest thousandth:

$$2e^{0.49x} = 15$$


- 224 When observed by researchers under a microscope, a smartphone screen contained approximately 11,000 bacteria per square inch. Bacteria, under normal conditions, double in population every 20 minutes.

- a) Assuming an initial value of 11,000 bacteria, write a function, $p(t)$, that can be used to model the population of bacteria, p , on a smartphone screen, where t represents the time in minutes after it is first observed under a microscope.
- b) Using $p(t)$ from part a, determine algebraically, to the nearest hundredth of a minute, the amount of time it would take for a smartphone screen that was not touched or cleaned to have a population of 1,000,000 bacteria per square inch.

- 225 Joette is playing a carnival game. To win a prize, one has to correctly guess which of five equally sized regions a spinner will land on, as shown in the diagram below.




She complains that the game is unfair because her favorite number, 2, has only been spun once in ten times she played the game. State the proportion of 2's that were spun. State the theoretical probability of spinning a 2. The simulation output below shows the results of simulating ten spins of a fair spinner, repeated 100 times.



Does the output indicate that the carnival game was unfair? Explain your answer.

- 226 In order to decrease the percentage of its residents who drive to work, a large city launches a campaign to encourage people to use public transportation instead. Before starting the campaign, the city's Department of Transportation uses census data to estimate that 65% of its residents drive to work. The Department of Transportation conducts a simulation, shown below, run 400 times based on this estimate. Each dot represents the proportion of 200 randomly selected residents who drive to work.



Use the simulation results to construct a plausible interval containing the middle 95% of the data. Round your answer to the *nearest hundredth*. One year after launching the campaign, the Department of Transportation conducts a survey of 200 randomly selected city residents and finds that 122 of them drive to work. Should the department conclude that the city's campaign was effective? Use statistical evidence from the simulation to explain your answer.

- 227 The population of Austin, Texas from 1850 to 2010 is summarized in the table below.

Year	1850	1870	1890	1910	1930	1950	1970	1990	2010
Population	629	4428	14,575	29,860	53,120	132,459	251,808	494,290	790,390

Over which period of time was the average rate of change in population the greatest?

- 1) 1850 to 1910
 - 2) 1990 to 2010
 - 3) 1950 to 1970
 - 4) 1890 to 1970
- 228 A recursive formula for the sequence 64, 48, 36, ... is
- 1) $a_n = 64(0.75)^{n-1}$
 - 2) $a_1 = 64$
 - $a_n = a_{n-1} - 16$
 - 3) $a_n = 64 + (n-1)(-16)$
 - 4) $a_1 = 64$
 - $a_n = 0.75a_{n-1}$
- 229 Algebraically solve the system:
- $$(x-2)^2 + (y-3)^2 = 20$$
- $$y = -2x + 7$$
- 230 The business office of a local college wishes to determine the methods of payment that will be used by students when buying books at the beginning of a semester. Explain how the office can gather an appropriate sample that minimizes bias.
- 231 Biologists are studying a new bacterium. They create a culture with 100 of the bacteria and anticipate that the number of bacteria will double every 30 hours. Write an equation for the number of bacteria, B , in terms of the number of hours, t , since the experiment began.

- 232 Objects cool at different rates based on the formula below.

$$T = (T_0 - T_R)e^{-rt} + T_R$$

T_0 : initial temperature

T_R : room temperature

r : rate of cooling of the object

t : time in minutes that the object

cools to a temperature, T

Mark makes T-shirts using a hot press to transfer designs to the shirts. He removes a shirt from a press that heats the shirt to 400°F . The rate of cooling for the shirt is 0.0735 and the room temperature is 75°F . Using this information, write an equation for the temperature of the shirt, T , after t minutes. Use the equation to find the temperature of the shirt, to the *nearest degree*, after five minutes. At the same time, Mark's friend Jeanine removes a hoodie from a press that heats the hoodie to 450°F . After eight minutes, the hoodie measured 270°F . The room temperature is still 75°F . Determine the rate of cooling of the hoodie, to the *nearest ten thousandth*. The T-shirt and hoodie were removed at the same time. Determine when the temperature will be the same, to the *nearest minute*.

- 233 Given $x \neq -3$, the expression $\frac{2x^3 + 7x^2 - 3x - 25}{x + 3}$ is equivalent to


- 1) $2x^2 + x - 6 - \frac{7}{x + 3}$
- 2) $2x^2 + 13x - 36 + \frac{83}{x + 3}$
- 3) $2x^2 + x - 13$
- 4) $x^2 + 4x - 15 + \frac{20}{x + 3}$

- 234 For all real values of x , if $f(x) = (x - 3)^2$ and $g(x) = (x + 3)^2$, what is $f(x) - g(x)$?
- 1) -18
 - 2) 0
 - 3) $-12x$
 - 4) $2x^2 - 12x - 18$

- 235 The expression $\frac{x^2 + 6}{x^2 + 4}$ is equivalent to

- 1) $\frac{6}{4}$
- 2) $1 + \frac{10}{x^2 + 4}$
- 3) $1 - \frac{2}{x^2 + 4}$
- 4) $1 + \frac{2}{x^2 + 4}$

- 236 Graph the following function on the axes below.
 $f(x) = \log_3(2 - x)$



State the domain of f . State the equation of the asymptote.

- 237 What is the solution set of the equation

$$\frac{4}{k^2 - 8k + 12} = \frac{k}{k - 2} + \frac{1}{k - 6}?$$


- 1) $\{-1, 6\}$
- 2) $\{1, -6\}$
- 3) $\{-1\}$
- 4) $\{1\}$

- 238 Factor the expression $x^3 - 2x^2 - 9x + 18$ completely.

- 239 Given $x \neq -2$, the expression $\frac{2x^2 + 5x + 8}{x + 2}$ is equivalent to

- 1) $2x^2 + \frac{9}{x+2}$
- 2) $2x + \frac{7}{x+2}$
- 3) $2x + 1 + \frac{6}{x+2}$
- 4) $2x + 9 - \frac{10}{x+2}$

- 240 Graph $y = 2 \cos\left(\frac{1}{2}x\right) + 5$ on the interval $[0, 2\pi]$, using the axes below.



- 241 A parabola has a directrix of $y = 3$ and a vertex at $(2, 1)$. Which ordered pair is the focus of the parabola?
- 1) $(2, -1)$
 - 2) $(2, 0)$
 - 3) $(2, 2)$
 - 4) $(2, 5)$

- 242 Solve for x algebraically:

$$\frac{1}{x-6} + \frac{x}{x-2} = \frac{4}{x^2 - 8x + 12}$$

- 243 A tree farm initially has 150 trees. Each year, 20% of the trees are cut down and 80 seedlings are planted. Which recursive formula models the number of trees, a_n , after n years?

- 1) $a_1 = 150$
 $a_n = a_{n-1}(0.2) + 80$
- 2) $a_1 = 150$
 $a_n = a_{n-1}(0.8) + 80$
- 3) $a_n = 150(0.2)^n + 80$
- 4) $a_n = 150(0.8)^n + 80$

- 244 What is the value of $\tan \theta$ when $\sin \theta = \frac{2}{5}$ and θ is in quadrant II?

- 1) $\frac{-\sqrt{21}}{5}$
- 2) $\frac{-\sqrt{21}}{2}$
- 3) $\frac{-2}{\sqrt{21}}$
- 4) $\frac{2}{\sqrt{21}}$

- 245 What is the solution for the system of equations below?

$$x + y + z = 2$$


$$x - 2y - z = -4$$

$$x - 9y + z = -18$$

- 1) $(-2, 2, 2)$
- 2) $(-2, -2, 6)$
- 3) $(0, 2, 0)$
- 4) $(0, 2, 4)$

- 246 Given $f(x) = 3x^3 - 4x^2 + 2x - 1$ and $g(x) = x - 4$, state the quotient and remainder of $\frac{f(x)}{g(x)}$, in the form $q(x) + \frac{r(x)}{g(x)}$. Is $x = 4$ a root of $f(x)$? Explain your answer.

247 Consider the graph of g and the table representing t below.



Over the interval $[2, 4]$, which statement regarding the average rate of change for g and t is true?


- 1) g has a greater average rate of change.
- 2) The average rates of change are equal.
- 3) The average rate of change for g is twice the average rate of change for t .
- 4) The average rate of change for g is half the average rate of change for t .

248 Consider the end behavior description below.


- as $x \rightarrow -\infty, f(x) \rightarrow \infty$
- as $x \rightarrow \infty, f(x) \rightarrow -\infty$

Which function satisfies the given conditions?

- 1) $f(x) = x^4 + 2x^2 + 1$



- 2)
- 3) $f(x) = -x^3 + 2x - 6$



- 4)


249 According to the USGS, an agency within the Department of Interior of the United States, the frog population in the U.S. is decreasing at the rate of 3.79% per year. A student created a model, $P = 12,150(0.962)^t$, to estimate the population in a pond after t years. The student then created a model that would predict the population after d decades. This model is best represented by

- 1) $P = 12,150(0.461)^d$
- 2) $P = 12,150(0.679)^d$
- 3) $P = 12,150(0.996)^d$
- 4) $P = 12,150(0.998)^d$

250 The expression $(x^2 + 3)^2 - 2(x^2 + 3) - 24$ is equivalent to

- 1) $(x^2 + 9)(x^2 - 1)$
- 2) $(x^2 - 3)(x^2 + 7)$
- 3) $x^4 - 2x^2 - 21$
- 4) $x^4 + 4x^2 - 9$


- 251 Betty conducted a survey of her class to see if they like pizza. She gathered 200 responses and 65% of the voters said they did like pizza. Betty then ran a simulation of 400 more surveys, each with 200 responses, assuming that 65% of the voters would like pizza. The output of the simulation is shown below.



Considering the middle 95% of the data, what is the margin of error for the simulation?

- | | |
|---------|---------|
| 1) 0.01 | 3) 0.05 |
| 2) 0.02 | 4) 0.07 |

- 252 Patricia creates a cubic polynomial function, $p(x)$, with a leading coefficient of 1. The zeros of the function are 2, 3, and -6 . Write an equation for $p(x)$. Sketch $y = p(x)$ on the set of axes below.



- 253 The polynomial function $g(x) = x^3 + ax^2 - 5x + 6$ has a factor of $(x - 3)$. Determine the value of a .
- 254 The population of bacteria, $P(t)$, in hundreds, after t hours can be modeled by the function $P(t) = 37e^{0.0532t}$. Determine whether the population is increasing or decreasing over time. Explain your reasoning.


- 255 Which expression is equivalent to $\frac{2x^3 + 2x - 7}{2x + 4}$?

- 1) $x^2 - 2x + 5 - \frac{27}{2x+4}$
- 2) $x^2 - 1 - \frac{3}{2x+4}$
- 3) $x^2 + 2x + 5 + \frac{13}{2x+4}$
- 4) $x^2 + 2x - 3 + \frac{5}{2x+4}$


- 256 Does the equation $x^2 - 4x + 13 = 0$ have imaginary solutions? Justify your answer.

- 257 The growth of a \$500 investment can be modeled by the function $P(t) = 500(1.03)^t$, where t represents time in years. In terms of the monthly rate of growth, the value of the investment can be best approximated by
- 1) $P(t) = 500(1.00247)^{12t}$
 - 2) $P(t) = 500(1.00247)^t$
 - 3) $P(t) = 500(1.03)^{12t}$
 - 4) $P(t) = 500(1.03)^{\frac{t}{12}}$

- 258 Which function has a maximum y-value of 4 and a midline of $y = 1$?



- 1)
2) $g(x) = -3\cos(x) + 1$



- 3)
4) $j(x) = 4\sin(x) + 1$

- 259 Given the inverse function $f^{-1}(x) = \frac{2}{3}x + \frac{1}{6}$, which function represents $f(x)$?

- 1) $f(x) = -\frac{2}{3}x + \frac{1}{6}$
2) $f(x) = -\frac{3}{2}x + \frac{1}{4}$
3) $f(x) = \frac{3}{2}x - \frac{1}{4}$
4) $f(x) = \frac{3}{2}x - \frac{1}{6}$

- 260 The expression $\left(a^3\sqrt{2b^2}\right)\left(\sqrt[3]{4a^2b}\right)$ is equivalent to

- 1) $2ab^3\sqrt{a^2}$
2) $2ab$
3) $2ab^3\sqrt[3]{2a^2}$
4) $2a^2b^3\sqrt[3]{2b}$

- 261 Which expression is an equivalent form of $a^5\sqrt[5]{a^4}$?

- 1) a
2) $a^{\frac{9}{5}}$
3) $a^{\frac{2}{4}}$
4) $a^{\frac{1}{5}}$

- 262 On a certain tropical island, there are currently 500 palm trees and 200 flamingos. Suppose the palm tree population is decreasing at an annual rate of 3% per year and the flamingo population is growing at a continuous rate of 2% per year. Write two functions, $P(x)$ and $F(x)$, that represent the number of palm trees and flamingos on this island, respectively, x years from now. State the solution to the equation $P(x) = F(x)$, rounded to the *nearest year*. Interpret the meaning of this value within the given context.

- 263 Which equation has roots of $3 + i$ and $3 - i$?

- 1) $x^2 - 6x + 10 = 0$
2) $x^2 + 6x - 10 = 0$
3) $x^2 - 10x + 6 = 0$
4) $x^2 + 10x - 6 = 0$

- 264 Consider a cubic polynomial with the characteristics below.

- exactly one real root
- as $x \rightarrow \infty, f(x) \rightarrow -\infty$

Given $a > 0$ and $b > 0$, which equation represents a cubic polynomial with these characteristics?

- 1) $f(x) = (x - a)(x^2 + b)$
2) $f(x) = (a - x)(x^2 + b)$
3) $f(x) = (a - x^2)(x^2 + b)$
4) $f(x) = (x - a)(b - x^2)$


Algebra II Regents Exam Questions at Random Worksheet # 39

- 265 The table below gives air pressures in kPa at selected altitudes above sea level measured in kilometers.

x	Altitude (km)	0	1	2	3	4	5
y	Air Pressure (kPa)	101	90	79	70	62	54

Write an exponential regression equation that models these data rounding all values to the *nearest thousandth*. Use this equation to algebraically determine the altitude, to the *nearest hundredth* of a kilometer, when the air pressure is 29 kPa.

- 266 Two classes of students were entered into an experiment to see whether using an interactive whiteboard leads to better grades. It was observed that the mean grade of students in the class with the interactive whiteboard was 0.6 points higher than the class without it. To determine if the observed difference is statistically significant, the classes were rerandomized 5000 times to study these random differences in the mean grades. The output of the simulation is summarized in the histogram below.



Determine an interval containing the middle 95% of the simulation results. Round your answer to the *nearest hundredth*. Does the interval indicate that the difference between the classes' grades is significant? Explain.

- 267 According to a study, 45% of Americans have type O blood. If a random number generator produces three-digit values from 000 to 999, which values would represent those having type O blood?
- 1) between 000 and 045, inclusive
 - 2) between 000 and 444, inclusive
 - 3) between 000 and 449, inclusive
 - 4) between 000 and 450, inclusive

- 268 A recursive formula for the sequence 40, 30, 22.5, ... is

- 1) $g_n = 40\left(\frac{3}{4}\right)^n$
- 2) $g_1 = 40$
 $g_n = g_{n-1} - 10$
- 3) $g_n = 40\left(\frac{3}{4}\right)^{n-1}$
- 4) $g_1 = 40$
 $g_n = \frac{3}{4}g_{n-1}$

- 269 The scores on a collegiate mathematics readiness assessment are approximately normally distributed with a mean of 680 and a standard deviation of 120. Determine the percentage of scores between 690 and 900, to the *nearest percent*.

- 270 To the *nearest tenth*, the solution to the equation $4300e^{0.07x} - 123 = 5000$ is

- 1) 1.1
- 2) 2.5
- 3) 6.3
- 4) 68.5

- 271 How many equations below are identities?

- $x^2 + y^2 = (x^2 - y^2) + (2xy)^2$
- $x^3 + y^3 = (x - y) + (x^2 - xy + y^2)$
- $x^4 + y^4 = (x - y)(x - y)(x^2 + y^2)$

- 1) 1
- 2) 2
- 3) 3
- 4) 0

- 272 The relative frequency table shows the proportion of a population who have a given eye color and the proportion of the same population who wear glasses.

	Wear Glasses	Don't Wear Glasses
Blue Eyes	0.14	0.26
Brown Eyes	0.11	0.24
Green Eyes	0.10	0.15

Given the data, are the events of having blue eyes and wearing glasses independent? Justify your answer.

- 273 If $f(t) = 50(.5)^{\frac{t}{5715}}$ represents a mass, in grams, of carbon-14 remaining after t years, which statement(s) must be true?

I. The mass of the carbon-14 is decreasing by half each year.

II. The mass of the original sample is 50 g.

- 1) I, only
 2) II, only
 3) I and II
 4) neither I nor II
- 274 Which expression is *not* a solution to the equation $2^t = \sqrt{10}$?
- 1) $\frac{1}{2} \log_2 10$
 2) $\log_2 \sqrt{10}$
 3) $\log_4 10$
 4) $\log_{10} 4$
- 275 Given $a > 0$, solve the equation $a^{x+1} = \sqrt[3]{a^2}$ for x algebraically.

- 276 If $\cos A = \frac{\sqrt{5}}{3}$ and $\tan A < 0$, what is the value of $\sin A$?
- 1) $\frac{2}{3}$
 2) $-\frac{\sqrt{5}}{3}$
 3) $-\frac{2}{3}$
 4) $\frac{3}{\sqrt{5}}$

- 277 Mia has a student loan that is in deferment, meaning that she does not need to make payments right now. The balance of her loan account during her deferment can be represented by the function $f(x) = 35,000(1.0325)^x$, where x is the number of years since the deferment began. If the bank decides to calculate her balance showing a monthly growth rate, an approximately equivalent function would be

- 1) $f(x) = 35,000(1.0027)^{12x}$
 2) $f(x) = 35,000(1.0027)^{\frac{x}{12}}$
 3) $f(x) = 35,000(1.0325)^{12x}$
 4) $f(x) = 35,000(1.0325)^{\frac{x}{12}}$

- 278 Which expression is *not* equivalent to $36x^6 - 25y^4$?

- 1) $6^2(x^3)^2 - 5^2(y^2)^2$
 2) $(6x^3 - 5y^2)(6x^3 + 5y^2)$
 3) $(6x^6 - 5y^4)(6x^6 + 5y^4)$
 4) $(3 \bullet 2x^3 - 5y^2)(3 \bullet 2x^3 + 5y^2)$

- 279 Consider the system of equations below?

$$x + 2y - z = 1$$


$$-x - 3y + 2z = 0$$

$$2x - 4y + z = 10$$

What is the solution to the given system of equations?


- 1) (1, 1, 2)
 2) (3, -1, 0)
 3) (5, -1, 2)
 4) (3, 5, 8)

- 280 Visible light can be represented by sinusoidal waves. Three visible light waves are shown in the graph below. The midline of each wave is labeled ℓ .



Based on the graph, which light wave has the longest period? Justify your answer.

- 281 A cyclist pedals a bike at a rate of 60 revolutions per minute. The height, h , of a pedal at time t , in seconds, is plotted below.



The graph can be modeled by the function $h(t) = 5 \sin(kt)$, where k is equal to

- 1) 1
- 2) 2π
- 3) 60
- 4) $\frac{\pi}{30}$

- 282 Algebraically solve the following system of equations.

$$(x-2)^2 + (y-3)^2 = 16$$

$$x + y - 1 = 0$$

- 283 Emmeline is working on one side of a polynomial identity proof used to form Pythagorean triples. Her work is shown below:

$$(5x)^2 + (5x^2 - 5)^2$$

Step 1: $25x^2 + (5x^2 - 5)^2$

Step 2: $25x^2 + 25x^2 + 25$

Step 3: $50x^2 + 25$

Step 4: $75x^2$


What statement is true regarding Emmeline's work?

- 1) Emmeline's work is entirely correct.
- 2) There is a mistake in step 2, only.
- 3) There are mistakes in step 2 and step 4.
- 4) There is a mistake in step 4, only.

- 284 If $(6 - ki)^2 = 27 - 36i$, the value of k is

- 1) -36
- 2) -3
- 3) 3
- 4) 6


- 285 The J& B candy company claims that 45% of the candies it produces are blue, 30% are brown, and 25% are yellow. Each bag holds 65 candies. A simulation was run 200 times, each of sample size 65, based on the premise that 45% of the candies are blue. The results of the simulation are shown below.



Bonnie purchased a bag of J& B's candy and counted 24 blue candies. What inference can be made regarding a bag of J& B's with only 24 blue candies?

- 1) The company is not meeting their production standard.
 - 2) Bonnie's bag was a rarity and the company should not be concerned.
 - 3) The company should change their claim to 37% blue candies are produced.
 - 4) Bonnie's bag is within the middle 95% of the simulated data supporting the company's claim.
- 286 A local university has a current enrollment of 12,000 students. The enrollment is increasing continuously at a rate of 2.5% each year. Which logarithm is equal to the number of years it will take for the population to increase to 15,000 students?
- 1) $\frac{\ln 1.25}{0.25}$
 - 2) $\frac{\ln 3000}{0.025}$
 - 3) $\frac{\ln 1.25}{2.5}$
 - 4) $\frac{\ln 1.25}{0.025}$
- 287 Monthly mortgage payments can be found using the formula below, where M is the monthly payment, P is the amount borrowed, r is the annual interest rate, and n is the total number of monthly payments.
- $$M = \frac{P\left(\frac{r}{12}\right)\left(1 + \frac{r}{12}\right)^n}{\left(1 + \frac{r}{12}\right)^n - 1}$$
- If Adam takes out a 15-year mortgage, borrowing \$240,000 at an annual interest rate of 4.5%, his monthly payment will be
- 1) \$1379.09
 - 2) \$1604.80
 - 3) \$1835.98
 - 4) \$9011.94
- 288 Consider the parabola given by $y = \frac{1}{4}x^2 + x + 8$ with vertex $(-2, 7)$ and focus $(-2, 8)$. Use this information to explain how to determine the equation of the directrix.
- 289 Which investigation technique is most often used to determine if a single variable has an impact on a given population?
- 1) observational study
 - 2) random survey
 - 3) controlled experiment
 - 4) formal interview
- 290 The heights of the 3300 students at Oceanview High School are approximately normally distributed with a mean of 65.5 inches and a standard deviation of 2.9 inches. The number of students at Oceanview who are between 64 and 68 inches tall is closest to
- 1) 1660
 - 2) 1070
 - 3) 2244
 - 4) 1640
- 291 Solve algebraically for all values of x :
- $$\sqrt{4x + 1} = 11 - x$$


- 292 A technology company is comparing two plans for speeding up its technical support time. Plan A can be modeled by the function $A(x) = 15.7(0.98)^x$ and plan B can be modeled by the function $B(x) = 11(0.99)^x$ where x is the number of customer service representatives employed by the company and $A(x)$ and $B(x)$ represent the average wait time, in minutes, of each customer. Graph $A(x)$ and $B(x)$ in the interval $0 \leq x \leq 100$ on the set of axes below.



To the *nearest integer*, solve the equation $A(x) = B(x)$. Determine, to the *nearest minute*, $B(100) - A(100)$. Explain what this value represents in the given context.

- 293 What are the zeros of $s(x) = x^4 - 9x^2 + 3x^3 - 27x - 10x^2 + 90$?
- 1) $\{-3, -2, 5\}$
 - 2) $\{-2, 3, 5\}$
 - 3) $\{-3, -2, 3, 5\}$
 - 4) $\{-5, -3, 2, 3\}$
- 294 Expressed in simplest $a + bi$ form, $(7 - 3i) + (x - 2i)^2 - (4i + 2x^2)$ is
- 1) $(3 - x^2) - (4x + 7)i$
 - 2) $(3 + 3x^2) - (4x + 7)i$
 - 3) $(3 - x^2) - 7i$
 - 4) $(3 + 3x^2) - 7i$

- 295 Which sketch could represent the function $m(x) = -\log_{100}(x - 2)$?



- 296 A function is defined as $a_n = a_{n-1} + \log_{n+1}(n - 1)$, where $a_1 = 8$. What is the value of a_3 ?
- 1) 8
 - 2) 8.5
 - 3) 9.2
 - 4) 10

- 297 Consider the system below.


$$x + y + z = 9$$

$$x - y - z = -1$$

$$x - y + z = 21$$

Which value is *not* in the solution, (x, y, z) , of the system?

- 1) -8
 - 2) -6
 - 3) 11
 - 4) 4
- 298 In the diagram of a unit circle below, point A, $\left(-\frac{\sqrt{3}}{2}, \frac{1}{2}\right)$, represents the point where the terminal side of θ intersects the unit circle.




What is $m\angle\theta$?

- 1) 30°
 - 2) 120°
 - 3) 135°
 - 4) 150°
- 299 Susan won \$2,000 and invested it into an account with an annual interest rate of 3.2%. If her investment were compounded monthly, which expression best represents the value of her investment after t years?
- 1) $2000(1.003)^{12t}$
 - 2) $2000(1.032)^{\frac{t}{12}}$
 - 3) $2064^{\frac{t}{12}}$
 - 4) $\frac{2000(1.032)^t}{12}$

300 The solution to the equation $5e^{x+2} = 7$ is

- 1) $-2 + \ln\left(\frac{7}{5}\right)$
- 2) $\left(\frac{\ln 7}{\ln 5}\right) - 2$
- 3) $\frac{-3}{5}$
- 4) $-2 + \ln(2)$

301 On the coordinate plane below, sketch *at least one cycle* of a cosine function with a midline at $y = -2$, an amplitude of 3, and a period of $\frac{\pi}{2}$.



302 Sarah is fighting a sinus infection. Her doctor prescribed a nasal spray and an antibiotic to fight the infection. The active ingredients, in milligrams, remaining in the bloodstream from the nasal spray, $n(t)$, and the antibiotic, $a(t)$, are modeled in the functions below, where t is the time in hours since the medications were taken.

$$n(t) = \frac{t+1}{t+5} + \frac{18}{t^2 + 8t + 15}$$

$$a(t) = \frac{9}{t+3}$$

Determine which drug is made with a greater initial amount of active ingredient. Justify your answer. Sarah's doctor told her to take both drugs at the same time. Determine algebraically the number of hours after taking the medications when both medications will have the same amount of active ingredient remaining in her bloodstream.

303 Given $p(\theta) = 3\sin\left(\frac{1}{2}\theta\right)$ on the interval

$-\pi < \theta < \pi$, the function p

- 1) decreases, then increases
- 2) increases, then decreases
- 3) decreases throughout the interval
- 4) increases throughout the interval

304 A retailer advertises that items will be discounted by 10% every Monday until they are sold. In how many weeks will an item costing \$50 first be sold for under half price?

- 1) 7
- 2) 6
- 3) 5
- 4) 4

305 Given $\cos A = \frac{3}{\sqrt{10}}$ and $\cot A = -3$, determine the value of $\sin A$ in radical form.

306 For which approximate value(s) of x will $\log(x+5) = |x-1| - 3$?

- 1) 5, 1
- 2) -2.41, 0.41
- 3) -2.41, 5
- 4) 5, only

307 For all positive values of x , which expression is

equivalent to $x^{\frac{3}{4}}$?

- 1) $\sqrt[4]{x^3}$
- 2) $\sqrt[3]{x^4}$
- 3) $(x^3)^4$
- 4) $3(x^4)$

308 The expression $(x+a)^2 + 5(x+a) + 4$ is equivalent to

- 1) $(a+1)(a+4)$
- 2) $(x+1)(x+4)$
- 3) $(x+a+1)(x+a+4)$
- 4) $x^2 + a^2 + 5x + 5a + 4$

Algebra II Regents Exam Questions at Random Worksheet # 46

309 Which equation represents a parabola with a focus of $(4, -3)$ and directrix of $y = 1$?

- 1) $(x - 1)^2 = 4(y + 3)$
- 2) $(x - 1)^2 = -8(y - 3)$
- 3) $(x + 4)^2 = 4(y - 3)$
- 4) $(x - 4)^2 = -8(y + 1)$

310 Which expression is equivalent to $(x + 2)^2 - 5(x + 2) + 6$?

- 1) $x(x - 1)$
- 2) $(x - 3)(x - 2)$
- 3) $(x - 4)(x + 3)$
- 4) $(x - 6)(x + 1)$

311 If $f(x) = 12x - 4$, then the inverse function $f^{-1}(x)$ is

- 1) $f^{-1}(x) = \frac{x + 1}{3}$
- 2) $f^{-1}(x) = \frac{x}{3} + 1$
- 3) $f^{-1}(x) = \frac{x + 4}{12}$
- 4) $f^{-1}(x) = \frac{x}{12} + 4$

312 Natalia's teacher has given her the following information about angle θ .

$$\bullet \pi < \theta < 2\pi$$

$$\bullet \cos \theta = \frac{\sqrt{3}}{4}$$

Explain how Natalia can determine if the value of $\tan \theta$ is positive or negative.

313 In a survey of people who recently bought a laptop, 45% said they were looking for a large screen, 31% said they were looking for a fast processor, and 58% said they wanted a large screen or a fast processor. If a survey respondent is selected at random, what is the probability that the respondent wanted both a large screen and a fast processor?

- 1) 76%
- 2) 14%
- 3) 77%
- 4) 18%

314 Factor the expression $2x^3 - 3x^2 - 18x + 27$ completely.

315 Over the set of integers, completely factor $x^4 - 5x^2 + 4$.

316 For the polynomial $p(x)$, if $p(3) = 0$, it can be concluded that

- 1) $x + 3$ is a factor of $p(x)$
- 2) $x - 3$ is a factor of $p(x)$
- 3) when $p(x)$ is divided by 3, the remainder is zero
- 4) when $p(x)$ is divided by -3 , the remainder is zero


317 For all values of x for which the expression is defined, write the expression below in simplest form.

$$\frac{2x^3 + x^2 - 18x - 9}{3x - x^2}$$

318 On the set of axes below, graph $y = f(x)$ and $y = g(x)$ for the given functions.

$$f(x) = x^3 - 3x^2$$

$$g(x) = 2x - 5$$



State the number of solutions to the equation $f(x) = g(x)$.


319 Consider the data in the table below.

	Right Handed	Left Handed
Male	87	13
Female	89	11

What is the probability that a randomly selected person is male given the person is left handed?

- 1) $\frac{13}{200}$ 3) $\frac{13}{50}$
 2) $\frac{13}{100}$ 4) $\frac{13}{24}$


320 The graph of the function $f(x)$ is shown below.



In which interval is $f(x)$ always positive?

- 1) $(-2, 4)$
 2) $(0, 10)$
 3) $(-12, -5)$
 4) $(-10, 0)$
- 321 A group of high school students wanted to collect information on how many times per week students exercised. If they want the *least* biased results they should survey every fifth student at the school who is
- 1) entering the gym
 2) in the junior class
 3) entering the library
 4) entering the building
- 322 The Hot and Tasty Coffee chain conducts a survey of its customers at its location at the Staten Island ferry terminal. After the survey is completed, the statistical consultant states that 70% of customers who took the survey said the most important factor in choosing where to get their coffee is how fast they are served. Based on this result, Hot and Tasty Coffee can infer that
- 1) most of its customers in New York State care most about being served quickly
 2) coffee drinkers care less about taste and more about being served quickly
 3) most of its customers at the Staten Island ferry terminal care most about being served quickly
 4) most of its customers at transportation terminals and stations care most about being served quickly
- 323 The function $N(x) = 90(0.86)^x + 69$ can be used to predict the temperature of a cup of hot chocolate in degrees Fahrenheit after x minutes. What is the approximate average rate of change of the temperature of the hot chocolate, in degrees per minute, over the interval $[0, 6]$?
- 1) -8.93
 2) -0.11
 3) 0.11
 4) 8.93
- 324 Which expression is a factor of $x^4 - x^3 - 11x^2 + 5x + 30$?
- 1) $x + 2$
 2) $x - 2$
 3) $x + 5$
 4) $x - 5$

- 325 Sketch
- $p(x) = -\log_2(x+3) + 2$
- on the axes below.

Describe the end behavior of $p(x)$ as $x \rightarrow -3$.Describe the end behavior of $p(x)$ as $x \rightarrow \infty$

- 326 A researcher wants to determine if room-darkening shades cause people to sleep longer. Which method of data collection is most appropriate?

- 1) census
- 2) survey
- 3) observation study
- 4) controlled experiment

- 327 The Manford family started savings accounts for their twins, Abby and Brett, on the day they were born. They invested \$8000 in an account for each child. Abby's account pays 4.2% annual interest compounded quarterly. Brett's account pays 3.9% annual interest compounded continuously. Write a function, $A(t)$, for Abby's account and a function, $B(t)$, for Brett's account that calculates the value of each account after t years. Determine who will have more money in their account when the twins turn 18 years old, and find the difference in the amounts in the accounts to the *nearest cent*. Algebraically determine, to the *nearest tenth of a year*, how long it takes for Brett's account to triple in value.

- 328 Algebraically determine the zeros of the function below.

$$r(x) = 3x^3 + 12x^2 - 3x - 12$$

- 329 If
- $f(x) = (x^2 + 3x + 2)(x^2 - 4x + 3)$
- and
- $g(x) = x^2 - 9$
- , then how many real solutions are there to the equation
- $f(x) = g(x)$
- ?

- 1) 1
- 2) 2
- 3) 6
- 4) 4

- 330 As
- θ
- increases from
- $-\frac{\pi}{2}$
- to 0 radians, the value of
- $\cos \theta$
- will

- 1) decrease from 1 to 0
- 2) decrease from 0 to -1
- 3) increase from -1 to 0
- 4) increase from 0 to 1

- 331 Luminescence is the emission of light that is not caused by heat. A luminescent substance decays according to the function below.

$$I = I_0 e^{3\left(-\frac{t}{0.6}\right)}$$

This function can be best approximated by

- 1) $I = I_0 e^{\left(-\frac{t}{0.18}\right)}$
- 2) $I = I_0 e^{5t}$
- 3) $I = I_0 (0.0067)^t$
- 4) $I = I_0 (0.0497)^{0.6t}$

- 332 Solve the following system of equations algebraically for
- x
- ,
- y
- , and
- z
- .

$$2x + 4y - 3z = 12$$


$$3x - 2y + 2z = -9$$

$$-x + y - 3z = 0$$

- 333 If
- $f(x) = \frac{1}{2}x + 2$
- , then the inverse function is

- 1) $f^{-1}(x) = -\frac{1}{2}x - 2$
- 2) $f^{-1}(x) = \frac{1}{2}x - 1$
- 3) $f^{-1}(x) = 2x - 4$
- 4) $f^{-1}(x) = 2x + 2$

334 Which equation best represents the graph below?



- 1) $h(x) = \log(x + a) + c$
- 2) $h(x) = \log(x - a) + c$
- 3) $h(x) = \log(x + a) - c$
- 4) $h(x) = \log(x - a) - c$

335 What is the total number of points of intersection of the graphs of the equations $y = e^x$ and $xy = 20$?

- 1) 1
- 2) 2
- 3) 3
- 4) 0

336 The solution set of $\frac{x+3}{x-5} + \frac{6}{x+2} = \frac{6+10x}{(x-5)(x+2)}$ is

- 1) $\{-6\}$
- 2) $\{5\}$
- 3) $\{-6, 5\}$
- 4) $\{-5, 6\}$

337 The element Americium has a half-life of 25 minutes. Given an initial amount, A_0 , which expression could be used to determine the amount of Americium remaining after t minutes?

- 1) $A_0 \left(\frac{1}{2} \right)^{\frac{t}{25}}$
- 2) $A_0 (25)^{\frac{t}{2}}$
- 3) $25 \left(\frac{1}{2} \right)^t$
- 4) $A_0 \left(\frac{1}{2} \right)^{25t}$

338 Which value, to the *nearest tenth*, is the *smallest* solution of $f(x) = g(x)$ if $f(x) = 3 \sin \left(\frac{1}{2}x \right) - 1$ and

$$g(x) = x^3 - 2x + 1?$$


- 1) -3.6
- 2) -2.1
- 3) -1.8
- 4) 1.4

339 For n and $p > 0$, is the expression

$$\left(p^2 n^{\frac{1}{2}} \right)^8 \sqrt{p^5 n^4} \text{ equivalent to } p^{18} n^6 \sqrt{p}?$$

Justify your answer.

340 State officials claim 82% of a community want to repeal the 30 mph speed limit on an expressway. A community organization devises a simulation based on the claim that 82% of the community supports the repeal. Each dot on the graph below represents the proportion of community members who support the repeal. The graph shows 200 simulated surveys, each of sample size 60.



Based on the simulation, determine an interval containing the middle 95% of plausible proportions. Round your answer to the *nearest thousandth*. The community organization conducted its own sample survey of 60 people and found 70% supported the repeal. Based on the results of the simulation, explain why the organization should question the State officials' claim.


- 341 A popular celebrity tracks the number of people, in thousands, who have followed her on social media since January 1, 2015. A summary of the data she recorded is shown in the table below:

Number of Months Since January 2015	2	11	16	20	27	35	47	50	52
Number of Social Media Followers (thousands)	3.1	7.5	29.7	49.7	200.3	680.3	5200.3	8109.3	12,107.1


The celebrity uses an exponential regression equation to model the data. According to the model, about how many followers did she have on June 1, 2018?

- 1) 13,000,000
 - 2) 5,420,000
 - 3) 1,850,000
 - 4) 790,000
- 342 Write a recursive formula for the sequence 189, 63, 21, 7, ...
- 343 What is the solution set of $x = \sqrt{3x + 40}$?
- 1) $\{-5, 8\}$
 - 2) $\{8\}$
 - 3) $\{-4, 10\}$
 - 4) $\{ \}$
- 344 The roots of the equation $x^2 - 4x = -13$ are
- 1) $2 \pm 3i$
 - 2) $2 \pm 6i$
 - 3) $2 \pm \sqrt{17}$
 - 4) $2 \pm \sqrt{13}$
- 345 Which situation best describes conditional probability?
- 1) finding the probability of an event occurring two or more times
 - 2) finding the probability of an event occurring only once
 - 3) finding the probability of two independent events occurring at the same time
 - 4) finding the probability of an event occurring given another event had already occurred
- 346 The expression $3i(ai - 6i^2)$ is equivalent to
- 1) $3a + 18i$
 - 2) $3a - 18i$
 - 3) $-3a + 18i$
 - 4) $-3a - 18i$
- 347 John and Margaret deposit \$500 into a savings account for their son on his first birthday. They continue to make a deposit of \$500 on the child's birthday, with the last deposit being made on the child's 21st birthday. If the account pays 4% annual interest, which equation represents the amount of money in the account after the last deposit is made?
- 1) $S_{21} = 500(1.04)^{21}$
 - 2) $S_{21} = \frac{500(1 - 1.04^{21})}{1 - 1.04}$
 - 3) $S_{21} = 500(1.04)^{20} + 500$
 - 4) $S_{21} = \frac{500(1 - 0.04^{21})}{1 - 1.04}$
- 348 The initial push of a child on a swing causes the swing to travel a total of 6 feet. Each successive swing travels 80% of the distance of the previous swing. Determine the total distance, to the *nearest hundredth of a foot*, a child travels in the first five swings.
- 349 Which function represents exponential decay?
- 1) $p(x) = \left(\frac{1}{4}\right)^{-x}$
 - 2) $q(x) = 1.8^{-x}$
 - 3) $r(x) = 2.3^{2x}$
 - 4) $s(x) = 4^{\frac{x}{2}}$


- 350 Which graph shows a quadratic function with two imaginary zeros?




1)



2)



3)



- 351 The inverse of $f(x) = -6x + \frac{1}{2}$ is

- 1) $f^{-1}(x) = 6x - \frac{1}{2}$
- 2) $f^{-1}(x) = \frac{1}{-6x + \frac{1}{2}}$
- 3) $f^{-1}(x) = -\frac{1}{6}x + \frac{1}{12}$
- 4) $f^{-1}(x) = -\frac{1}{6}x + 2$

- 352 The amount of a substance, $A(t)$, that remains after t days can be given by the equation

$A(t) = A_0(0.5)^{\frac{t}{0.0803}}$, where A_0 represents the initial amount of the substance. An equivalent form of this equation is

- 1) $A(t) = A_0(0.000178)^t$
- 2) $A(t) = A_0(0.945861)^t$
- 3) $A(t) = A_0(0.04015)^t$
- 4) $A(t) = A_0(1.08361)^t$

- 353 Solve the equation $3x^2 + 5x + 8 = 0$. Write your solution in $a + bi$ form.

- 354 The average monthly temperature, $T(m)$, in degrees Fahrenheit, over a 12 month period, can be

modeled by $T(m) = -23 \cos\left(\frac{\pi}{6}m\right) + 56$, where m is

in months. What is the range of temperatures, in degrees Fahrenheit, of this function?

- 1) $[-23, 23]$
- 2) $[33, 79]$
- 3) $[-23, 56]$
- 4) $[-79, 33]$

- 355 Given $f(x) = -\frac{2}{5}x + 4$, which statement is true of the inverse function $f^{-1}(x)$?

- 1) $f^{-1}(x)$ is a line with slope $\frac{5}{2}$.
- 2) $f^{-1}(x)$ is a line with slope $\frac{2}{5}$.
- 3) $f^{-1}(x)$ passes through the point $(6, -5)$.
- 4) $f^{-1}(x)$ has a y-intercept at $(0, -4)$.

- 356 Which statement below about the graph of $f(x) = -\log(x + 4) + 2$ is true?

- 1) $f(x)$ has a y-intercept at $(0, 2)$.
- 2) $-f(x)$ has a y-intercept at $(0, 2)$.
- 3) As $x \rightarrow \infty, f(x) \rightarrow \infty$.
- 4) $x \rightarrow -4, f(x) \rightarrow \infty$.

- 357 A company fired several employees in order to save money. The amount of money the company saved per year over five years following the loss of employees is shown in the table below.

Year	Amount Saved (in dollars)
1	59,000
2	64,900
3	71,390
4	78,529
5	86,381.9

Which expression determines the total amount of money saved by the company over 5 years?

- 1) $\frac{59,000 - 59,000(1.1)^5}{1 - 1.1}$ 3) $\sum_{n=1}^5 59,000(1.1)^n$
- 2) $\frac{59,000 - 59,000(0.1)^5}{1 - 0.1}$ 4) $\sum_{n=1}^5 59,000(0.1)^{n-1}$
- 358 If a solution of $2(2x - 1) = 5x^2$ is expressed in simplest $a + bi$ form, the value of b is

- 1) $\frac{\sqrt{6}}{5}i$
- 2) $\frac{\sqrt{6}}{5}$
- 3) $\frac{1}{5}i$
- 4) $\frac{1}{5}$


- 359 The population, in millions of people, of the United States can be represented by the recursive formula below, where a_0 represents the population in 1910 and n represents the number of years since 1910.

$$a_0 = 92.2$$

$$a_n = 1.015a_{n-1}$$

Identify the percentage of the annual rate of growth from the equation $a_n = 1.015a_{n-1}$. Write an exponential function, P , where $P(t)$ represents the United States population in millions of people, and t is the number of years since 1910. According to this model, determine algebraically the number of years it takes for the population of the United States to be approximately 300 million people. Round your answer to the *nearest year*.

- 360 Graph $c(x) = -9(3)^{x-4} + 2$ on the axes below.



Describe the end behavior of $c(x)$ as x approaches positive infinity. Describe the end behavior of $c(x)$ as x approaches negative infinity.

- 361 Given $P(A) = \frac{1}{3}$ and $P(B) = \frac{5}{12}$, where A and B are independent events, determine $P(A \cap B)$.

Algebra II Regents Exam Questions at Random Worksheet # 53

362 What is the solution of $2(3^{x+4}) = 56$?

- 1) $x = \log_3(28) - 4$
- 2) $x = -1$
- 3) $x = \log(25) - 4$
- 4) $x = \frac{\log(56)}{\log(6)} - 4$

363 Given the geometric series $300 + 360 + 432 + 518.4 + \dots$, write a geometric series formula, S_n , for the sum of the first n terms. Use the formula to find the sum of the first 10 terms, to the *nearest tenth*.

364 Given $x > 0$, the expression $\frac{x^{\frac{1}{5}}}{x^{\frac{1}{2}}}$ can be rewritten as

- 1) $\sqrt[3]{x}$
- 2) $-\sqrt[10]{x^3}$
- 3) $\frac{1}{\sqrt[10]{x^3}}$
- 4) $\sqrt[3]{x^{10}}$

365 In a group of 40 people, 20 have brown hair, 22 have blue eyes, and 15 have both brown hair and blue eyes. How many people have neither brown hair nor blue eyes?

- 1) 0
- 2) 13
- 3) 27
- 4) 32

366 The expression $\sqrt[4]{81x^8y^6}$ is equivalent to

- 1) $3x^2y^{\frac{3}{2}}$
- 2) $3x^4y^2$
- 3) $9x^2y^{\frac{3}{2}}$
- 4) $9x^4y^2$

367 Completely factor the following expression:
 $x^2 + 3xy + 3x^3 + y$


368 Stone Manufacturing has developed a cost model, $C(x) = 0.18x^3 + 0.02x^2 + 4x + 180$, where x is the number of sprockets sold, in thousands. The sales price can be modeled by $S(x) = 95.4 - 6x$ and the company's revenue by $R(x) = x \cdot S(x)$. The company's profits, $R(x) - C(x)$, could be modeled by

- 1) $0.18x^3 + 6.02x^2 + 91.4x + 180$
- 2) $0.18x^3 - 5.98x^2 - 91.4x + 180$
- 3) $-0.18x^3 - 6.02x^2 + 91.4x - 180$
- 4) $0.18x^3 + 5.98x^2 + 99.4x + 180$

369 The solution set for the equation $\sqrt{3(x+6)} = x$ is

- 1) $\{6, -3\}$
- 2) $\{-6, 3\}$
- 3) $\{6\}$
- 4) $\{-3\}$

370 Graph $y = x^3 - 4x^2 + 2x + 7$ on the set of axes below.




371 For which values of x , rounded to the *nearest hundredth*, will $|x^2 - 9| - 3 = \log_3 x$?

- 1) 2.29 and 3.63
- 2) 2.37 and 3.54
- 3) 2.84 and 3.17
- 4) 2.92 and 3.06

Algebra II Regents at Random Worksheets

- 372 Some smart-phone applications contain "in-app" purchases, which allow users to purchase special content within the application. A random sample of 140 users found that 35 percent made in-app purchases. A simulation was conducted with 200 samples of 140 users assuming 35 percent of the samples make in-app purchases. The approximately normal results are shown below.



Considering the middle 95% of the data, determine the margin of error, to the *nearest hundredth*, for the simulated results. In the given context, explain what this value represents.

- 373 A number, minus twenty times its reciprocal, equals eight. The number is
- 1) 10 or -2
 - 2) 10 or 2
 - 3) -10 or -2
 - 4) -10 or 2
- 374 A study of black bears in the Adirondacks reveals that their population can be represented by the function $P(t) = 3500(1.025)^t$, where t is the number of years since the study began. Which function is correctly rewritten to reveal the monthly growth rate of the black bear population?
- 1) $P(t) = 3500(1.00206)^{12t}$
 - 2) $P(t) = 3500(1.00206)^{\frac{t}{12}}$
 - 3) $P(t) = 3500(1.34489)^{12t}$
 - 4) $P(t) = 3500(1.34489)^{\frac{t}{12}}$
- 375 The Wells family is looking to purchase a home in a suburb of Rochester with a 30-year mortgage that has an annual interest rate of 3.6%. The house the family wants to purchase is \$152,500 and they will make a \$15,250 down payment and borrow the remainder. Use the formula below to determine their monthly payment, to the *nearest dollar*.
- $$M = \frac{P\left(\frac{r}{12}\right)\left(1 + \frac{r}{12}\right)^n}{\left(1 + \frac{r}{12}\right)^n - 1}$$
- M = monthly payment
 P = amount borrowed
 r = annual interest rate
 n = total number of monthly payments

Algebra II Regents Exam Questions at Random Worksheet # 55

376 Consider the system of equations below:

$$x + y - z = 6$$

$$2x - 3y + 2z = -19$$

$$-x + 4y - z = 17$$

Which number is *not* the value of any variable in the solution of the system?

- 1) -1
- 2) 2
- 3) 3
- 4) -4

377 Algebraically solve for x : $\frac{7}{2x} - \frac{2}{x+1} = \frac{1}{4}$

378 A random sample of 100 people that would best estimate the proportion of all registered voters in a district who support improvements to the high school football field should be drawn from registered voters in the district at a

- 1) football game
- 2) supermarket
- 3) school fund-raiser
- 4) high school band concert

379 A veterinary pharmaceutical company plans to test a new drug to treat a common intestinal infection among puppies. The puppies are randomly assigned to two equal groups. Half of the puppies will receive the drug, and the other half will receive a placebo. The veterinarians monitor the puppies. This is an example of which study method?

- 1) census
- 2) observational study
- 3) survey
- 4) controlled experiment

380 Solve the following system of equations algebraically. $x^2 + y^2 = 400$

$$y = x - 28$$

381 Rowan is training to run in a race. He runs 15 miles in the first week, and each week following, he runs 3% more than the week before. Using a geometric series formula, find the total number of miles Rowan runs over the first ten weeks of training, rounded to the *nearest thousandth*.

382 Which table best represents an exponential relationship?

x	y
1	8
2	4
3	2
4	1
5	$\frac{1}{2}$

1)

x	y
8	0
4	1
0	2
-4	3
-8	4

2)

x	y
0	0
1	1
2	4
3	9
4	16

3)

x	y
1	1
2	8
3	27
4	64
5	125

4)

383 If $f(x) = a^x$ where $a > 1$, then the inverse of the function is

- 1) $f^{-1}(x) = \log_x a$
- 2) $f^{-1}(x) = a \log x$
- 3) $f^{-1}(x) = \log_a x$
- 4) $f^{-1}(x) = x \log a$

Algebra II Regents Exam Questions at Random Worksheet # 56

- 384 A runner is using a nine-week training app to prepare for a "fun run." The table below represents the amount of the program completed, A , and the distance covered in a session, D , in miles.

A	$\frac{4}{9}$	$\frac{5}{9}$	$\frac{6}{9}$	$\frac{8}{9}$	1
D	2	2	2.25	3	3.25

Based on these data, write an exponential regression equation, rounded to the *nearest thousandth*, to model the distance the runner is able to complete in a session as she continues through the nine-week program.

- 385 A savings account, S , has an initial value of \$50. The account grows at a 2% interest rate compounded n times per year, t , according to the function below.

$$S(t) = 50 \left(1 + \frac{.02}{n} \right)^{nt}$$

Which statement about the account is correct?

- As the value of n increases, the amount of interest per year decreases.
- As the value of n increases, the value of the account approaches the function $S(t) = 50e^{0.02t}$.
- As the value of n decreases to one, the amount of interest per year increases.
- As the value of n decreases to one, the value of the account approaches the function $S(t) = 50(1 - 0.02)^t$.

- 386 Solve the following system of equations algebraically for all values of x , y , and z :

$$2x + 3y - 4z = -1$$

$$x - 2y + 5z = 3$$

$$-4x + y + z = 16$$

- 387 The function below models the average price of gas in a small town since January 1st.

$$G(t) = -0.0049t^4 + 0.0923t^3 - 0.56t^2 + 1.166t + 3.23,$$

where $0 \leq t \leq 10$.

If $G(t)$ is the average price of gas in dollars and t represents the number of months since January 1st, the absolute maximum $G(t)$ reaches over the given domain is about

- \$1.60
- \$3.92
- \$4.01
- \$7.73

- 388 Which description could represent the graph of

$$f(x) = 4x^2(x + a) - x - a, \text{ if } a \text{ is an integer?}$$

- As $x \rightarrow -\infty, f(x) \rightarrow \infty$, as $x \rightarrow \infty, f(x) \rightarrow \infty$, and the graph has 3 x -intercepts.
- As $x \rightarrow -\infty, f(x) \rightarrow -\infty$, as $x \rightarrow \infty, f(x) \rightarrow \infty$, and the graph has 3 x -intercepts.
- As $x \rightarrow -\infty, f(x) \rightarrow \infty$, as $x \rightarrow \infty, f(x) \rightarrow -\infty$, and the graph has 4 x -intercepts.
- As $x \rightarrow -\infty, f(x) \rightarrow -\infty$, as $x \rightarrow \infty, f(x) \rightarrow \infty$, and the graph has 4 x -intercepts.

- 389 Given $y > 0$, the expression $\sqrt{3x^2y} \bullet \sqrt[3]{27x^3y^2}$ is equivalent to

- $81x^5y^3$
- $3^{1.5}x^2y$
- $3^{\frac{5}{2}}x^2y^{\frac{5}{3}}$
- $3^{\frac{3}{2}}x^2y^{\frac{7}{6}}$

- 390 Savannah just got contact lenses. Her doctor said she can wear them 2 hours the first day, and can then increase the length of time by 30 minutes each day. If this pattern continues, which formula would *not* be appropriate to determine the length of time, in either minutes or hours, she could wear her contact lenses on the n th day?

$$1) \quad a_1 = 120$$

$$a_n = a_{n-1} + 30$$

$$2) \quad a_n = 90 + 30n$$

$$3) \quad a_1 = 2$$

$$a_n = a_{n-1} + 0.5$$

$$4) \quad a_n = 2.5 + 0.5n$$

Algebra II Regents Exam Questions at Random Worksheet # 57

- 391 The value(s) of x that satisfy

$$\sqrt{x^2 - 4x - 5} = 2x - 10 \text{ are}$$

- 1) $\{5\}$
- 2) $\{7\}$
- 3) $\{5, 7\}$
- 4) $\{3, 5, 7\}$

- 392 There are 400 students in the senior class at Oak Creek High School. All of these students took the SAT. The distribution of their SAT scores is approximately normal. The number of students who scored within 2 standard deviations of the mean is approximately

- 1) 75
- 2) 95
- 3) 300
- 4) 380

- 393 If $n = \sqrt{a^5}$ and $m = a$, where $a > 0$, an expression for $\frac{n}{m}$ could be

- 1) $a^{\frac{5}{2}}$
- 2) a^4
- 3) $\sqrt[3]{a^2}$
- 4) $\sqrt{a^3}$

- 394 Consider the probability statements regarding events A and B below.

$$P(A \text{ or } B) = 0.3;$$

$$P(A \text{ and } B) = 0.2; \text{ and}$$

$$P(A|B) = 0.8$$


What is $P(B)$?

- 1) 0.1
- 2) 0.25
- 3) 0.375
- 4) 0.667

- 395 The solution set for the equation $b = \sqrt{2b^2 - 64}$ is

- 1) $\{-8\}$
- 2) $\{8\}$
- 3) $\{\pm 8\}$
- 4) $\{ \}$

- 396 Data for the students enrolled in a local high school are shown in the Venn diagram below.



If a student from the high school is selected at random, what is the probability that the student is a sophomore given that the student is enrolled in Algebra II?

- 1) $\frac{85}{210}$
- 2) $\frac{85}{295}$
- 3) $\frac{85}{405}$
- 4) $\frac{85}{1600}$

- 397 What are the solution(s) to the system of equations shown below?

$$x^2 + y^2 = 5$$

$$y = 2x$$

- 1) $x = 1$ and $x = -1$
- 2) $x = 1$
- 3) $(1, 2)$ and $(-1, -2)$
- 4) $(1, 2)$, only

- 398 Suppose events A and B are independent and $P(A \text{ and } B)$ is 0.2. Which statement could be true?

- 1) $P(A) = 0.4$, $P(B) = 0.3$, $P(A \text{ or } B) = 0.5$
- 2) $P(A) = 0.8$, $P(B) = 0.25$
- 3) $P(A|B) = 0.2$, $P(B) = 0.2$
- 4) $P(A) = 0.15$, $P(B) = 0.05$

- 399 If $f(x) = x^2 + 9$ and $g(x) = x + 3$, which operation would not result in a polynomial expression?

- 1) $f(x) + g(x)$
- 2) $f(x) - g(x)$
- 3) $f(x) \bullet g(x)$
- 4) $f(x) \div g(x)$

400 The table below shows the number of hours of daylight on the first day of each month in Rochester, NY.

Month	Hours of Daylight
Jan.	9.4
Feb.	10.6
March	11.9
April	13.9
May	14.7
June	15.4
July	15.1
Aug.	13.9
Sept.	12.5
Oct.	11.1
Nov.	9.7
Dec.	9.0

Given the data, what is the average rate of change in hours of daylight per month from January 1st to April 1st? Interpret what this means in the context of the problem.

- 401 Perry invested in property that cost him \$1500. Five years later it was worth \$3000, and 10 years from his original purchase, it was worth \$6000. Assuming the growth rate remains the same, which type of function could he create to find the value of his investment 30 years from his original purchase?
- 1) exponential function
 - 2) linear function
 - 3) quadratic function
 - 4) trigonometric function

- 402 Solve algebraically for all values of x :
- $$\sqrt{6-2x} + x = 2(x+15) - 9$$

- 403 The expression $\frac{9x^2-2}{3x+1}$ is equivalent to


- 1) $3x - 1 - \frac{1}{3x+1}$
- 2) $3x - 1 + \frac{1}{3x+1}$
- 3) $3x + 1 - \frac{1}{3x+1}$
- 4) $3x + 1 + \frac{1}{3x+1}$

- 404 Determine, to the *nearest tenth of a year*, how long it would take an investment to double at a $3\frac{3}{4}\%$ interest rate, compounded continuously.

- 405 A student is chosen at random from the student body at a given high school. The probability that the student selects Math as the favorite subject is $\frac{1}{4}$. The probability that the student chosen is a junior is $\frac{116}{459}$. If the probability that the student selected is a junior or that the student chooses Math as the favorite subject is $\frac{47}{108}$, what is the exact probability that the student selected is a junior whose favorite subject is Math? Are the events "the student is a junior" and "the student's favorite subject is Math" independent of each other? Explain your answer.

- 406 If the function $g(x) = ab^x$ represents exponential growth, which statement about $g(x)$ is *false*?
- 1) $a > 0$ and $b > 1$
 - 2) The y -intercept is $(0, a)$.
 - 3) The asymptote is $y = 0$.
 - 4) The x -intercept is $(b, 0)$.


- 407 Griffin is riding his bike down the street in Churchville, N.Y. at a constant speed, when a nail gets caught in one of his tires. The height of the nail above the ground, in inches, can be represented by the trigonometric function $f(t) = -13\cos(0.8\pi t) + 13$, where t represents the time (in seconds) since the nail first became caught in the tire. Determine the period of $f(t)$. Interpret what the period represents in this context. On the grid below, graph *at least one* cycle of $f(t)$ that includes the y-intercept of the function.



Does the height of the nail ever reach 30 inches above the ground? Justify your answer.

- 408 For all values of x for which the expression is defined, $\frac{x^3 + 2x^2 - 9x - 18}{x^3 - x^2 - 6x}$, in simplest form, is equivalent to
- 1) 3
 - 2) $-\frac{17}{2}$
 - 3) $\frac{x+3}{x}$
 - 4) $\frac{x^2 - 9}{x(x-3)}$

- 409 The hours of daylight, y , in Utica in days, x , from January 1, 2013 can be modeled by the equation $y = 3.06\sin(0.017x - 1.40) + 12.23$. How many hours of daylight, to the *nearest tenth*, does this model predict for February 14, 2013?
- 1) 9.4
 - 2) 10.4
 - 3) 12.1
 - 4) 12.2
- 410 The half-life of iodine-131 is 8 days. The percent of the isotope left in the body d days after being introduced is $I = 100\left(\frac{1}{2}\right)^{\frac{d}{8}}$. When this equation is written in terms of the number e , the base of the natural logarithm, it is equivalent to $I = 100e^{kd}$. What is the approximate value of the constant, k ?
- 1) -0.087
 - 2) 0.087
 - 3) -11.542
 - 4) 11.542
- 411 Sketch the graphs of $r(x) = \frac{1}{x}$ and $a(x) = |x| - 3$ on the set of axes below. Determine, to the *nearest tenth*, the positive solution of $r(x) = a(x)$.



Algebra II Regents Exam Questions at Random Worksheet # 60


412 If $A = -3 + 5i$, $B = 4 - 2i$, and $C = 1 + 6i$, where i is the imaginary unit, then $A - BC$ equals

- 1) $5 - 17i$
- 2) $5 + 27i$
- 3) $-19 - 17i$
- 4) $-19 + 27i$

413 Which expression is equivalent to $(2x - i)^2 - (2x - i)(2x + 3i)$ where i is the imaginary unit and x is a real number?

- 1) $-4 - 8xi$
- 2) $-4 - 4xi$
- 3) 2
- 4) $8x - 4i$

414 Determine an equation for the parabola with focus $(4, -1)$ and directrix $y = -5$. (Use of the grid below is optional.)



415 Tides are a periodic rise and fall of ocean water. On a typical day at a seaport, to predict the time of the next high tide, the most important value to have would be the

- 1) time between consecutive low tides
- 2) time when the tide height is 20 feet
- 3) average depth of water over a 24-hour period
- 4) difference between the water heights at low and high tide

416 A manufacturing plant produces two different-sized containers of peanuts. One container weighs x ounces and the other weighs y pounds. If a gift set can hold one of each size container, which expression represents the number of gift sets needed to hold 124 ounces?

- 1) $\frac{124}{16x + y}$
- 2) $\frac{x + 16y}{124}$
- 3) $\frac{124}{x + 16y}$
- 4) $\frac{16x + y}{124}$

417 What is the solution set of the equation

$$\frac{2}{x} - \frac{3x}{x+3} = \frac{x}{x+3}?$$

- 1) $\{3\}$
- 2) $\left\{\frac{3}{2}\right\}$
- 3) $\{-2, 3\}$
- 4) $\left\{-1, \frac{3}{2}\right\}$

418 What is the solution set of the equation

$$\frac{2}{3x+1} = \frac{1}{x} - \frac{6x}{3x+1}?$$

- 1) $\left\{-\frac{1}{3}, \frac{1}{2}\right\}$
- 2) $\left\{-\frac{1}{3}\right\}$
- 3) $\left\{\frac{1}{2}\right\}$
- 4) $\left\{\frac{1}{3}, -2\right\}$

419 What is the inverse of the function $y = 4x + 5$?

- 1) $x = \frac{1}{4}y - \frac{5}{4}$
- 2) $y = \frac{1}{4}x - \frac{5}{4}$
- 3) $y = 4x - 5$
- 4) $y = \frac{1}{4x+5}$

Algebra II Regents Exam Questions at Random Worksheet # 61

- 420 Sodium iodide-131, used to treat certain medical conditions, has a half-life of 1.8 hours. The data table below shows the amount of sodium iodide-131, rounded to the nearest thousandth, as the dose fades over time.

Number of Half Lives	1	2	3	4	5
Amount of Sodium Iodide-131	139.000	69.500	34.750	17.375	8.688

What approximate amount of sodium iodide-131 will remain in the body after 18 hours?

- | | |
|----------|----------|
| 1) 0.001 | 3) 0.271 |
| 2) 0.136 | 4) 0.543 |

- 421 Determine for which polynomial(s) $(x + 2)$ is a factor. Explain your answer.

$$P(x) = x^4 - 3x^3 - 16x - 12$$

$$O(x) = x^3 - 3x^2 - 16x - 12$$

- 422 On July 21, 2016, the water level in Puget Sound, WA reached a high of 10.1 ft at 6 a.m. and a low of -2 ft at 12:30 p.m. Across the country in Long Island, NY, Shinnecock Bay's water level reached a high of 2.5 ft at 10:42 p.m. and a low of -0.1 ft at 5:31 a.m. The water levels of both locations are affected by the tides and can be modeled by sinusoidal functions. Determine the difference in amplitudes, in feet, for these two locations.

- 423 Explain what a rational exponent, such as $\frac{5}{2}$ means.

Use this explanation to evaluate $9^{\frac{5}{2}}$.

- 424 The height above ground for a person riding a
Ferris wheel after t seconds is modeled by

$h(t) = 150 \sin\left(\frac{\pi}{45}t + 67.5\right) + 160$ feet. How many

seconds does it take to go from the bottom of the wheel to the top of the wheel?

- 1) 10
- 2) 45
- 3) 90
- 4) 150

- 425 Which expression is equivalent to

$$\frac{2x^4 + 8x^3 - 25x^2 - 6x + 14}{x + 6}?$$

- 1) $2x^3 + 4x^2 + x - 12 + \frac{86}{x+6}$
- 2) $2x^3 - 4x^2 - x + 14$
- 3) $2x^3 - 4x^2 - x + \frac{14}{x+6}$
- 4) $2x^3 - 4x^2 - x$


- 426 The operator of the local mall wants to find out how many of the mall's employees make purchases in the food court when they are working. She hopes to use these data to increase the rent and attract new food vendors. In total, there are 1023 employees who work at the mall. The best method to obtain a random sample of the employees would be to survey
- 1) all 170 employees at each of the larger stores
 - 2) 50% of the 90 employees of the food court
 - 3) every employee
 - 4) every 30th employee entering each mall entrance for one week

- 427 Which statement is true about the graph of

$$f(x) = \left(\frac{1}{8}\right)^x ?$$

- 1) The graph is always increasing.
- 2) The graph is always decreasing.
- 3) The graph passes through $(1,0)$.
- 4) The graph has an asymptote, $x = 0$.


- 428 Website popularity ratings are often determined using models that incorporate the number of visits per week a website receives. One model for ranking websites is $P(x) = \log(x - 4)$, where x is the number of visits per week in thousands and $P(x)$ is the website's popularity rating. According to this model, if a website is visited 16,000 times in one week, what is its popularity rating, rounded to the *nearest tenth*? Graph $y = P(x)$ on the axes below.



An alternative rating model is represented by $R(x) = \frac{1}{2}x - 6$, where x is the number of visits per week in thousands. Graph $R(x)$ on the same set of axes. For what number of weekly visits will the two models provide the same rating?

- 429 If $ae^{bt} = c$, where a , b , and c are positive, then t equals
- 1) $\ln\left(\frac{c}{ab}\right)$
 - 2) $\ln\left(\frac{cb}{a}\right)$
 - 3) $\frac{\ln\left(\frac{c}{a}\right)}{b}$
 - 4) $\frac{\ln\left(\frac{c}{a}\right)}{\ln b}$
- 430 Determine the quotient and remainder when $(6a^3 + 11a^2 - 4a - 9)$ is divided by $(3a - 2)$. Express your answer in the form $q(a) + \frac{r(a)}{d(a)}$.
- 431 If $(a^3 + 27) = (a + 3)(a^2 + ma + 9)$, then m equals
- 1) -9
 - 2) -3
 - 3) 3
 - 4) 6
- 432 Explain why $81^{\frac{3}{4}}$ equals 27.

- 433 The results of simulating tossing a coin 10 times, recording the number of heads, and repeating this 50 times are shown in the graph below.




Based on the results of the simulation, which statement is *false*?

- 1) Five heads occurred most often, which is consistent with the theoretical probability of obtaining a heads.
 - 2) Eight heads is unusual, as it falls outside the middle 95% of the data.
 - 3) Obtaining three heads or fewer occurred 28% of the time.
 - 4) Seven heads is not unusual, as it falls within the middle 95% of the data.
- 434 The parabola described by the equation $y = \frac{1}{12}(x-2)^2 + 2$ has the directrix at $y = -1$. The focus of the parabola is
- 1) $(2, -1)$
 - 2) $(2, 2)$
 - 3) $(2, 3)$
 - 4) $(2, 5)$
- 435 A 7-year lease for office space states that the annual rent is \$85,000 for the first year and will increase by 6% each additional year of the lease. What will the total rent expense be for the entire 7-year lease?
- 1) \$42,809.63
 - 2) \$90,425.53
 - 3) \$595,000.00
 - 4) \$713,476.20

- 436 Researchers in a local area found that the population of rabbits with an initial population of 20 grew continuously at the rate of 5% per month. The fox population had an initial value of 30 and grew continuously at the rate of 3% per month. Find, to the *nearest tenth of a month*, how long it takes for these populations to be equal.

- 437 At Andrew Jackson High School, students are only allowed to enroll in AP U.S. History if they have already taken AP World History or AP European History. Out of 825 incoming seniors, 165 took AP World History, 66 took AP European History, and 33 took both. Given this information, determine the probability a randomly selected incoming senior is allowed to enroll in AP U.S. History.

- 438 Consider the function $p(x) = 3x^3 + x^2 - 5x$ and the graph of $y = m(x)$ below.



Which statement is true?

- 1) $p(x)$ has three real roots and $m(x)$ has two real roots.
 - 2) $p(x)$ has one real root and $m(x)$ has two real roots.
 - 3) $p(x)$ has two real roots and $m(x)$ has three real roots.
 - 4) $p(x)$ has three real roots and $m(x)$ has four real roots.
- 439 Over the set of integers, factor the expression $x^4 - 4x^2 - 12$.

Algebra II Regents Exam Questions at Random Worksheet # 64

- 440 Jake wants to buy a car and hopes to save at least \$5000 for a down payment. The table below summarizes the amount of money he plans to save each week.

Week	1	2	3	4	5
Money Saved, in Dollars	2	5	12.5	31.25	...

Based on this plan, which expression should he use to determine how much he has saved in n weeks?

- 1) $\frac{2 - 2(2.5^n)}{1 - 2.5}$ 3) $\frac{1 - 2.5^n}{1 - 2.5}$
 2) $\frac{2 - 2(2.5^{n-1})}{1 - 2.5}$ 4) $\frac{1 - 2.5^{n-1}}{1 - 2.5}$

- 441 After Roger's surgery, his doctor administered pain medication in the following amounts in milligrams over four days.

Day (n)	1	2	3	4
Dosage (m)	2000	1680	1411.2	1185.4

How can this sequence best be modeled recursively?

- 1) $m_1 = 2000$ 3) $m_1 = 2000$
 $m_n = m_{n-1} - 320$ $m_n = (0.84)m_{n-1}$
 2) $m_n = 2000(0.84)^{n-1}$ 4) $m_n = 2000(0.84)^{n+1}$

- 442 A population of 950 bacteria grows continuously at a rate of 4.75% per day. Write an exponential function, $N(t)$, that represents the bacterial population after t days and explain the reason for your choice of base. Determine the bacterial population after 36 hours, to the *nearest bacterium*.

- 443 If $f(x)$ is an even function, which function must also be even?

- 1) $f(x - 2)$
 2) $f(x) + 3$
 3) $f(x + 1)$
 4) $f(x + 1) + 3$

- 444 Which equation represents a parabola with a focus of $(-2, 5)$ and a directrix of $y = 9$?

- 1) $(y - 7)^2 = 8(x + 2)$
 2) $(y - 7)^2 = -8(x + 2)$
 3) $(x + 2)^2 = 8(y - 7)$
 4) $(x + 2)^2 = -8(y - 7)$

- 445 The average monthly high temperature in Buffalo, in degrees Fahrenheit, can be modeled by the function $B(t) = 25.29 \sin(0.4895t - 1.9752) + 55.2877$, where t is the month number (January = 1). State, to the *nearest tenth*, the average monthly rate of temperature change between August and November. Explain its meaning in the given context.

- 446 When the function $p(x)$ is divided by $x - 1$ the quotient is $x^2 + 7 + \frac{5}{x - 1}$. State $p(x)$ in standard form.

- 447 Evaluate $j(-1)$ given $j(x) = 2x^4 - x^3 - 35x^2 + 16x + 48$. Explain what your answer tells you about $x + 1$ as a factor. Algebraically find the remaining zeros of $j(x)$.

- 448 Given $a(x) = x^4 + 2x^3 + 4x - 10$ and $b(x) = x + 2$, determine $\frac{a(x)}{b(x)}$ in the form $q(x) + \frac{r(x)}{b(x)}$. Is $b(x)$ a factor of $a(x)$? Explain.

- 449 A formula for work problems involving two people is shown below.

$$\frac{1}{t_1} + \frac{1}{t_2} = \frac{1}{t_b}$$

t_1 = the time taken by the first person to complete the job

t_2 = the time taken by the second person to complete the job

t_b = the time it takes for them working together to complete the job

Fred and Barney are carpenters who build the same model desk. It takes Fred eight hours to build the desk while it only takes Barney six hours. Write an equation that can be used to find the time it would take both carpenters working together to build a desk. Determine, to the *nearest tenth of an hour*, how long it would take Fred and Barney working together to build a desk.

- 450 Algebraically solve for x : $\frac{-3}{x+3} + \frac{1}{2} = \frac{x}{6} - \frac{1}{2}$

- 451 How many solutions exist for

$$\frac{1}{1-x^2} = -|3x-2| + 5?$$

- 1) 1
- 2) 2
- 3) 3
- 4) 4

- 452 Carla wants to start a college fund for her daughter Lila. She puts \$63,000 into an account that grows at a rate of 2.55% per year, compounded monthly. Write a function, $C(t)$, that represents the amount of money in the account t years after the account is opened, given that no more money is deposited into or withdrawn from the account. Calculate algebraically the number of years it will take for the account to reach \$100,000, to the *nearest hundredth of a year*.

- 453 Which statement(s) are true for all real numbers?

I $(x-y)^2 = x^2 + y^2$

II $(x+y)^3 = x^3 + 3xy + y^3$

- 1) I, only
- 2) II, only
- 3) I and II
- 4) neither I nor II

- 454 The weights of bags of Graseck's Chocolate Candies are normally distributed with a mean of 4.3 ounces and a standard deviation of 0.05 ounces. What is the probability that a bag of these chocolate candies weighs less than 4.27 ounces?


- 1) 0.2257
- 2) 0.2743
- 3) 0.7257
- 4) 0.7757

- 455 What is the solution set for x in the equation below?

$$\sqrt{x+1} - 1 = x$$

- 1) {1}
- 2) {0}
- 3) {-1,0}
- 4) {0,1}

- 456 A scatterplot showing the weight, w , in grams, of each crystal after growing t hours is shown below.



The relationship between weight, w , and time, t , is best modeled by

- 1) $w = 4^t + 5$
- 2) $w = (1.4)^t + 2$
- 3) $w = 5(2.1)^t$
- 4) $w = 8(.75)^t$


- 457 The populations of two small towns at the beginning of 2018 and their annual population growth rate are shown in the table below.

Town	Population	Annual Population Growth Rate
Jonesville	1240	6% increase
Williamstown	890	11% increase

Assuming the trend continues, approximately how many years after the beginning of 2018 will it take for the populations to be equal?

- | | |
|-------|--------|
| 1) 7 | 3) 68 |
| 2) 20 | 4) 125 |

- 458 Graph $f(x) = \log_2(x + 6)$ on the set of axes below.



- 459 Which equation represents the equation of the parabola with focus $(-3, 3)$ and directrix $y = 7$?

- 1) $y = \frac{1}{8}(x+3)^2 - 5$
- 2) $y = \frac{1}{8}(x-3)^2 + 5$
- 3) $y = -\frac{1}{8}(x+3)^2 + 5$
- 4) $y = -\frac{1}{8}(x-3)^2 + 5$

- 460 Given: $f(x) = 2x^2 + x - 3$ and $g(x) = x - 1$
Express $f(x) \bullet g(x) - [f(x) + g(x)]$ as a polynomial
in standard form.

- 461 Evan graphed a cubic function,
 $f(x) = ax^3 + bx^2 + cx + d$, and determined the roots
of $f(x)$ to be ± 1 and 2. What is the value of b , if
 $a = 1$?

- 1) 1
- 2) 2
- 3) -1
- 4) -2

- 462 Which expression(s) are equivalent to $\frac{x^2 - 4x}{2x}$, where $x \neq 0$?

I. $\frac{x}{2} - 2$ II. $\frac{x-4}{2}$ III. $\frac{x-1}{2} - \frac{3}{2}$

- 1) II, only
- 2) I and II
- 3) II and III
- 4) I, II, and III

- 463 A fast-food restaurant analyzes data to better serve its customers. After its analysis, it discovers that the events D , that a customer uses the drive-thru, and F , that a customer orders French fries, are independent. The following data are given in a report:

$$P(F) = 0.8$$

$$P(F \cap D) = 0.456$$


Given this information, $P(F|D)$ is

- 1) 0.344
- 2) 0.3648
- 3) 0.57
- 4) 0.8

- 464 On a given school day, the probability that Nick oversleeps is 48% and the probability he has a pop quiz is 25%. Assuming these two events are independent, what is the probability that Nick oversleeps and has a pop quiz on the same day?

1) 73%
2) 36%
3) 23%
4) 12%

- 465 Consider $f(x) = 4x^2 + 6x - 3$, and $p(x)$ defined by the graph below.



The difference between the values of the maximum of p and minimum of f is

1) 0.25
2) 1.25
3) 3.25
4) 10.25


- 466 The average monthly temperature of a city can be modeled by a cosine graph. Melissa has been living in Phoenix, Arizona, where the average annual temperature is 75°F . She would like to move, and live in a location where the average annual temperature is 62°F . When examining the graphs of the average monthly temperatures for various locations, Melissa should focus on the

1) amplitude
2) horizontal shift
3) period
4) midline

- 467 The resting blood pressure of an adult patient can be modeled by the function P below, where $P(t)$ is the pressure in millimeters of mercury after time t in seconds.

$$P(t) = 24\cos(3\pi t) + 120$$

On the set of axes below, graph $y = P(t)$ over the domain $0 \leq t \leq 2$.




Determine the period of P . Explain what this value represents in the given context. Normal resting blood pressure for an adult is 120 over 80. This means that the blood pressure oscillates between a maximum of 120 and a minimum of 80. Adults with high blood pressure (above 140 over 90) and adults with low blood pressure (below 90 over 60) may be at risk for health disorders. Classify the given patient's blood pressure as low, normal, or high and explain your reasoning.

- 468 What is the solution when the equation $wx^2 + w = 0$ is solved for x , where w is a positive integer?

1) -1
2) 0
3) 6
4) $\pm i$

- 469 Mary bought a pack of candy. The manufacturer claims that 30% of the candies manufactured are red. In her pack, 14 of the 60 candies are red. She ran a simulation of 300 samples, assuming the manufacturer is correct. The results are shown below.



Based on the simulation, determine the middle 95% of plausible values that the proportion of red candies in a pack is within. Based on the simulation, is it unusual that Mary's pack had 14 red candies out of a total of 60? Explain.


- 470 A survey about television-viewing preferences was given to randomly selected freshmen and seniors at Fairport High School. The results are shown in the table below.

Favorite Type of Program			
	Sports	Reality Show	Comedy Series
Senior	83	110	67
Freshmen	119	103	54

A student response is selected at random from the results. State the *exact* probability the student response is from a freshman, given the student prefers to watch reality shows on television.

- 471 When a ball bounces, the heights of consecutive bounces form a geometric sequence. The height of the first bounce is 121 centimeters and the height of the third bounce is 64 centimeters. To the *nearest centimeter*, what is the height of the fifth bounce?
- 1) 25
 - 2) 34
 - 3) 36
 - 4) 42
- 472 Which function's graph has a period of 8 and reaches a maximum height of 1 if at least one full period is graphed?
- 1) $y = -4 \cos\left(\frac{\pi}{4}x\right) - 3$
 - 2) $y = -4 \cos\left(\frac{\pi}{4}x\right) + 5$
 - 3) $y = -4 \cos(8x) - 3$
 - 4) $y = -4 \cos(8x) + 5$

- 473 A group of students was trying to determine the proportion of candies in a bag that are blue. The company claims that 24% of candies in bags are blue. A simulation was run 100 times with a sample size of 50, based on the premise that 24% of the candies are blue. The approximately normal results of the simulation are shown in the dot plot below.




The simulation results in a mean of 0.254 and a standard deviation of 0.060. Based on this simulation, what is a plausible interval containing the middle 95% of the data?

- 1) (0.194, 0.314)
 - 2) (0.134, 0.374)
 - 3) (-0.448, 0.568)
 - 4) (0.254, 0.374)
- 474 The profit function, $p(x)$, for a company is the cost function, $c(x)$, subtracted from the revenue function, $r(x)$. The profit function for the Acme Corporation is $p(x) = -0.5x^2 + 250x - 300$ and the revenue function is $r(x) = -0.3x^2 + 150x$. The cost function for the Acme Corporation is
- 1) $c(x) = 0.2x^2 - 100x + 300$
 - 2) $c(x) = 0.2x^2 + 100x + 300$
 - 3) $c(x) = -0.2x^2 + 100x - 300$
 - 4) $c(x) = -0.8x^2 + 400x - 300$
- 475 The graph of $y = \log_2 x$ is translated to the right 1 unit and down 1 unit. The coordinates of the x -intercept of the translated graph are
- 1) (0, 0)
 - 2) (1, 0)
 - 3) (2, 0)
 - 4) (3, 0)
- 476 A person's lung capacity can be modeled by the function $C(t) = 250 \sin\left(\frac{2\pi}{5}t\right) + 2450$, where $C(t)$ represents the volume in mL present in the lungs after t seconds. State the maximum value of this function over one full cycle, and explain what this value represents.
- 477 The completely factored form of $n^4 - 9n^2 + 4n^3 - 36n - 12n^2 + 108$ is
- 1) $(n^2 - 9)(n + 6)(n - 2)$
 - 2) $(n + 3)(n - 3)(n + 6)(n - 2)$
 - 3) $(n - 3)(n - 3)(n + 6)(n - 2)$
 - 4) $(n + 3)(n - 3)(n - 6)(n + 2)$
- 478 Kelly-Ann has \$20,000 to invest. She puts half of the money into an account that grows at an annual rate of 0.9% compounded monthly. At the same time, she puts the other half of the money into an account that grows continuously at an annual rate of 0.8%. Which function represents the value of Kelly-Ann's investments after t years?
- 1) $f(t) = 10,000(1.9)^t + 10,000e^{0.8t}$
 - 2) $f(t) = 10,000(1.009)^t + 10,000e^{0.008t}$
 - 3) $f(t) = 10,000(1.075)^{12t} + 10,000e^{0.8t}$
 - 4) $f(t) = 10,000(1.00075)^{12t} + 10,000e^{0.008t}$
- 479 If $p(x) = 2x^3 - 3x + 5$, what is the remainder of $p(x) \div (x - 5)$?
- 1) -230
 - 2) 0
 - 3) 40
 - 4) 240

- 480 The temperature, in degrees Fahrenheit, in Times Square during a day in August can be predicted by the function $T(x) = 8\sin(0.3x - 3) + 74$, where x is the number of hours after midnight. According to this model, the predicted temperature, to the nearest degree Fahrenheit, at 7 P.M. is


1) 68
2) 74
3) 77
4) 81

- 481 Tony is evaluating his retirement savings. He currently has \$318,000 in his account, which earns an interest rate of 7% compounded annually. He wants to determine how much he will have in the account in the future, even if he makes no additional contributions to the account. Write a function, $A(t)$, to represent the amount of money that will be in his account in t years. Graph $A(t)$ where $0 \leq t \leq 20$ on the set of axes below.



Tony's goal is to save \$1,000,000. Determine algebraically, to the nearest year, how many years it will take for him to achieve his goal. Explain how your graph of $A(t)$ confirms your answer.

- 482 The depth of the water at a marker 20 feet from the shore in a bay is depicted in the graph below.



If the depth, d , is measured in feet and time, t , is measured in hours since midnight, what is an equation for the depth of the water at the marker?

- 1) $d = 5 \cos\left(\frac{\pi}{6}t\right) + 9$
2) $d = 9 \cos\left(\frac{\pi}{6}t\right) + 5$
3) $d = 9 \sin\left(\frac{\pi}{6}t\right) + 5$
4) $d = 5 \sin\left(\frac{\pi}{6}t\right) + 9$

- 483 A certain pain reliever is taken in 220 mg dosages and has a half-life of 12 hours. The function

$$A = 220\left(\frac{1}{2}\right)^{\frac{t}{12}}$$

can be used to model this situation,


where A is the amount of pain reliever in milligrams remaining in the body after t hours. According to this function, which statement is true?

- 1) Every hour, the amount of pain reliever remaining is cut in half.
2) In 12 hours, there is no pain reliever remaining in the body.
3) In 24 hours, there is no pain reliever remaining in the body.
4) In 12 hours, 110 mg of pain reliever is remaining.

484 To determine if the type of music played while taking a quiz has a relationship to results, 16 students were randomly assigned to either a room softly playing classical music or a room softly playing rap music. The results on the quiz were as follows:

Classical: 74, 83, 77, 77, 84, 82, 90, 89
Rap: 77, 80, 78, 74, 69, 72, 78, 69

John correctly rounded the difference of the means of his experimental groups as 7. How did John obtain this value and what does it represent in the given context? Justify your answer. To determine if there is any significance in this value, John rerandomized the 16 scores into two groups of 8, calculated the difference of the means, and simulated this process 250 times as shown below.




Does the simulation support the theory that there may be a significant difference in quiz scores? Explain.

485 Consider the function $h(x) = 2 \sin(3x) + 1$ and the function q represented in the table below.

x	$q(x)$
-2	-8
-1	0
0	0
1	-2
2	0

Determine which function has the *smaller* minimum value for the domain $[-2,2]$. Justify your answer.

- 486 Robin flips a coin 100 times. It lands heads up 43 times, and she wonders if the coin is unfair. She runs a computer simulation of 750 samples of 100 fair coin flips. The output of the proportion of heads is shown below.



Do the results of the simulation provide strong evidence that Robin's coin is unfair? Explain your answer.

- 487 When the expression $(x + 2)^2 + 4(x + 2) + 3$ is rewritten as the product of two binomials, the result is
- 1) $(x + 3)(x + 1)$
 - 2) $(x + 5)(x + 3)$
 - 3) $(x + 2)(x + 2)$
 - 4) $(x + 6)(x + 1)$

- 488 The height, $h(t)$ in cm, of a piston, is given by the equation $h(t) = 12 \cos\left(\frac{\pi}{3}t\right) + 8$, where t represents the number of seconds since the measurements began. Determine the average rate of change, in cm/sec, of the piston's height on the interval $1 \leq t \leq 2$. At what value(s) of t , to the *nearest tenth of a second*, does $h(t) = 0$ in the interval $1 \leq t \leq 5$? Justify your answer.

- 489 If $x - 1$ is a factor of $x^3 - kx^2 + 2x$, what is the value of k ?
- 1) 0
 - 2) 2
 - 3) 3
 - 4) -3


- 490 The solutions to $x + 3 - \frac{4}{x-1} = 5$ are

- 1) $\frac{3}{2} \pm \frac{\sqrt{17}}{2}$
- 2) $\frac{3}{2} \pm \frac{\sqrt{17}}{2}i$
- 3) $\frac{3}{2} \pm \frac{\sqrt{33}}{2}$
- 4) $\frac{3}{2} \pm \frac{\sqrt{33}}{2}i$

- 491 Camryn puts \$400 into a savings account that earns 6% annually. The amount in her account can be modeled by $C(t) = 400(1.06)^t$ where t is the time in years. Which expression best approximates the amount of money in her account using a weekly growth rate?

- 1) $400(1.001153846)^t$
- 2) $400(1.001121184)^t$
- 3) $400(1.001153846)^{52t}$
- 4) $400(1.001121184)^{52t}$


- 492 A sketch of $r(x)$ is shown below.




An equation for $r(x)$ could be

- 1) $r(x) = (x - a)(x + b)(x + c)$
- 2) $r(x) = (x + a)(x - b)(x - c)^2$
- 3) $r(x) = (x + a)(x - b)(x - c)$
- 4) $r(x) = (x - a)(x + b)(x + c)^2$


- 493 Which graph represents a polynomial function that contains $x^2 + 2x + 1$ as a factor?




1)



2)



3)



4)

- 494 Point $M\left(t, \frac{4}{7}\right)$ is located in the second quadrant on the unit circle. Determine the exact value of t .

- 495 For $x > 0$, which expression is equivalent to

$$\frac{\sqrt[3]{x^2} \cdot \sqrt{x^5}}{\sqrt[6]{x}}?$$

- 1) x
- 2) $x^{\frac{3}{2}}$
- 3) x^3
- 4) x^{10}

- 496 The world population was 2560 million people in 1950 and 3040 million in 1960 and can be modeled by the function $p(t) = 2560e^{0.017185t}$, where t is time in years after 1950 and $p(t)$ is the population in millions. Determine the average rate of change of $p(t)$ in millions of people per year, from $4 \leq t \leq 8$. Round your answer to the nearest hundredth.

- 497 For a given time, x , in seconds, an electric current, y , can be represented by $y = 2.5(1 - 2.7^{-.10x})$.

Which equation is *not* equivalent?

- 1) $y = 2.5 - 2.5(2.7^{-.10x})$
- 2) $y = 2.5 - 2.5\left((2.7^2)^{-.05x}\right)$
- 3) $y = 2.5 - 2.5\left(\frac{1}{2.7^{.10x}}\right)$
- 4) $y = 2.5 - 2.5(2.7^{-2})(2.7^{.05x})$

- 498 Erin and Christa were working on cubing binomials for math homework. Erin believed they could save time with a shortcut. She wrote down the rule below for Christa to follow.

$$(a + b)^3 = a^3 + b^3$$

Does Erin's shortcut always work? Justify your result algebraically.

- 499 Solve the given equation algebraically for all values of x . $3\sqrt{x} - 2x = -5$

Algebra II Regents Exam Questions at Random Worksheet # 74

500 Given $f(x) = \frac{1}{2}x + 8$, which equation represents the inverse, $g(x)$?

- 1) $g(x) = 2x - 8$
- 2) $g(x) = 2x - 16$
- 3) $g(x) = -\frac{1}{2}x + 8$
- 4) $g(x) = -\frac{1}{2}x - 16$

501 There are 440 students at Thomas Paine High School enrolled in U.S. History. On the April report card, the students' grades are approximately normally distributed with a mean of 79 and a standard deviation of 7. Students who earn a grade less than or equal to 64.9 must attend summer school. The number of students who must attend summer school for U.S. History is closest to

- 1) 3
- 2) 5
- 3) 10
- 4) 22


502 Which situation could be modeled using a geometric sequence?

- 1) A cell phone company charges \$30.00 per month for 2 gigabytes of data and \$12.50 for each additional gigabyte of data.
- 2) The temperature in your car is 79° . You lower the temperature of your air conditioning by 2° every 3 minutes in order to find a comfortable temperature.
- 3) David's parents have set a limit of 50 minutes per week that he may play online games during the school year. However, they will increase his time by 5% per week for the next ten weeks.
- 4) Sarah has \$100.00 in her piggy bank and saves an additional \$15.00 each week.

503 Given $c(m) = m^3 - 2m^2 + 4m - 8$, the solution of $c(m) = 0$ is

- 1) ± 2
- 2) 2, only
- 3) $2i, 2$
- 4) $\pm 2i, 2$


504 Graph $t(x) = 3\sin(2x) + 2$ over the domain $[0, 2\pi]$ on the set of axes below.



505 Which representation of a quadratic has imaginary roots?


x	y
-2.5	2
-2.0	0
-1.5	-1
-1.0	-1
-0.5	0
0.0	2

- 1)
- 2) $2(x + 3)^2 = 64$



- 3)
- 4) $2x^2 + 32 = 0$

- 506 The graph of $y = f(x)$ is shown below. The function has a leading coefficient of 1.



Write an equation for $f(x)$. The function g is formed by translating function f left 2 units. Write an equation for $g(x)$.

- 507 An angle, θ , is in standard position and its terminal side passes through the point $(2, -1)$. Find the *exact* value of $\sin \theta$.

- 508 The average depreciation rate of a new boat is approximately 8% per year. If a new boat is purchased at a price of \$75,000, which model is a recursive formula representing the value of the boat n years after it was purchased?

1) $a_n = 75,000(0.08)^n$

2) $a_0 = 75,000$

$a_n = (0.92)^n$

3) $a_n = 75,000(1.08)^n$

4) $a_0 = 75,000$

$a_n = 0.92(a_{n-1})$


- 509 Given $\tan \theta = \frac{7}{24}$, and θ terminates in Quadrant III, determine the value of $\cos \theta$.

- 510 Chuck's Trucking Company has decided to initiate an Employee of the Month program. To determine the recipient, they put the following sign on the back of each truck.



The driver who receives the highest number of positive comments will win the recognition. Explain *one* statistical bias in this data collection method.

- 511 The zeros of a quartic polynomial function h are $-1, \pm 2$, and 3. Sketch a graph of $y = h(x)$ on the grid below.



- 512 Where i is the imaginary unit, the expression $(x + 3i)^2 - (2x - 3i)^2$ is equivalent to


1) $-3x^2$

2) $-3x^2 - 18$


3) $-3x^2 + 18xi$

4) $-3x^2 - 6xi - 18$

513 On the grid below, graph the function $y = \log_2(x - 3) + 1$



514 Write a recursive formula, a_n , to describe the sequence graphed below.



515 The mean intelligence quotient (IQ) score is 100, with a standard deviation of 15, and the scores are normally distributed. Given this information, the approximate percentage of the population with an IQ greater than 130 is closest to

- 1) 2%
- 2) 31%
- 3) 48%
- 4) 95%

516 Factor completely over the set of integers:

$16x^4 - 81$. Sara graphed the polynomial $y = 16x^4 - 81$ and stated “All the roots of $y = 16x^4 - 81$ are real.” Is Sara correct? Explain your reasoning.

517 The terminal side of θ , an angle in standard

position, intersects the unit circle at $P\left(-\frac{1}{3}, -\frac{\sqrt{8}}{3}\right)$.

What is the value of $\sec \theta$?

- 1) -3
- 2) $-\frac{3\sqrt{8}}{8}$
- 3) $-\frac{1}{3}$
- 4) $-\frac{\sqrt{8}}{3}$

- 518 Given $\cos \theta = \frac{7}{25}$, where θ is an angle in standard position terminating in quadrant IV, and $\sin^2 \theta + \cos^2 \theta = 1$, what is the value of $\tan \theta$?

- 1) $-\frac{24}{25}$
- 2) $-\frac{24}{7}$
- 3) $\frac{24}{25}$
- 4) $\frac{24}{7}$

- 519 Brian deposited 1 cent into an empty non-interest bearing bank account on the first day of the month. He then additionally deposited 3 cents on the second day, 9 cents on the third day, and 27 cents on the fourth day. What would be the total amount of money in the account at the end of the 20th day if the pattern continued?

- 1) \$11,622,614.67
- 2) \$17,433,922.00
- 3) \$116,226,146.80
- 4) \$1,743,392,200.00

- 520 Given the following polynomials

$$x = (a + b + c)^2$$

$$y = a^2 + b^2 + c^2$$

$$z = ab + bc + ac$$

Which identity is true?

- 1) $x = y - z$
- 2) $x = y + z$
- 3) $x = y - 2z$
- 4) $x = y + 2z$

- 521 What is the solution set of the following system of equations?

$$y = 3x + 6$$

$$y = (x + 4)^2 - 10$$

- 1) $\{(-5, -9)\}$
- 2) $\{(5, 21)\}$
- 3) $\{(0, 6), (-5, -9)\}$
- 4) $\{(0, 6), (5, 21)\}$

- 522 Written in simplest form, $\frac{c^2 - d^2}{d^2 + cd - 2c^2}$ where $c \neq d$, is equivalent to

- 1) $\frac{c + d}{d + 2c}$
- 2) $\frac{c - d}{d + 2c}$
- 3) $\frac{-c - d}{d + 2c}$
- 4) $\frac{-c + d}{d + 2c}$

- 523 On average, college seniors graduating in 2012 could compute their growing student loan debt using the function $D(t) = 29,400(1.068)^t$, where t is time in years. Which expression is equivalent to $29,400(1.068)^t$ and could be used by students to identify an approximate daily interest rate on their loans?

- 1) $29,400 \left(1.068^{\frac{1}{365}} \right)^t$
- 2) $29,400 \left(\frac{1.068}{365} \right)^{365t}$
- 3) $29,400 \left(1 + \frac{0.068}{365} \right)^t$
- 4) $29,400 \left(1.068^{\frac{1}{365}} \right)^{365t}$

- 524 The probability that a resident of a housing community opposes spending money for community improvement on plumbing issues is 0.8. The probability that a resident favors spending money on improving walkways given that the resident opposes spending money on plumbing issues is 0.85. Determine the probability that a randomly selected resident opposes spending money on plumbing issues and favors spending money on walkways.

Algebra II Regents Exam Questions at Random Worksheet # 78

- 525 For the system shown below, what is the value of z ?

$$y = -2x + 14$$

$$3x - 4z = 2$$

$$3x - y = 16$$

- 1) 5
2) 2
3) 6
4) 4
- 526 An estimate of the number of milligrams of a medication in the bloodstream t hours after 400 mg has been taken can be modeled by the function below.

$$I(t) = 0.5t^4 + 3.45t^3 - 96.65t^2 + 347.7t,$$

$$\text{where } 0 \leq t \leq 6$$

Over what time interval does the amount of medication in the bloodstream strictly increase?

- 1) 0 to 2 hours
2) 0 to 3 hours
3) 2 to 6 hours
4) 3 to 6 hours
- 527 The graphs of the equations $y = x^2 + 4x - 1$ and $y + 3 = x$ are drawn on the same set of axes. One solution of this system is
- 1) $(-5, -2)$
2) $(-1, -4)$
3) $(1, 4)$
4) $(-2, -1)$

- 528 A researcher randomly divides 50 bean plants into two groups. He puts one group by a window to receive natural light and the second group under artificial light. He records the growth of the plants weekly. Which data collection method is described in this situation?

- 1) observational study
2) controlled experiment
3) survey
4) systematic sample

- 529 Express the fraction $\frac{2x^{\frac{3}{2}}}{(16x^4)^{\frac{1}{4}}}$ in simplest radical form.


- 530 Suppose two sets of test scores have the same mean, but different standard deviations, σ_1 and σ_2 , with $\sigma_2 > \sigma_1$. Which statement best describes the variability of these data sets?

- 1) Data set one has the greater variability.
2) Data set two has the greater variability.
3) The variability will be the same for each data set.
4) No conclusion can be made regarding the variability of either set.

- 531 Judith puts \$5000 into an investment account with interest compounded continuously. Which approximate annual rate is needed for the account to grow to \$9110 after 30 years?

- 1) 2%
2) 2.2%
3) 0.02%
4) 0.022%

- 532 On the grid below, graph the function $f(x) = x^3 - 6x^2 + 9x + 6$ on the domain $-1 \leq x \leq 4$.



533 What is the quotient when $10x^3 - 3x^2 - 7x + 3$ is divided by $2x - 1$?

- 1) $5x^2 + x + 3$
- 2) $5x^2 - x + 3$
- 3) $5x^2 - x - 3$
- 4) $5x^2 + x - 3$


534 The scores on a mathematics college-entry exam are normally distributed with a mean of 68 and standard deviation 7.2. Students scoring higher than one standard deviation above the mean will not be enrolled in the mathematics tutoring program. How many of the 750 incoming students can be expected to be enrolled in the tutoring program?

- 1) 631
- 2) 512
- 3) 238
- 4) 119

535 The function $N(t) = 100e^{-0.023t}$ models the number of grams in a sample of cesium-137 that remain after t years. On which interval is the sample's average rate of decay the fastest?

- 1) $[1, 10]$
- 2) $[10, 20]$
- 3) $[15, 25]$
- 4) $[1, 30]$

536 The zeros of a quartic polynomial function are 2, -2 , 4, and -4 . Use the zeros to construct a possible sketch of the function, on the set of axes below.



537 The expression $2 - \frac{x-1}{x+2}$ is equivalent to

- 1) $1 - \frac{3}{x+2}$
- 2) $1 + \frac{3}{x+2}$
- 3) $1 - \frac{1}{x+2}$
- 4) $1 + \frac{1}{x+2}$

538 The half-life of a radioactive substance is 15 years. Write an equation that can be used to determine the amount, $s(t)$, of 200 grams of this substance that remains after t years. Determine algebraically, to the *nearest year*, how long it will take for $\frac{1}{10}$ of this substance to remain.

539 At her job, Pat earns \$25,000 the first year and receives a raise of \$1000 each year. The explicit formula for the n th term of this sequence is $a_n = 25,000 + (n-1)1000$. Which rule best represents the equivalent recursive formula?

- 1) $a_n = 24,000 + 1000n$
- 2) $a_n = 25,000 + 1000n$
- 3) $a_1 = 25,000, a_n = a_{n-1} + 1000$
- 4) $a_1 = 25,000, a_n = a_{n+1} + 1000$

540 The Fahrenheit temperature, $F(t)$, of a heated object at time t , in minutes, can be modeled by the function below. F_s is the surrounding temperature, F_0 is the initial temperature of the object, and k is a constant.

$$F(t) = F_s + (F_0 - F_s)e^{-kt}$$

Coffee at a temperature of 195°F is poured into a container. The room temperature is kept at a constant 68°F and $k = 0.05$. Coffee is safe to drink when its temperature is, at most, 120°F . To the *nearest minute*, how long will it take until the coffee is safe to drink?

- 1) 7
- 2) 10
- 3) 11
- 4) 18


541 Selected values for the functions f and g are shown in the tables below.

x	f(x)		x	g(x)
-3.12	-4.88		-2.01	-1.01
0	-6		0	0.58
1.23	-4.77		8.52	2.53
8.52	2.53		13.11	3.01
9.01	3.01		16.52	3.29

A solution to the equation $f(x) = g(x)$ is


- 1) 0 3) 3.01
2) 2.53 4) 8.52

542 In a random sample of 250 men in the United States, age 21 or older, 139 are married. The graph below simulated samples of 250 men, 200 times, assuming that 139 of the men are married.



- Based on the simulation, create an interval in which the middle 95% of the number of married men may fall. Round your answer to the *nearest integer*.
- A study claims "50 percent of men 21 and older in the United States are married." Do your results from part a contradict this claim? Explain.

544 Write an equation for a sine function with an amplitude of 2 and a period of $\frac{\pi}{2}$. On the grid below, sketch the graph of the equation in the interval 0 to 2π .



545 Solve the following system of equations algebraically for all values of a , b , and c .

$$a + 4b + 6c = 23$$

$$a + 2b + c = 2$$

$$6b + 2c = a + 14$$

543 What is the solution set of the equation

$$\frac{10}{x^2 - 2x} + \frac{4}{x} = \frac{5}{x - 2}?$$

- 1) $\{0, 2\}$
- 2) $\{0\}$
- 3) $\{2\}$
- 4) $\{\}$

- 546 The Beaufort Wind Scale was devised by British Rear Admiral Sir Francis Beaufort, in 1805 based upon observations of the effects of the wind. Beaufort numbers, B , are determined by the equation $B = 1.69\sqrt{s + 4.45} - 3.49$, where s is the speed of the wind in mph, and B is rounded to the nearest integer from 0 to 12.

Beaufort Wind Scale	
Beaufort Number	Force of Wind
0	Calm
1	Light air
2	Light breeze
3	Gentle breeze
4	Moderate breeze
5	Fresh breeze
6	Steady breeze
7	Moderate gale
8	Fresh gale
9	Strong gale
10	Whole gale
11	Storm
12	Hurricane


Using the table above, classify the force of wind at a speed of 30 mph. Justify your answer. In 1946, the scale was extended to accommodate strong hurricanes. A strong hurricane received a B value of exactly 15. Algebraically determine the value of s , to the *nearest mph*. Any B values that round to 10 receive a Beaufort number of 10. Using technology, find an approximate range of wind speeds, to the *nearest mph*, associated with a Beaufort number of 10.

- 547 Stephanie found that the number of white-winged cross bills in an area can be represented by the formula $C = 550(1.08)^t$, where t represents the number of years since 2010. Which equation correctly represents the number of white-winged cross bills in terms of the monthly rate of population growth?
- 1) $C = 550(1.00643)^t$
 - 2) $C = 550(1.00643)^{12t}$
 - 3) $C = 550(1.00643)^{\frac{t}{12}}$
 - 4) $C = 550(1.00643)^{t+12}$
- 548 If \$5000 is put into a savings account that pays 3.5% interest compounded monthly, how much money, to the *nearest ten cents*, would be in that account after 6 years, assuming no money was added or withdrawn?
- 1) \$5177.80
 - 2) \$5941.30
 - 3) \$6146.30
 - 4) \$6166.50
- 549 Which expression can be rewritten as $(x+7)(x-1)$?
- 1) $(x+3)^2 - 16$
 - 2) $(x+3)^2 - 10(x+3) - 2(x+3) + 20$
 - 3) $\frac{(x-1)(x^2 - 6x - 7)}{(x+1)}$
 - 4) $\frac{(x+7)(x^2 + 4x + 3)}{(x+3)}$

- 550 Joseph was curious to determine if scent improves memory. A test was created where better memory is indicated by higher test scores. A controlled experiment was performed where one group was given the test on scented paper and the other group was given the test on unscented paper. The summary statistics from the experiment are given below.

	Scented Paper	Unscented Paper
\bar{x}	23	18
s_x	2.898	2.408

Calculate the difference in means in the experimental test grades (scented -unscented). A simulation was conducted in which the subjects' scores were rerandomized into two groups 1000 times. The differences of the group means were calculated each time. The results are shown below.



Use the simulation results to determine the interval representing the middle 95% of the difference in means, to the nearest hundredth. Is the difference in means in Joseph's experiment statistically significant based on the simulation? Explain.

- 551 Write a recursive formula for the sequence
6, 9, 13.5, 20.25, ...

Algebra II Regents Exam Questions at Random Worksheet # 83

- 552 The scores of a recent test taken by 1200 students had an approximately normal distribution with a mean of 225 and a standard deviation of 18. Determine the number of students who scored between 200 and 245.
- 553 Given $P(x) = x^3 - 3x^2 - 2x + 4$, which statement is true?
- 1) $(x - 1)$ is a factor because $P(-1) = 2$.
 - 2) $(x + 1)$ is a factor because $P(-1) = 2$.
 - 3) $(x + 1)$ is a factor because $P(1) = 0$.
 - 4) $(x - 1)$ is a factor because $P(1) = 0$.
- 554 What is the equation of the directrix for the parabola $-8(y - 3) = (x + 4)^2$?
- 1) $y = 5$
 - 2) $y = 1$
 - 3) $y = -2$
 - 4) $y = -6$
- 555 The first term of a geometric sequence is 8 and the fourth term is 216. What is the sum of the first 12 terms of the corresponding series?
- 1) 236,192
 - 2) 708,584
 - 3) 2,125,760
 - 4) 6,377,288
- 556 The recursive formula to describe a sequence is shown below.
- $$a_1 = 3$$
- $$a_n = 1 + 2a_{n-1}$$
- State the first four terms of this sequence. Can this sequence be represented using an explicit geometric formula? Justify your answer.
- 557 A sociologist reviews randomly selected surveillance videos from a public park over a period of several years and records the amount of time people spent on a smartphone. The statistical procedure the sociologist used is called
- 1) a census
 - 2) an experiment
 - 3) an observational study
 - 4) a sample survey
- 558 The roots of the equation $3x^2 + 2x = -7$ are
- 1) $-2, -\frac{1}{3}$
 - 2) $-\frac{7}{3}, 1$
 - 3) $-\frac{1}{3} \pm \frac{2i\sqrt{5}}{3}$
 - 4) $-\frac{1}{3} \pm \frac{\sqrt{11}}{3}$
- 559 The expression $6 - (3x - 2i)^2$ is equivalent to
- 1) $-9x^2 + 12xi + 10$
 - 2) $9x^2 - 12xi + 2$
 - 3) $-9x^2 + 10$
 - 4) $-9x^2 + 12xi - 4i + 6$
- 560 When factoring to reveal the roots of the equation $x^3 + 2x^2 - 9x - 18 = 0$, which equations can be used?
- I. $x^2(x + 2) - 9(x + 2) = 0$
 - II. $x(x^2 - 9) + 2(x^2 - 9) = 0$
 - III. $(x - 2)(x^2 - 9) = 0$
- 1) I and II, only
 - 2) I and III, only
 - 3) II and III, only
 - 4) I, II, and III
- 561 Which function is even?
- 1) $f(x) = \sin x$
 - 2) $f(x) = x^2 - 4$
 - 3) $f(x) = |x - 2| + 5$
 - 4) $f(x) = x^4 + 3x^3 + 4$
- 562 Justify why $\frac{\sqrt[3]{x^2 y^5}}{\sqrt[4]{x^3 y^4}}$ is equivalent to $x^{\frac{-1}{12}} y^{\frac{2}{3}}$ using properties of rational exponents, where $x \neq 0$ and $y \neq 0$.

- 563 Juan and Filipe practice at the driving range before playing golf. The number of wins and corresponding practice times for each player are shown in the table below.

	Juan Wins	Felipe Wins
Short Practice Time	8	10
Long Practice Time	15	12

Given that the practice time was long, determine the exact probability that Filipe wins the next match. Determine whether or not the two events “Filipe wins” and “long practice time” are independent. Justify your answer.

- 564 What is the inverse of $f(x) = -6(x - 2)$?

- 1) $f^{-1}(x) = -2 - \frac{x}{6}$
- 2) $f^{-1}(x) = 2 - \frac{x}{6}$
- 3) $f^{-1}(x) = \frac{1}{-6(x - 2)}$
- 4) $f^{-1}(x) = 6(x + 2)$

- 565 Elizabeth tried to find the product of $(2 + 4i)$ and $(3 - i)$, and her work is shown below.

$$\begin{aligned}
 &(2 + 4i)(3 - i) \\
 &= 6 - 2i + 12i - 4i^2 \\
 &= 6 + 10i - 4i^2 \\
 &= 6 + 10i - 4(1) \\
 &= 6 + 10i - 4 \\
 &= 2 + 10i
 \end{aligned}$$


Identify the error in the process shown and determine the correct product of $(2 + 4i)$ and $(3 - i)$.

- 566 Kenzie believes that for $x \geq 0$, the expression $\left(\sqrt[7]{x^2}\right)\left(\sqrt[5]{x^3}\right)$ is equivalent to $\sqrt[35]{x^6}$. Is she correct? Justify your response algebraically.

- 567 If $p(x) = 2\ln(x) - 1$ and $m(x) = \ln(x + 6)$, then what is the solution for $p(x) = m(x)$?

- 1) 1.65
- 2) 3.14
- 3) 5.62
- 4) no solution

- 568 Irma initially ran one mile in over ten minutes. She then began a training program to reduce her one-mile time. She recorded her one-mile time once a week for twelve consecutive weeks, as modeled in the graph below.



Which statement regarding Irma's one-mile training program is correct?

- 1) Her one-mile speed increased as the number of weeks increased.
- 2) Her one-mile speed decreased as the number of weeks increased.
- 3) If the trend continues, she will run under a six-minute mile by week thirteen.
- 4) She reduced her one-mile time the most between weeks ten and twelve.

- 569 If $f(x) = \log_3 x$ and $g(x)$ is the image of $f(x)$ after a translation five units to the left, which equation represents $g(x)$?


- 1) $g(x) = \log_3(x + 5)$
- 2) $g(x) = \log_3 x + 5$
- 3) $g(x) = \log_3(x - 5)$
- 4) $g(x) = \log_3 x - 5$

Algebra II Regents Exam Questions at Random Worksheet # 85

- 570 A radio station claims to its advertisers that the mean number of minutes commuters listen to the station is 30. The station conducted a survey of 500 of their listeners who commute. The sample statistics are shown below.

\bar{x}	29.11
s_x	20.718

A simulation was run 1000 times based upon the results of the survey. The results of the simulation appear below.



Based on the simulation results, is the claim that commuters listen to the station on average 30 minutes plausible? Explain your response including an interval containing the middle 95% of the data, rounded to the *nearest hundredth*.

- 571 The expression $(x^2 + y^2)^2$ is *not* equivalent to

- 1) $(x^2 - y^2)^2 + (2xy)^2$
- 2) $(x + y)^4 + 2(xy)^2$
- 3) $x^2(x^2 + 2y^2) + (y^2)^2$
- 4) $(2x^2 + y^2)^2 - (3x^4 + 2x^2y^2)$

- 573 a) Algebraically determine the roots, in simplest $a + bi$ form, to the equation below.

$$x^2 - 2x + 7 = 4x - 10$$

- b) Consider the system of equations below.

$$y = x^2 - 2x + 7$$

$$y = 4x - 10$$

The graph of this system confirms the solution from part a is imaginary. Explain why.

- 572 If $\cos \theta = -\frac{3}{4}$ and θ is in Quadrant III, then $\sin \theta$ is equivalent to

- 1) $-\frac{\sqrt{7}}{4}$
- 2) $\frac{\sqrt{7}}{4}$
- 3) $-\frac{5}{4}$
- 4) $\frac{5}{4}$

- 574 The solutions to the equation $5x^2 - 2x + 13 = 9$ are

- 1) $\frac{1}{5} \pm \frac{\sqrt{21}}{5}$
- 2) $\frac{1}{5} \pm \frac{\sqrt{19}}{5}i$
- 3) $\frac{1}{5} \pm \frac{\sqrt{66}}{5}i$
- 4) $\frac{1}{5} \pm \frac{\sqrt{66}}{5}$

- 575 Julia deposits \$2000 into a savings account that earns 4% interest per year. The exponential function that models this savings account is $y = 2000(1.04)^t$, where t is the time in years. Which equation correctly represents the amount of money in her savings account in terms of the monthly growth rate?

- 1) $y = 166.67(1.04)^{0.12t}$
- 2) $y = 2000(1.01)^t$
- 3) $y = 2000(1.0032737)^{12t}$
- 4) $y = 166.67(1.0032737)^t$


- 576 For positive values of x , which expression is

equivalent to $\sqrt{16x^2} \cdot x^{\frac{2}{3}} + \sqrt[3]{8x^5}$

- 1) $6\sqrt[5]{x^3}$
- 2) $6\sqrt[3]{x^5}$
- 3) $4\sqrt[3]{x^2} + 2\sqrt[3]{x^5}$
- 4) $4\sqrt{x^3} + 2\sqrt[5]{x^3}$

- 577 Write $-\frac{1}{2}i^3(\sqrt{-9} - 4) - 3i^2$ in simplest $a + bi$ form.


- 578 The graph of $y = f(x)$ is shown below.




Which expression defines $f(x)$?

- 1) $2x$
- 2) $5(2^x)$
- 3) $5(2^{\frac{x}{2}})$
- 4) $5(2^{2x})$


- 579 A 4th degree polynomial has zeros -5 , 3 , i , and $-i$. Which graph could represent the function defined by this polynomial?




1)



2)



3)




4)

- 580 Solve the equation $2x^2 + 5x + 8 = 0$. Express the answer in $a + bi$ form.

581 The solution of $87e^{0.3x} = 5918$, to the nearest thousandth, is

- 1) 0.583
- 2) 1.945
- 3) 4.220
- 4) 14.066

582 The graph below represents national and New York State average gas prices.



If New York State's gas prices are modeled by $G(x)$ and $C > 0$, which expression best approximates the national average x months from August 2014?

- 1) $G(x + C)$
- 2) $G(x) + C$
- 3) $G(x - C)$
- 4) $G(x) - C$


583 What is the inverse of $f(x) = x^3 - 2$?

- 1) $f^{-1}(x) = \sqrt[3]{x} + 2$
- 2) $f^{-1}(x) = \pm \sqrt[3]{x} + 2$
- 3) $f^{-1}(x) = \sqrt[3]{x + 2}$
- 4) $f^{-1}(x) = \pm \sqrt[3]{x + 2}$

584 The expression $(x + a)(x + b)$ can *not* be written as

- 1) $a(x + b) + x(x + b)$
- 2) $x^2 + abx + ab$
- 3) $x^2 + (a + b)x + ab$
- 4) $x(x + a) + b(x + a)$

585 The function $f(x) = a \cos bx + c$ is plotted on the graph shown below.



What are the values of a , b , and c ?

- 1) $a = 2, b = 6, c = 3$
- 2) $a = 2, b = 3, c = 1$
- 3) $a = 4, b = 6, c = 5$
- 4) $a = 4, b = \frac{\pi}{3}, c = 3$





586 What is the inverse of $f(x) = \frac{x}{x+2}$, where $x \neq -2$?

- 1) $f^{-1}(x) = \frac{2x}{x-1}$
- 2) $f^{-1}(x) = \frac{-2x}{x-1}$
- 3) $f^{-1}(x) = \frac{x}{x-2}$
- 4) $f^{-1}(x) = \frac{-x}{x-2}$

587 For $x \geq 0$, which equation is *false*?

- 1) $(x^{\frac{3}{2}})^2 = 4\sqrt{x^3}$
- 2) $(x^3)^{\frac{1}{4}} = 4\sqrt{x^3}$
- 3) $(x^{\frac{3}{2}})^{\frac{1}{2}} = 4\sqrt{x^3}$
- 4) $(x^{\frac{2}{3}})^2 = 3\sqrt{x^4}$

588 Which sketch best represents the graph of $x = 3^y$?

- 1) 
- 2) 
- 3) 
- 4) 

589 After examining the functions $f(x) = \ln(x + 2)$ and $g(x) = e^{x-1}$ over the interval $(-2, 3]$, Lexi determined that the correct number of solutions to the equation $f(x) = g(x)$ is


- 1) 1
2) 2
3) 3
4) 0

590 A major car company analyzes its revenue, $R(x)$, and costs $C(x)$, in millions of dollars over a fifteen-year period. The company represents its revenue and costs as a function of time, in years, x , using the given functions.

$$R(x) = 550x^3 - 12,000x^2 + 83,000x + 7000$$

$$C(x) = 880x^3 - 21,000x^2 + 150,000x - 160,000$$

The company's profits can be represented as the difference between its revenue and costs. Write the profit function, $P(x)$, as a polynomial in standard form. Graph $y = P(x)$ on the set of axes below over the domain $2 \leq x \leq 16$.



Over the given domain, state when the company was the least profitable and the most profitable, to the nearest year. Explain how you determined your answer.


591 Which expression is equivalent to

$$x^6y^4(x^4 - 16) - 9(x^4 - 16)?$$

- 1) $x^{10}y^4 - 16x^6y^4 - 9x^4 - 144$
2) $(x^6y^4 - 9)(x + 2)^3(x - 2)$
3) $(x^3y^2 + 3)(x^3y^2 - 3)(x + 2)^2(x - 2)^2$
4) $(x^3y^2 + 3)(x^3y^2 - 3)(x^2 + 4)(x^2 - 4)$


Algebra II Regents at Random Worksheets

- 592 The graph below represents the height above the ground, h , in inches, of a point on a triathlete's bike wheel during a training ride in terms of time, t , in seconds.



Identify the period of the graph and describe what the period represents in this context.

- 593 Given $f(9) = -2$, which function can be used to generate the sequence $-8, -7.25, -6.5, -5.75, \dots$?
- 1) $f(n) = -8 + 0.75n$
 - 2) $f(n) = -8 - 0.75(n - 1)$
 - 3) $f(n) = -8.75 + 0.75n$
 - 4) $f(n) = -0.75 + 8(n - 1)$
- 594 Explain how $(-8)^{\frac{4}{3}}$ can be evaluated using properties of rational exponents to result in an integer answer.
- 595 What is the completely factored form of $k^4 - 4k^2 + 8k^3 - 32k + 12k^2 - 48$?
- 1) $(k - 2)(k - 2)(k + 3)(k + 4)$
 - 2) $(k - 2)(k - 2)(k + 6)(k + 2)$
 - 3) $(k + 2)(k - 2)(k + 3)(k + 4)$
 - 4) $(k + 2)(k - 2)(k + 6)(k + 2)$
- 596 A house purchased 5 years ago for \$100,000 was just sold for \$135,000. Assuming exponential growth, approximate the annual growth rate, to the nearest percent.
- 597 Given $r(x) = x^3 - 4x^2 + 4x - 6$, find the value of $r(2)$. What does your answer tell you about $x - 2$ as a factor of $r(x)$? Explain.
- 598 Solve the equation $\sqrt{2x - 7} + x = 5$ algebraically, and justify the solution set.
- 599 On the axes below, sketch a possible function $p(x) = (x - a)(x - b)(x + c)$, where a , b , and c are positive, $a > b$, and $p(x)$ has a positive y -intercept of d . Label all intercepts.



- 600 The results of a survey of the student body at Central High School about television viewing preferences are shown below.


	Comedy Series	Drama Series	Reality Series	Total
Males	95	65	70	230
Females	80	70	110	260
Total	175	135	180	490

Are the events “student is a male” and “student prefers reality series” independent of each other? Justify your answer.

- 601 The eighth and tenth terms of a sequence are 64 and 100. If the sequence is either arithmetic or geometric, the ninth term can *not* be
- 1) -82
 - 2) -80
 - 3) 80
 - 4) 82
- 602 Which equation has $1 - i$ as a solution?
- 1) $x^2 + 2x - 2 = 0$
 - 2) $x^2 + 2x + 2 = 0$
 - 3) $x^2 - 2x - 2 = 0$
 - 4) $x^2 - 2x + 2 = 0$
- 603 Given $f(x) = 3x^2 + 7x - 20$ and $g(x) = x - 2$, state the quotient and remainder of $\frac{f(x)}{g(x)}$, in the form $q(x) + \frac{r(x)}{g(x)}$.
- 604 To solve $\frac{2x}{x-2} - \frac{11}{x} = \frac{8}{x^2 - 2x}$, Ren multiplied both sides by the least common denominator. Which statement is true?
- 1) 2 is an extraneous solution.
 - 2) $\frac{7}{2}$ is an extraneous solution.
 - 3) 0 and 2 are extraneous solutions.
 - 4) This equation does not contain any extraneous solutions.
- 605 Seth’s parents gave him \$5000 to invest for his 16th birthday. He is considering two investment options. Option A will pay him 4.5% interest compounded annually. Option B will pay him 4.6% compounded quarterly. Write a function of option A and option B that calculates the value of each account after n years. Seth plans to use the money after he graduates from college in 6 years. Determine how much more money option B will earn than option A to the *nearest cent*. Algebraically determine, to the *nearest tenth of a year*, how long it would take for option B to double Seth’s initial investment.
- 606 The expression $\frac{6x^3 + 17x^2 + 10x + 2}{2x + 3}$ equals
- 1) $3x^2 + 4x - 1 + \frac{5}{2x + 3}$
 - 2) $6x^2 + 8x - 2 + \frac{5}{2x + 3}$
 - 3) $6x^2 - x + 13 - \frac{37}{2x + 3}$
 - 4) $3x^2 + 13x + \frac{49}{2} + \frac{151}{2x + 3}$
- 607 The expression $\frac{x^3 + 2x^2 + x + 6}{x + 2}$ is equivalent to
- 1) $x^2 + 3$
 - 2) $x^2 + 1 + \frac{4}{x + 2}$
 - 3) $2x^2 + x + 6$
 - 4) $2x^2 + 1 + \frac{4}{x + 2}$

- 608 A polynomial equation of degree three, $p(x)$, is used to model the volume of a rectangular box. The graph of $p(x)$ has x intercepts at -2 , 10 , and 14 . Which statements regarding $p(x)$ could be true?
- The equation of $p(x) = (x - 2)(x + 10)(x + 14)$.
 - The equation of $p(x) = -(x + 2)(x - 10)(x - 14)$.
 - The maximum volume occurs when $x = 10$.
 - The maximum volume of the box is approximately 56.
- A and C
 - A and D
 - B and C
 - B and D

- 609 A study conducted in 2004 in New York City found that 212 out of 1334 participants had hypertension. Kim ran a simulation of 100 studies based on these data. The output of the simulation is shown in the diagram below.




At a 95% confidence level, the proportion of New York City residents with hypertension and the margin of error are closest to

- proportion $\approx .16$; margin of error $\approx .01$
 - proportion $\approx .16$; margin of error $\approx .02$
 - proportion $\approx .01$; margin of error $\approx .16$
 - proportion $\approx .02$; margin of error $\approx .16$
- 610 Algebraically prove that the difference of the squares of any two consecutive integers is an odd integer.
- 611 Write $\sqrt[3]{x} \cdot \sqrt{x}$ as a single term with a rational exponent.
- 612 Given that $\sin^2 \theta + \cos^2 \theta = 1$ and $\sin \theta = -\frac{\sqrt{2}}{5}$, what is a possible value of $\cos \theta$?
- $\frac{5 + \sqrt{2}}{5}$
 - $\frac{\sqrt{23}}{5}$
 - $\frac{3\sqrt{3}}{5}$
 - $\frac{\sqrt{35}}{5}$
- 613 An equation to represent the value of a car after t months of ownership is $v = 32,000(0.81)^{\frac{t}{12}}$. Which statement is *not* correct?
- The car lost approximately 19% of its value each month.
 - The car maintained approximately 98% of its value each month.
 - The value of the car when it was purchased was \$32,000.
 - The value of the car 1 year after it was purchased was \$25,920.
- 614 What is the solution set of the equation $\frac{3x+25}{x+7} - 5 = \frac{3}{x}$?
- $\left\{\frac{3}{2}, 7\right\}$
 - $\left\{\frac{7}{2}, -3\right\}$
 - $\left\{-\frac{3}{2}, 7\right\}$
 - $\left\{-\frac{7}{2}, -3\right\}$
- 615 Which expression has been rewritten correctly to form a true statement?
- $(x+2)^2 + 2(x+2) - 8 = (x+6)x$
 - $x^4 + 4x^2 + 9x^2y^2 - 36y^2 = (x+3y)^2(x-2)^2$
 - $x^3 + 3x^2 - 4xy^2 - 12y^2 = (x-2y)(x+3)^2$
 - $(x^2-4)^2 - 5(x^2-4) - 6 = (x^2-7)(x^2-6)$


616 Which equation represents a parabola with the focus at $(0, -1)$ and the directrix of $y = 1$?

- 1) $x^2 = -8y$
- 2) $x^2 = -4y$
- 3) $x^2 = 8y$
- 4) $x^2 = 4y$

617 Which sinusoid has the greatest amplitude?



- 1)
- 2) $y = 3 \sin(\theta - 3) + 5$




- 3)
- 4) $y = -5 \sin(\theta - 1) - 3$

618 Pedro and Bobby each own an ant farm. Pedro starts with 100 ants and says his farm is growing exponentially at a rate of 15% per month. Bobby starts with 350 ants and says his farm is steadily decreasing by 5 ants per month. Assuming both boys are accurate in describing the population of their ant farms, after how many months will they both have approximately the same number of ants?

- 1) 7
- 2) 8
- 3) 13
- 4) 36

619 For the function $f(x) = (x - 3)^3 + 1$, find $f^{-1}(x)$.

620 Drugs break down in the human body at different rates and therefore must be prescribed by doctors carefully to prevent complications, such as overdosing. The breakdown of a drug is represented by the function $N(t) = N_0(e)^{-rt}$, where $N(t)$ is the amount left in the body, N_0 is the initial dosage, r is the decay rate, and t is time in hours. Patient A, $A(t)$, is given 800 milligrams of a drug with a decay rate of 0.347. Patient B, $B(t)$, is given 400 milligrams of another drug with a decay rate of 0.231. Write two functions, $A(t)$ and $B(t)$, to represent the breakdown of the respective drug given to each patient. Graph each function on the set of axes below.



To the *nearest hour*, t , when does the amount of the given drug remaining in patient B begin to exceed the amount of the given drug remaining in patient A? The doctor will allow patient A to take another 800 milligram dose of the drug once only 15% of the original dose is left in the body. Determine, to the *nearest tenth of an hour*, how long patient A will have to wait to take another 800 milligram dose of the drug.

- 621 The x -value of which function's x -intercept is larger, f or h ? Justify your answer.

$$f(x) = \log(x - 4)$$

x	$h(x)$
-1	6
0	4
1	2
2	0
3	-2

- 622 The speed of a tidal wave, s , in hundreds of miles per hour, can be modeled by the equation $s = \sqrt{t} - 2t + 6$, where t represents the time from its origin in hours. Algebraically determine the time when $s = 0$. How much faster was the tidal wave traveling after 1 hour than 3 hours, to the nearest mile per hour? Justify your answer.

- 623 In 2010, the population of New York State was approximately 19,378,000 with an annual growth rate of 1.5%. Assuming the growth rate is maintained for a large number of years, which equation can be used to predict the population of New York State t years after 2010?

- 1) $P_t = 19,378,000(1.5)^t$
- 2) $P_0 = 19,378,000$
 $P_t = 19,378,000 + 1.015P_{t-1}$
- 3) $P_t = 19,378,000(1.015)^{t-1}$
- 4) $P_0 = 19,378,000$
 $P_t = 1.015P_{t-1}$

- 624 Relative to the graph of $y = 3 \sin x$, what is the shift of the graph of $y = 3 \sin\left(x + \frac{\pi}{3}\right)$?

- 1) $\frac{\pi}{3}$ right
- 2) $\frac{\pi}{3}$ left
- 3) $\frac{\pi}{3}$ up
- 4) $\frac{\pi}{3}$ down

- 625 If $g(c) = 1 - c^2$ and $m(c) = c + 1$, then which statement is *not* true?

- 1) $g(c) \cdot m(c) = 1 + c - c^2 - c^3$
- 2) $g(c) + m(c) = 2 + c - c^2$
- 3) $m(c) - g(c) = c + c^2$
- 4) $\frac{m(c)}{g(c)} = \frac{-1}{1 - c}$


- 626 The expression $\frac{-3x^2 - 5x + 2}{x^3 + 2x^2}$ can be rewritten as

- 1) $\frac{-3x - 3}{x^2 + 2x}$
- 2) $\frac{-3x - 1}{x^2}$
- 3) $-3x^{-1} + 1$
- 4) $-3x^{-1} + x^{-2}$

- 627 Which expression is equivalent to $(3k - 2i)^2$, where i is the imaginary unit?

- 1) $9k^2 - 4$
- 2) $9k^2 + 4$
- 3) $9k^2 - 12ki - 4$
- 4) $9k^2 - 12ki + 4$

- 628 Gabriel performed an experiment to see if planting 13 tomato plants in black plastic mulch leads to larger tomatoes than if 13 plants are planted without mulch. He observed that the average weight of the tomatoes from tomato plants grown in black plastic mulch was 5 ounces greater than those from the plants planted without mulch. To determine if the observed difference is statistically significant, he rerandomized the tomato groups 100 times to study these random differences in the mean weights. The output of his simulation is summarized in the dotplot below.



Given these results, what is an appropriate inference that can be drawn?


- | | |
|---|---|
| <p>1) There was no effect observed between the two groups.</p> <p>2) There was an effect observed that could be due to the random assignment of plants to the groups.</p> | <p>3) There is strong evidence to support the hypothesis that tomatoes from plants planted in black plastic mulch are larger than those planted without mulch.</p> <p>4) There is strong evidence to support the hypothesis that tomatoes from plants planted without mulch are larger than those planted in black plastic mulch.</p> |
|---|---|
-
- | | |
|--|---|
| <p>629 The Rickerts decided to set up an account for their daughter to pay for her college education. The day their daughter was born, they deposited \$1000 in an account that pays 1.8% compounded annually. Beginning with her first birthday, they deposit an additional \$750 into the account on each of her birthdays. Which expression correctly represents the amount of money in the account n years after their daughter was born?</p> <p>1) $a_n = 1000(1.018)^n + 750$</p> <p>2) $a_n = 1000(1.018)^n + 750n$</p> <p>3) $a_0 = 1000$</p> <p style="padding-left: 40px;">$a_n = a_{n-1}(1.018) + 750$</p> <p>4) $a_0 = 1000$</p> <p style="padding-left: 40px;">$a_n = a_{n-1}(1.018) + 750n$</p> | <p>630 If $p(x) = ab^x$ and $r(x) = cd^x$, then $p(x) \bullet r(x)$ equals</p> <p>1) $ac(b + d)^x$</p> <p>2) $ac(b + d)^{2x}$</p> <p>3) $ac(bd)^x$</p> <p>4) $ac(bd)^{x^2}$</p> <p>631 Which binomial is a factor of $x^4 - 4x^2 - 4x + 8$?</p> <p>1) $x - 2$</p> <p>2) $x + 2$</p> <p>3) $x - 4$</p> <p>4) $x + 4$</p> |
|--|---|

- 632 Using a microscope, a researcher observed and recorded the number of bacteria spores on a large sample of uniformly sized pieces of meat kept at room temperature. A summary of the data she recorded is shown in the table below.

Hours (x)	Average Number of Spores (y)
0	4
0.5	10
1	15
2	60
3	260
4	1130
6	16,380


Using these data, write an exponential regression equation, rounding all values to the *nearest thousandth*. The researcher knows that people are likely to suffer from food-borne illness if the number of spores exceeds 100. Using the exponential regression equation, determine the maximum amount of time, to the *nearest quarter hour*, that the meat can be kept at room temperature safely.

- 633 Which equation represents the set of points equidistant from line ℓ and point R shown on the graph below?



- 1) $y = -\frac{1}{8}(x+2)^2 + 1$
- 2) $y = -\frac{1}{8}(x+2)^2 - 1$
- 3) $y = -\frac{1}{8}(x-2)^2 + 1$
- 4) $y = -\frac{1}{8}(x-2)^2 - 1$


- 634 The price of a postage stamp in the years since the end of World War I is shown in the scatterplot below.



The equation that best models the price, in cents, of a postage stamp based on these data is


- 1) $y = 0.59x - 14.82$
- 2) $y = 1.04(1.43)^x$
- 3) $y = 1.43(1.04)^x$
- 4) $y = 24 \sin(14x) + 25$

- 635 Charlie's Automotive Dealership is considering implementing a new check-in procedure for customers who are bringing their vehicles for routine maintenance. The dealership will launch the procedure if 50% or more of the customers give the new procedure a favorable rating when compared to the current procedure. The dealership devises a simulation based on the minimal requirement that 50% of the customers prefer the new procedure. Each dot on the graph below represents the proportion of the customers who preferred the new check-in procedure, each of sample size 40, simulated 100 times.




Assume the set of data is approximately normal and the dealership wants to be 95% confident of its results. Determine an interval containing the plausible sample values for which the dealership will launch the new procedure. Round your answer to the *nearest hundredth*. Forty customers are selected randomly to undergo the new check-in procedure and the proportion of customers who prefer the new procedure is 32.5%. The dealership decides *not* to implement the new check-in procedure based on the results of the study. Use statistical evidence to explain this decision.

- 636 On the axes below, graph *one* cycle of a cosine function with amplitude 3, period $\frac{\pi}{2}$, midline $y = -1$, and passing through the point (0,2).



- 637 A manufacturing company has developed a cost model, $C(x) = 0.15x^3 + 0.01x^2 + 2x + 120$, where x is the number of items sold, in thousands. The sales price can be modeled by $S(x) = 30 - 0.01x$. Therefore, revenue is modeled by $R(x) = x \cdot S(x)$. The company's profit, $P(x) = R(x) - C(x)$, could be modeled by
- 1) $0.15x^3 + 0.02x^2 - 28x + 120$
 - 2) $-0.15x^3 - 0.02x^2 + 28x - 120$
 - 3) $-0.15x^3 + 0.01x^2 - 2.01x - 120$
 - 4) $-0.15x^3 + 32x + 120$
- 638 In 2013, approximately 1.6 million students took the Critical Reading portion of the SAT exam. The mean score, the modal score, and the standard deviation were calculated to be 496, 430, and 115, respectively. Which interval reflects 95% of the Critical Reading scores?
- 1) 430 ± 115
 - 2) 430 ± 230
 - 3) 496 ± 115
 - 4) 496 ± 230

- 639 The ocean tides near Carter Beach follow a repeating pattern over time, with the amount of time between each low and high tide remaining relatively constant. On a certain day, low tide occurred at 8:30 a.m. and high tide occurred at 3:00 p.m. At high tide, the water level was 12 inches above the average local sea level; at low tide it was 12 inches below the average local sea level. Assume that high tide and low tide are the maximum and minimum water levels each day, respectively. Write a cosine function of the form $f(t) = A \cos(Bt)$, where A and B are real numbers, that models the water level, $f(t)$, in inches above or below the average Carter Beach sea level, as a function of the time measured in t hours since 8:30 a.m. On the grid below, graph one cycle of this function.



People who fish in Carter Beach know that a certain species of fish is most plentiful when the water level is increasing. Explain whether you would recommend fishing for this species at 7:30 p.m. or 10:30 p.m. using evidence from the given context.

- 640 In New York State, the minimum wage has grown exponentially. In 1966, the minimum wage was \$1.25 an hour and in 2015, it was \$8.75. Algebraically determine the rate of growth to the nearest percent.

- 641 Simon lost his library card and has an overdue library book. When the book was 5 days late, he owed \$2.25 to replace his library card and pay the fine for the overdue book. When the book was 21 days late, he owed \$6.25 to replace his library card and pay the fine for the overdue book. Suppose the total amount Simon owes when the book is n days late can be determined by an arithmetic sequence. Determine a formula for a_n , the n th term of this sequence. Use the formula to determine the amount of money, in dollars, Simon needs to pay when the book is 60 days late.

- 642 Verify the following Pythagorean identity for all values of x and y :

$$(x^2 + y^2)^2 = (x^2 - y^2)^2 + (2xy)^2$$

- 643 Explain how $\left(3^{\frac{1}{5}}\right)^2$ can be written as the equivalent radical expression $\sqrt[5]{9}$.

- 644 What is the solution to $8(2^{x+3}) = 48$?

- 1) $x = \frac{\ln 6}{\ln 2} - 3$
- 2) $x = 0$
- 3) $x = \frac{\ln 48}{\ln 16} - 3$
- 4) $x = \ln 4 - 3$

- 645 According to a pricing website, Indroid phones lose 58% of their cash value over 1.5 years. Which expression can be used to estimate the value of a \$300 Indroid phone in 1.5 years?

- 1) $300e^{-0.87}$
- 2) $300e^{-0.63}$
- 3) $300e^{-0.58}$
- 4) $300e^{-0.42}$

- 646 Rewrite the expression

$$\left(4x^2 + 5x\right)^2 - 5\left(4x^2 + 5x\right) - 6$$

as a product of four linear factors.

647 The expression $\frac{4x^3 + 5x + 10}{2x + 3}$ is equivalent to

- 1) $2x^2 + 3x - 7 + \frac{31}{2x + 3}$
- 2) $2x^2 - 3x + 7 - \frac{11}{2x + 3}$
- 3) $2x^2 + 2.5x + 5 + \frac{15}{2x + 3}$
- 4) $2x^2 - 2.5x - 5 - \frac{20}{2x + 3}$

648 Solve the system of equations shown below algebraically.

$$(x - 3)^2 + (y + 2)^2 = 16$$

$$2x + 2y = 10$$

649 Write $(5 + 2yi)(4 - 3i) - (5 - 2yi)(4 - 3i)$ in $a + bi$ form, where y is a real number.

650 Using the formula below, determine the monthly payment on a 5-year car loan with a monthly percentage rate of 0.625% for a car with an original cost of \$21,000 and a \$1000 down payment, to the nearest cent.

$$P_n = PMT \left(\frac{1 - (1 + i)^{-n}}{i} \right)$$

P_n = present amount borrowed

n = number of monthly pay periods

PMT = monthly payment


i = interest rate per month

The affordable monthly payment is \$300 for the same time period. Determine an appropriate down payment, to the nearest dollar.

651 When $g(x) = \frac{2}{x+2}$ and $h(x) = \log(x+1) + 3$ are graphed on the same set of axes, which coordinates best approximate their point of intersection?

- 1) $(-0.9, 1.8)$
- 2) $(-0.9, 1.9)$
- 3) $(1.4, 3.3)$
- 4) $(1.4, 3.4)$

652 Fifty-five students attending the prom were randomly selected to participate in a survey about the music choice at the prom. Sixty percent responded that a DJ would be preferred over a band. Members of the prom committee thought that the vote would have 50% for the DJ and 50% for the band. A simulation was run 200 times, each of sample size 55, based on the premise that 60% of the students would prefer a DJ. The approximate normal simulation results are shown below.



Using the results of the simulation, determine a plausible interval containing the middle 95% of the data. Round all values to the nearest hundredth. Members of the prom committee are concerned that a vote of all students attending the prom may produce a 50% – 50% split. Explain what statistical evidence supports this concern.


653 The solution to the equation $4x^2 + 98 = 0$ is

- 1) ± 7
- 2) $\pm 7i$
- 3) $\pm \frac{7\sqrt{2}}{2}$
- 4) $\pm \frac{7i\sqrt{2}}{2}$


654 What are the zeros of $P(m) = (m^2 - 4)(m^2 + 1)$?

- 1) 2 and -2, only
- 2) 2, -2, and -4
- 3) -4, i , and $-i$
- 4) 2, -2, i , and $-i$


- 655 Which diagram represents an angle, α , measuring $\frac{13\pi}{20}$ radians drawn in standard position, and its reference angle, θ ?




1)



2)



3)



4)

- 656 What is the solution, if any, of the equation

$$\frac{2}{x+3} - \frac{3}{4-x} = \frac{2x-2}{x^2-x-12}?$$

- 1) -1
- 2) -5
- 3) all real numbers
- 4) no real solution

- 657 A recursive formula for the sequence 18, 9, 4.5, ... is

1) $g_1 = 18$

$$g_n = \frac{1}{2} g_{n-1}$$

2) $g_n = 18 \left(\frac{1}{2} \right)^{n-1}$

3) $g_1 = 18$

$$g_n = 2g_{n-1}$$

4) $g_n = 18(2)^{n-1}$

- 658 Solve for x : $\frac{1}{x} - \frac{1}{3} = -\frac{1}{3x}$

- 659 A radioactive substance has a mass of 140 g at 3 p.m. and 100 g at 8 p.m. Write an equation in the

form $A = A_0 \left(\frac{1}{2} \right)^{\frac{t}{h}}$ that models this situation,

where h is the constant representing the number of hours in the half-life, A_0 is the initial mass, and A is the mass t hours after 3 p.m. Using this equation, solve for h , to the *nearest ten thousandth*. Determine when the mass of the radioactive substance will be 40 g. Round your answer to the *nearest tenth of an hour*.

- 660 The solution set for the equation

$$\sqrt{x+14} - \sqrt{2x+5} = 1$$
 is

1) $\{-6\}$

2) $\{2\}$

3) $\{18\}$

4) $\{2, 22\}$

- 661 The lifespan of a 60-watt lightbulb produced by a company is normally distributed with a mean of 1450 hours and a standard deviation of 8.5 hours. If a 60-watt lightbulb produced by this company is selected at random, what is the probability that its lifespan will be between 1440 and 1465 hours?

1) 0.3803

2) 0.4612

3) 0.8415

4) 0.9612

- 662 The population of Jamesburg for the years 2010-2013, respectively, was reported as follows:
250,000 250,937 251,878 252,822

How can this sequence be recursively modeled?

1) $j_n = 250,000(1.00375)^{n-1}$

2) $j_n = 250,000 + 937^{(n-1)}$

3) $j_1 = 250,000$

$$j_n = 1.00375j_{n-1}$$

4) $j_1 = 250,000$

$$j_n = j_{n-1} + 937$$

- 663 When $b > 0$ and d is a positive integer, the

expression $(3b)^{\frac{2}{d}}$ is equivalent to

1) $\frac{1}{\left(\sqrt[d]{3b}\right)^2}$

2) $\left(\sqrt{3b}\right)^d$





3) $\frac{1}{\sqrt{3b^d}}$

4) $\left(\sqrt[d]{3b}\right)^2$

- 664 Titanium-44 is a radioactive isotope such that every 63 years, its mass decreases by half. For a sample of titanium-44 with an initial mass of 100 grams, write a function that will give the mass of the sample remaining after any amount of time. Define all variables. Scientists sometimes use the average yearly decrease in mass for estimation purposes. Use the average yearly decrease in mass of the sample between year 0 and year 10 to predict the amount of the sample remaining after 40 years. Round your answer to the *nearest tenth*. Is the actual mass of the sample or the estimated mass greater after 40 years? Justify your answer.

- 665 Describe how a controlled experiment can be created to examine the effect of ingredient X in a toothpaste.

- 666 Which graph represents a cosine function with no horizontal shift, an amplitude of 2, and a period of $\frac{2\pi}{3}$?



- 667 Kristin wants to increase her running endurance. According to experts, a gradual mileage increase of 10% per week can reduce the risk of injury. If Kristin runs 8 miles in week one, which expression can help her find the total number of miles she will have run over the course of her 6-week training program?

1) $\sum_{n=1}^6 8(1.10)^{n-1}$

2) $\sum_{n=1}^6 8(1.10)^n$


3) $\frac{8 - 8(1.10)^6}{0.90}$

4) $\frac{8 - 8(0.10)^n}{1.10}$

- 668 A study was designed to test the effectiveness of a new drug. Half of the volunteers received the drug. The other half received a sugar pill. The probability of a volunteer receiving the drug and getting well was 40%. What is the probability of a volunteer getting well, given that the volunteer received the drug?

669 Solve for all values of p : $\frac{3p}{p-5} - \frac{2}{p+3} = \frac{p}{p+3}$

670 Using the unit circle below, explain why $\csc \theta = \frac{1}{y}$.



- 671 The equation $t = \frac{1}{0.0105} \ln\left(\frac{A}{5000}\right)$ relates time, t , in years, to the amount of money, A , earned by a \$5000 investment. Which statement accurately describes the relationship between the average rates of change of t on the intervals $[6000, 8000]$ and $[9000, 12,000]$?
- 1) A comparison cannot be made because the intervals are different sizes.
 - 2) The average rate of change is equal for both intervals.
 - 3) The average rate of change is larger for the interval $[6000, 8000]$.
 - 4) The average rate of change is larger for the interval $[9000, 12,000]$.


672 Given the equal terms $\sqrt[3]{x^5}$ and $y^{\frac{5}{6}}$, determine and state y , in terms of x .

- 673 What is the inverse of the function $y = \log_3 x$?
- 1) $y = x^3$
 - 2) $y = \log_x 3$
 - 3) $y = 3^x$
 - 4) $x = 3^y$

674 If $\sin^2(32^\circ) + \cos^2(M) = 1$, then M equals

- 1) 32°
- 2) 58°
- 3) 68°
- 4) 72°

675 Graph $y = 400(.85)^{2x} - 6$ on the set of axes below.



676 A payday loan company makes loans between \$100 and \$1000 available to customers. Every 14 days, customers are charged 30% interest with compounding. In 2013, Remi took out a \$300 payday loan. Which expression can be used to calculate the amount she would owe, in dollars, after one year if she did not make payments?

- 1) $300(.30)^{\frac{14}{365}}$
- 2) $300(1.30)^{\frac{14}{365}}$
- 3) $300(.30)^{\frac{365}{14}}$
- 4) $300(1.30)^{\frac{365}{14}}$

677 Simplify $xi(i - 7i)^2$, where i is the imaginary unit.

678 Which function shown below has a greater average rate of change on the interval $[-2, 4]$? Justify your answer.

x	$f(x)$
-4	0.3125
-3	0.625
-2	1.25
-1	2.5
0	5
1	10
2	20
3	40
4	80
5	160
6	320

$$g(x) = 4x^3 - 5x^2 + 3$$


679 A public opinion poll was conducted on behalf of Mayor Ortega's reelection campaign shortly before the election. 264 out of 550 likely voters said they would vote for Mayor Ortega; the rest said they would vote for his opponent. Which statement is *least* appropriate to make, according to the results of the poll?

- 1) There is a 48% chance that Mayor Ortega will win the election.
- 2) The point estimate (\hat{p}) of voters who will vote for Mayor Ortega is 48%.
- 3) It is most likely that between 44% and 52% of voters will vote for Mayor Ortega.
- 4) Due to the margin of error, an inference cannot be made regarding whether Mayor Ortega or his opponent is most likely to win the election.

680 Which equation represents a parabola with a focus of $(0, 4)$ and a directrix of $y = 2$?

- 1) $y = x^2 + 3$
- 2) $y = -x^2 + 1$
- 3) $y = \frac{x^2}{2} + 3$
- 4) $y = \frac{x^2}{4} + 3$

681 Anne has a coin. She does not know if it is a fair coin. She flipped the coin 100 times and obtained 73 heads and 27 tails. She ran a computer simulation of 200 samples of 100 fair coin flips. The output of the proportion of heads is shown below.




Given the results of her coin flips and of her computer simulation, which statement is most accurate?

- 1) 73 of the computer's next 100 coin flips will be heads.
- 2) 50 of her next 100 coin flips will be heads.
- 3) Her coin is not fair.
- 4) Her coin is fair.

682 Using the identity $\sin^2 \theta + \cos^2 \theta = 1$, find the value of $\tan \theta$, to the *nearest hundredth*, if $\cos \theta$ is -0.7 and θ is in Quadrant II.

- 683 An orange-juice processing plant receives a truckload of oranges. The quality control team randomly chooses three pails of oranges, each containing 50 oranges, from the truckload. Identify the sample and the population in the given scenario. State *one* conclusion that the quality control team could make about the population if 5% of the sample was found to be unsatisfactory.

- 684 Find algebraically the zeros for $p(x) = x^3 + x^2 - 4x - 4$. On the set of axes below, graph $y = p(x)$.



- 685 Algebraically determine the values of x that satisfy the system of equations below.

$$y = -2x + 1$$

$$y = -2x^2 + 3x + 1$$

- 686 Which statement is *incorrect* for the graph of the function $y = -3 \cos \left[\frac{\pi}{3} (x - 4) \right] + 7$?

- 1) The period is 6.
- 2) The amplitude is 3.
- 3) The range is $[4, 10]$.
- 4) The midline is $y = -4$.

- 687 For $x \neq 0$, which expressions are equivalent to one divided by the sixth root of x ?

I. $\frac{\sqrt[6]{x}}{\sqrt[3]{x}}$ II. $\frac{x^{\frac{1}{6}}}{x^{\frac{1}{3}}}$ III. $x^{\frac{-1}{6}}$

- 1) I and II, only
- 2) I and III, only
- 3) II and III, only
- 4) I, II, and III

- 688 Which function represents exponential decay?

- 1) $y = 2^{0.3t}$
- 2) $y = 1.2^{3t}$
- 3) $y = \left(\frac{1}{2} \right)^{-t}$
- 4) $y = 5^{-t}$

- 689 After sitting out of the refrigerator for a while, a turkey at room temperature (68°F) is placed into an oven at 8 a.m., when the oven temperature is 325°F . Newton's Law of Heating explains that the temperature of the turkey will increase proportionally to the difference between the temperature of the turkey and the temperature of the oven, as given by the formula below:

$$T = T_a + (T_0 - T_a)e^{-kt}$$

T_a = the temperature surrounding the object

T_0 = the initial temperature of the object


t = the time in hours

T = the temperature of the object after t hours

k = decay constant


The turkey reaches the temperature of approximately 100°F after 2 hours. Find the value of k , to the *nearest thousandth*, and write an equation to determine the temperature of the turkey after t hours. Determine the Fahrenheit temperature of the turkey, to the *nearest degree*, at 3 p.m.

- 690 What is the solution to the system of equations $y = 3x - 2$ and $y = g(x)$ where $g(x)$ is defined by the function below?




- 1) $\{(0, -2)\}$
 - 2) $\{(0, -2), (1, 6)\}$
 - 3) $\{(1, 6)\}$
 - 4) $\{(1, 1), (6, 16)\}$
- 691 Based on climate data that have been collected in Bar Harbor, Maine, the average monthly temperature, in degrees F, can be modeled by the equation $B(x) = 23.914\sin(0.508x - 2.116) + 55.300$. The same governmental agency collected average monthly temperature data for Phoenix, Arizona, and found the temperatures could be modeled by the equation $P(x) = 20.238\sin(0.525x - 2.148) + 86.729$. Which statement can *not* be concluded based on the average monthly temperature models x months after starting data collection?
- 1) The average monthly temperature variation is more in Bar Harbor than in Phoenix.
 - 2) The midline average monthly temperature for Bar Harbor is lower than the midline temperature for Phoenix.
 - 3) The maximum average monthly temperature for Bar Harbor is 79°F , to the nearest degree.
 - 4) The minimum average monthly temperature for Phoenix is 20°F , to the nearest degree.

- 692 A game spinner is divided into 6 equally sized regions, as shown in the diagram below.



For Miles to win, the spinner must land on the number 6. After spinning the spinner 10 times, and losing all 10 times, Miles complained that the spinner is unfair. At home, his dad ran 100 simulations of spinning the spinner 10 times, assuming the probability of winning each spin is $\frac{1}{6}$.

The output of the simulation is shown in the diagram below.




Which explanation is appropriate for Miles and his dad to make?

- 1) The spinner was likely unfair, since the number 6 failed to occur in about 20% of the simulations.
- 2) The spinner was likely unfair, since the spinner should have landed on the number 6 by the sixth spin.
- 3) The spinner was likely not unfair, since the number 6 failed to occur in about 20% of the simulations.
- 4) The spinner was likely not unfair, since in the output the player wins once or twice in the majority of the simulations.

- 693 Seventy-two students are randomly divided into two equally-sized study groups. Each member of the first group (group 1) is to meet with a tutor after school twice each week for one hour. The second group (group 2), is given an online subscription to a tutorial account that they can access for a maximum of two hours each week. Students in both groups are given the same tests during the year. A summary of the two groups' final grades is shown below:

	Group 1	Group 2
\bar{x}	80.16	83.8
S_x	6.9	5.2

Calculate the mean difference in the final grades (group 1 – group 2) and explain its meaning in the context of the problem. A simulation was conducted in which the students' final grades were rerandomized 500 times. The results are shown below.



Use the simulation to determine if there is a significant difference in the final grades. Explain your answer.

- 694 A circle centered at the origin has a radius of 10 units. The terminal side of an angle, θ , intercepts the circle in Quadrant II at point C . The y-coordinate of point C is 8. What is the value of $\cos \theta$?
- 1) $-\frac{3}{5}$
 - 2) $-\frac{3}{4}$
 - 3) $\frac{3}{5}$
 - 4) $\frac{4}{5}$
- 695 While experimenting with her calculator, Candy creates the sequence 4, 9, 19, 39, 79, Write a recursive formula for Candy's sequence. Determine the eighth term in Candy's sequence.
- 696 A rabbit population doubles every 4 weeks. There are currently five rabbits in a restricted area. If t represents the time, in weeks, and $P(t)$ is the population of rabbits with respect to time, about how many rabbits will there be in 98 days?
- 1) 56
 - 2) 152
 - 3) 3688
 - 4) 81,920
- 697 A suburban high school has a population of 1376 students. The number of students who participate in sports is 649. The number of students who participate in music is 433. If the probability that a student participates in either sports or music is $\frac{974}{1376}$, what is the probability that a student participates in both sports and music?

- 698 Given events A and B , such that $P(A) = 0.6$, $P(B) = 0.5$, and $P(A \cup B) = 0.8$, determine whether A and B are independent or dependent.
- 699 Sally's high school is planning their spring musical. The revenue, R , generated can be determined by the function $R(t) = -33t^2 + 360t$, where t represents the price of a ticket. The production cost, C , of the musical is represented by the function $C(t) = 700 + 5t$. What is the highest ticket price, to the *nearest dollar*, they can charge in order to *not* lose money on the event?
- $t = 3$
 - $t = 5$
 - $t = 8$
 - $t = 11$
- 700 Alexa earns \$33,000 in her first year of teaching and earns a 4% increase in each successive year. Write a geometric series formula, S_n , for Alexa's total earnings over n years. Use this formula to find Alexa's total earnings for her first 15 years of teaching, to the *nearest cent*.
- 701 Cheap and Fast gas station is conducting a consumer satisfaction survey. Which method of collecting data would most likely lead to a biased sample?
- interviewing every 5th customer to come into the station
 - interviewing customers chosen at random by a computer at the checkout
 - interviewing customers who call an 800 number posted on the customers' receipts
 - interviewing every customer who comes into the station on a day of the week chosen at random out of a hat
- 702 The solution to the equation $18x^2 - 24x + 87 = 0$ is
- $-\frac{2}{3} \pm 6i\sqrt{158}$
 - $-\frac{2}{3} \pm \frac{1}{6}i\sqrt{158}$
 - $\frac{2}{3} \pm 6i\sqrt{158}$
 - $\frac{2}{3} \pm \frac{1}{6}i\sqrt{158}$
- 703 Mallory wants to buy a new window air conditioning unit. The cost for the unit is \$329.99. If she plans to run the unit three months out of the year for an annual operating cost of \$108.78, which function models the cost per year over the lifetime of the unit, $C(n)$, in terms of the number of years, n , that she owns the air conditioner.
- $C(n) = 329.99 + 108.78n$
 - $C(n) = 329.99 + 326.34n$
 - $C(n) = \frac{329.99 + 108.78n}{n}$
 - $C(n) = \frac{329.99 + 326.34n}{n}$
- 704 Jim is looking to buy a vacation home for \$172,600 near his favorite southern beach. The formula to compute a mortgage payment, M , is
- $$M = P \bullet \frac{r(1+r)^N}{(1+r)^N - 1}$$
- where P is the principal amount of the loan, r is the monthly interest rate, and N is the number of monthly payments. Jim's bank offers a monthly interest rate of 0.305% for a 15-year mortgage. With no down payment, determine Jim's mortgage payment, rounded to the *nearest dollar*. Algebraically determine and state the down payment, rounded to the *nearest dollar*, that Jim needs to make in order for his mortgage payment to be \$1100.
- 705 Over the set of integers, factor the expression $4x^3 - x^2 + 16x - 4$ completely.
- 706 The function $f(x) = 2^{-0.25x} \bullet \sin\left(\frac{\pi}{2}x\right)$ represents a damped sound wave function. What is the average rate of change for this function on the interval $[-7, 7]$, to the *nearest hundredth*?
- 3.66
 - 0.30
 - 0.26
 - 3.36

- 707 The set of data in the table below shows the results of a survey on the number of messages that people of different ages text on their cell phones each month.

Text Messages per Month			
Age Group	0-10	11-50	Over 50
15-18	4	37	68
19-22	6	25	87
23-60	25	47	157

If a person from this survey is selected at random, what is the probability that the person texts over 50 messages per month given that the person is between the ages of 23 and 60?

- 1) $\frac{157}{229}$
- 2) $\frac{157}{312}$
- 3) $\frac{157}{384}$
- 4) $\frac{157}{456}$

- 708 What does $\left(\frac{-54x^9}{y^4}\right)^{\frac{2}{3}}$ equal?

- $$\begin{aligned} 1) \quad & \frac{9ix^6\sqrt[3]{4}}{y^3\sqrt[3]{y^2}} \\ 2) \quad & \frac{9ix^6\sqrt[3]{4}}{y^2\sqrt[3]{y^2}} \\ 3) \quad & \frac{9x^6\sqrt[3]{4}}{y^3\sqrt[3]{y}} \\ 4) \quad & \frac{9x^6\sqrt[3]{4}}{y^2\sqrt[3]{y^2}} \end{aligned}$$

- 709 Which binomial is *not* a factor of the expression $x^3 - 11x^2 + 16x + 84$?

- 1) $x + 2$
- 2) $x + 4$
- 3) $x - 6$
- 4) $x - 7$

- 710 The weight of a bag of pears at the local market averages 8 pounds with a standard deviation of 0.5 pound. The weights of all the bags of pears at the market closely follow a normal distribution. Determine what percentage of bags, to the *nearest integer*, weighed *less* than 8.25 pounds.

- 711 The completely factored form of

$2d^4 + 6d^3 - 18d^2 - 54d$ is

- 1) $2d(d^2 - 9)(d + 3)$
- 2) $2d(d^2 + 9)(d + 3)$
- 3) $2d(d + 3)^2(d - 3)$
- 4) $2d(d - 3)^2(d + 3)$

- 712 When factored completely, $m^5 + m^3 - 6m$ is
equivalent to

- 1) $(m+3)(m-2)$
- 2) $(m^2+3m)(m^2-2)$
- 3) $m(m^4+m^2-6)$
- 4) $m(m^2+3)(m^2-2)$

- 713 Which statement about statistical analysis is *false*?

- 1) Experiments can suggest patterns and relationships in data.
- 2) Experiments can determine cause and effect relationships.
- 3) Observational studies can determine cause and effect relationships.
- 4) Observational studies can suggest patterns and relationships in data.

- 714 Given $z(x) = 6x^3 + bx^2 - 52x + 15$, $z(2) = 35$, and $z(-5) = 0$, algebraically determine all the zeros of $z(x)$.

715 Given $f^{-1}(x) = -\frac{3}{4}x + 2$, which equation represents $f(x)$?

- 1) $f(x) = \frac{4}{3}x - \frac{8}{3}$
- 2) $f(x) = -\frac{4}{3}x + \frac{8}{3}$
- 3) $f(x) = \frac{3}{4}x - 2$
- 4) $f(x) = -\frac{3}{4}x + 2$

716 Monthly mortgage payments can be found using the formula below:

$$M = \frac{P\left(\frac{r}{12}\right)\left(1 + \frac{r}{12}\right)^n}{\left(1 + \frac{r}{12}\right)^n - 1}$$

M = monthly payment
 P = amount borrowed
 r = annual interest rate
 n = number of monthly payments


The Banks family would like to borrow \$120,000 to purchase a home. They qualified for an annual interest rate of 4.8%. Algebraically determine the *fewest* number of whole years the Banks family would need to include in the mortgage agreement in order to have a monthly payment of no more than \$720.

717 Last year, the total revenue for Home Style, a national restaurant chain, increased 5.25% over the previous year. If this trend were to continue, which expression could the company's chief financial officer use to approximate their monthly percent increase in revenue? [Let m represent months.]

- 1) $(1.0525)^m$
- 2) $(1.0525)^{\frac{12}{m}}$
- 3) $(1.00427)^m$
- 4) $(1.00427)^{\frac{m}{12}}$

718 The value of a certain small passenger car based on its use in years is modeled by

$V(t) = 28482.698(0.684)^t$, where $V(t)$ is the value in dollars and t is the time in years. Zach had to take out a loan to purchase the small passenger car. The function $Z(t) = 22151.327(0.778)^t$, where $Z(t)$ is measured in dollars, and t is the time in years, models the unpaid amount of Zach's loan over time. Graph $V(t)$ and $Z(t)$ over the interval $0 \leq t \leq 5$, on the set of axes below.



State when $V(t) = Z(t)$, to the *nearest hundredth*, and interpret its meaning in the context of the problem. Zach takes out an insurance policy that requires him to pay a \$3000 deductible in case of a collision. Zach will cancel the collision policy when the value of his car equals his deductible. To the *nearest year*, how long will it take Zach to cancel this policy? Justify your answer.

719 One of the medical uses of Iodine-131 (I-131), a radioactive isotope of iodine, is to enhance x-ray images. The half-life of I-131 is approximately 8.02 days. A patient is injected with 20 milligrams of I-131. Determine, to the *nearest day*, the amount of time needed before the amount of I-131 in the patient's body is approximately 7 milligrams.

- 720 The Ferris wheel at the landmark Navy Pier in Chicago takes 7 minutes to make one full rotation. The height, H , in feet, above the ground of one of the six-person cars can be modeled by

$$H(t) = 70 \sin\left(\frac{2\pi}{7}(t - 1.75)\right) + 80, \text{ where } t \text{ is time,}$$

in minutes. Using $H(t)$ for one full rotation, this car's minimum height, in feet, is

- 1) 150
 - 2) 70
 - 3) 10
 - 4) 0
- 721 Elizabeth waited for 6 minutes at the drive thru at her favorite fast-food restaurant the last time she visited. She was upset about having to wait that long and notified the manager. The manager assured her that her experience was very unusual and that it would not happen again. A study of customers commissioned by this restaurant found an approximately normal distribution of results. The mean wait time was 226 seconds and the standard deviation was 38 seconds. Given these data, and using a 95% level of confidence, was Elizabeth's wait time unusual? Justify your answer.

- 722 The probability that Gary and Jane have a child with blue eyes is 0.25, and the probability that they have a child with blond hair is 0.5. The probability that they have a child with both blue eyes and blond hair is 0.125. Given this information, the events blue eyes and blond hair are

I: dependent
 II: independent
 III: mutually exclusive

- 1) I, only
 - 2) II, only
 - 3) I and III
 - 4) II and III
- 723 Determine if $x - 5$ is a factor of $2x^3 - 4x^2 - 7x - 10$. Explain your answer.

- 724 Solve algebraically for all values of x :
 $\sqrt{x-4} + x = 6$


- 725 Which statement about the graph of $c(x) = \log_6 x$ is false?

- 1) The asymptote has equation $y = 0$.
- 2) The graph has no y -intercept.
- 3) The domain is the set of positive reals.
- 4) The range is the set of all real numbers.

- 726 If the terminal side of angle θ , in standard position, passes through point $(-4, 3)$, what is the numerical value of $\sin \theta$?

- 1) $\frac{3}{5}$
- 2) $\frac{4}{5}$
- 3) $-\frac{3}{5}$
- 4) $-\frac{4}{5}$

- 727 On the grid below, sketch a cubic polynomial whose zeros are 1, 3, and -2.



- 728 The zeros for $f(x) = x^4 - 4x^3 - 9x^2 + 36x$ are

- 1) $\{0, \pm 3, 4\}$
- 2) $\{0, 3, 4\}$
- 3) $\{0, \pm 3, -4\}$
- 4) $\{0, 3, -4\}$


729 The results of a poll of 200 students are shown in the table below:

	Preferred Music Style		
	Techno	Rap	Country
Female	54	25	27
Male	36	40	18


For this group of students, do these data suggest that gender and preferred music styles are independent of each other? Justify your answer.

730 Which graph has the following characteristics?


- three real zeros
- as $x \rightarrow -\infty$, $f(x) \rightarrow -\infty$
- as $x \rightarrow \infty$, $f(x) \rightarrow \infty$




1)



2)



3)



4)

731 Algebraically determine whether the function

$$j(x) = x^4 - 3x^2 - 4$$

is odd, even, or neither.

732 Algebraically prove that $\frac{x^3 + 9}{x^3 + 8} = 1 + \frac{1}{x^3 + 8}$,
 where $x \neq -2$.

733 Mr. Farison gave his class the three mathematical rules shown below to either prove or disprove.
 Which rules can be proved for all real numbers?

I $(m+p)^2 = m^2 + 2mp + p^2$

II $(x+y)^3 = x^3 + 3xy + y^3$

III $(a^2 + b^2)^2 = (a^2 - b^2)^2 + (2ab)^2$

- 1) I, only
- 2) I and II
- 3) II and III
- 4) I and III

734 Express $(1 - i)^3$ in $a + bi$ form.


735 Given the parent function $p(x) = \cos x$, which phrase best describes the transformation used to obtain the graph of $g(x) = \cos(x + a) - b$, if a and b are positive constants?

- 1) right a units, up b units
- 2) right a units, down b units
- 3) left a units, up b units
- 4) left a units, down b units

- 736 Ayva designed an experiment to determine the effect of a new energy drink on a group of 20 volunteer students. Ten students were randomly selected to form group 1 while the remaining 10 made up group 2. Each student in group 1 drank one energy drink, and each student in group 2 drank one cola drink. Ten minutes later, their times were recorded for reading the same paragraph of a novel. The results of the experiment are shown below.

Group 1 (seconds)	Group 2 (seconds)
17.4	23.3
18.1	18.8
18.2	22.1
19.6	12.7
18.6	16.9
16.2	24.4
16.1	21.2
15.3	21.2
17.8	16.3
19.7	14.5
Mean = 17.7	Mean = 19.1

Ayva thinks drinking energy drinks makes students read faster. Using information from the experimental design or the results, explain why Ayva's hypothesis may be *incorrect*. Using the given results, Ayva randomly mixes the 20 reading times, splits them into two groups of 10, and simulates the difference of the means 232 times.



Ayva has decided that the difference in mean reading times is not an unusual occurrence. Support her decision using the results of the simulation. Explain your reasoning.

- 737 Joelle has a credit card that has a 19.2% annual interest rate compounded monthly. She owes a total balance of B dollars after m months. Assuming she makes no payments on her account, the table below illustrates the balance she owes after m months.

m	B
0	100.00
10	1172.00
19	1352.00
36	1770.80
60	2591.90
69	2990.00
72	3135.80
73	3186.00

Over which interval of time is her average rate of change for the balance on her credit card account the greatest?

- 1) month 10 to month 60 3) month 36 to month 72
 2) month 19 to month 69 4) month 60 to month 73
- 738 Data collected about jogging from students with two older siblings are shown in the table below.

	Neither Sibling Jogs	One Sibling Jogs	Both Siblings Jog
Student Does Not Jog	1168	1823	1380
Student Jogs	188	416	400

Using these data, determine whether a student with two older siblings is more likely to jog if one sibling jogs or if both siblings jog. Justify your answer.

- 739 The formula below can be used to model which scenario?

$$a_1 = 3000$$

$$a_n = 0.80a_{n-1}$$

- 1) The first row of a stadium has 3000 seats, and each row thereafter has 80 more seats than the row in front of it.
 2) The last row of a stadium has 3000 seats, and each row before it has 80 fewer seats than the row behind it.
 3) A bank account starts with a deposit of \$3000, and each year it grows by 80%.
 4) The initial value of a specialty toy is \$3000, and its value each of the following years is 20% less.

- 740 Solve the following system of equations algebraically for all values of x , y , and z :

$$x + 3y + 5z = 45$$

$$6x - 3y + 2z = -10$$

$$-2x + 3y + 8z = 72$$

- 741 The heights of women in the United States are normally distributed with a mean of 64 inches and a standard deviation of 2.75 inches. The percent of women whose heights are between 64 and 69.5 inches, to the *nearest whole percent*, is


- 1) 6
 2) 48
 3) 68
 4) 95

- 742 The distance needed to stop a car after applying the brakes varies directly with the square of the car's speed. The table below shows stopping distances for various speeds.

Speed (mph)	10	20	30	40	50	60	70
Distance (ft)	6.25	25	56.25	100	156.25	225	306.25

Determine the average rate of change in braking distance, in ft/mph, between one car traveling at 50 mph and one traveling at 70 mph. Explain what this rate of change means as it relates to braking distance.

- 743 a) On the axes below, sketch *at least one* cycle of a sine curve with an amplitude of 2, a midline at $y = -\frac{3}{2}$, and a period of 2π .



- b) Explain any differences between a sketch of $y = 2\sin\left(x - \frac{\pi}{3}\right) - \frac{3}{2}$ and the sketch from part a.


- 744 The expression $6xi^3(-4xi + 5)$ is equivalent to
- 1) $2x - 5i$
 - 2) $-24x^2 - 30xi$
 - 3) $-24x^2 + 30x - i$
 - 4) $26x - 24x^2i - 5i$

- 745 When $g(x)$ is divided by $x + 4$, the remainder is 0.

Given $g(x) = x^4 + 3x^3 - 6x^2 - 6x + 8$, which conclusion about $g(x)$ is true?

- 1) $g(4) = 0$
- 2) $g(-4) = 0$
- 3) $x - 4$ is a factor of $g(x)$.
- 4) No conclusion can be made regarding $g(x)$.





- 746 Graph $y = \log_2(x + 3) - 5$ on the set of axes below. Use an appropriate scale to include *both* intercepts.



Describe the behavior of the given function as x approaches -3 and as x approaches positive infinity.

- 747 The directrix of the parabola $12(y + 3) = (x - 4)^2$ has the equation $y = -6$. Find the coordinates of the focus of the parabola.





748 Which diagram shows an angle rotation of 1 radian on the unit circle?



749 A ball is dropped from a height of 32 feet. It bounces and rebounds 80% of the height from which it was falling. What is the total downward distance, in feet, the ball traveled up to the 12th bounce?

- 1) 29
- 2) 58
- 3) 120
- 4) 149

750 If a , b , and c are all positive real numbers, which graph could represent the sketch of the graph of $p(x) = -a(x+b)(x^2 - 2cx + c^2)$?



751 A solution of the equation $2x^2 + 3x + 2 = 0$ is

- 1) $-\frac{3}{4} + \frac{1}{4}i\sqrt{7}$
- 2) $-\frac{3}{4} + \frac{1}{4}i$
- 3) $-\frac{3}{4} + \frac{1}{4}\sqrt{7}$
- 4) $\frac{1}{2}$

752 Which statement(s) about statistical studies is true?

- I. A survey of all English classes in a high school would be a good sample to determine the number of hours students throughout the school spend studying.
- II. A survey of all ninth graders in a high school would be a good sample to determine the number of student parking spaces needed at that high school.
- III. A survey of all students in one lunch period in a high school would be a good sample to determine the number of hours adults spend on social media websites.
- IV. A survey of all Calculus students in a high school would be a good sample to determine the number of students throughout the school who don't like math.

- 1) I, only
- 2) II, only
- 3) I and III
- 4) III and IV

753 Use an appropriate procedure to show that $x - 4$ is a factor of the function $f(x) = 2x^3 - 5x^2 - 11x - 4$. Explain your answer.

754 Use the properties of rational exponents to determine the value of y for the equation:

$$\frac{\sqrt[3]{x^8}}{(x^4)^{\frac{1}{3}}} = x^y, x > 1$$

755 Which expression is equivalent to $\frac{4x^3 + 9x - 5}{2x - 1}$,

where $x \neq \frac{1}{2}$?

- 1) $2x^2 + x + 5$
- 2) $2x^2 + \frac{11}{2} + \frac{1}{2(2x - 1)}$
- 3) $2x^2 - x + 5$
- 4) $2x^2 - x + 4 + \frac{1}{2x - 1}$

756 Sean's team has a baseball game tomorrow. He pitches 50% of the games. There is a 40% chance of rain during the game tomorrow. If the probability that it rains given that Sean pitches is 40%, it can be concluded that these two events are

- 1) independent
- 2) dependent
- 3) mutually exclusive
- 4) complements

757 Solve the following system of equations algebraically for all values of x , y , and z :

$$x + y + z = 1$$

$$2x + 4y + 6z = 2$$

$$-x + 3y - 5z = 11$$

758 The expression $\left(\frac{m^2}{m^{\frac{1}{3}}}\right)^{-\frac{1}{2}}$ is equivalent to

- 1) $-\sqrt[6]{m^5}$
- 2) $\frac{1}{\sqrt[6]{m^5}}$
- 3) $-m^5\sqrt{m}$
- 4) $\frac{1}{m^5\sqrt{m}}$

759 Consider the system shown below.


$$2x - y = 4$$

$$(x + 3)^2 + y^2 = 8$$

The two solutions of the system can be described as

- 1) both imaginary
- 2) both irrational
- 3) both rational
- 4) one rational and one irrational

- 760 A candidate for political office commissioned a poll. His staff received responses from 900 likely voters and 55% of them said they would vote for the candidate. The staff then conducted a simulation of 1000 more polls of 900 voters, assuming that 55% of voters would vote for their candidate. The output of the simulation is shown in the diagram below.




Given this output, and assuming a 95% confidence level, the margin of error for the poll is closest to

- 1) 0.01
 - 2) 0.03
 - 3) 0.06
 - 4) 0.12
- 761 Two versions of a standardized test are given, an April version and a May version. The statistics for the April version show a mean score of 480 and a standard deviation of 24. The statistics for the May version show a mean score of 510 and a standard deviation of 20. Assume the scores are normally distributed. Joanne took the April version and scored in the interval 510-540. What is the probability, to the nearest ten thousandth, that a test paper selected at random from the April version scored in the same interval? Maria took the May version. In what interval must Maria score to claim she scored as well as Joanne?

- 762 Functions f , g , and h are given below.

$$f(x) = \sin(2x)$$

$$g(x) = f(x) + 1$$



Which statement is true about functions f , g , and h ?

- 1) $f(x)$ and $g(x)$ are odd, $h(x)$ is even.
 - 2) $f(x)$ and $g(x)$ are even, $h(x)$ is odd.
 - 3) $f(x)$ is odd, $g(x)$ is neither, $h(x)$ is even.
 - 4) $f(x)$ is even, $g(x)$ is neither, $h(x)$ is odd.
- 763 The function $p(t) = 110e^{0.03922t}$ models the population of a city, in millions, t years after 2010. As of today, consider the following two statements:
- I. The current population is 110 million.
 - II. The population increases continuously by approximately 3.9% per year.
- This model supports
- 1) I, only
 - 2) II, only
 - 3) both I and II
 - 4) neither I nor II

- 764 To the *nearest tenth*, the value of x that satisfies $2^x = -2x + 11$ is
 1) 2.5
 2) 2.6
 3) 5.8
 4) 5.9
- 765 The voltage used by most households can be modeled by a sine function. The maximum voltage is 120 volts, and there are 60 cycles *every second*. Which equation best represents the value of the voltage as it flows through the electric wires, where t is time in seconds?
 1) $V = 120\sin(t)$
 2) $V = 120\sin(60t)$
 3) $V = 120\sin(60\pi t)$
 4) $V = 120\sin(120\pi t)$
- 766 Which statement regarding the graphs of the functions below is *untrue*?
 $f(x) = 3\sin 2x$, from $-\pi < x < \pi$
 $g(x) = (x - 0.5)(x + 4)(x - 2)$
 $h(x) = \log_2 x$
 $j(x) = -|4x - 2| + 3$
 1) $f(x)$ and $j(x)$ have a maximum y -value of 3.
 2) $f(x)$, $h(x)$, and $j(x)$ have one y -intercept.
 3) $g(x)$ and $j(x)$ have the same end behavior as $x \rightarrow -\infty$.
 4) $g(x)$, $h(x)$, and $j(x)$ have rational zeros.
- 767 A parabola has its focus at $(1, 2)$ and its directrix is $y = -2$. The equation of this parabola could be
 1) $y = 8(x + 1)^2$
 2) $y = \frac{1}{8}(x + 1)^2$
 3) $y = 8(x - 1)^2$
 4) $y = \frac{1}{8}(x - 1)^2$
- 768 The function $f(x) = \frac{x - 3}{x^2 + 2x - 8}$ is undefined when x equals
 1) 2 or -4
 2) 4 or -2
 3) 3, only
 4) 2, only
- 769 Jasmine decides to put \$100 in a savings account each month. The account pays 3% annual interest, compounded monthly. How much money, S , will Jasmine have after one year?
 1) $S = 100(1.03)^{12}$
 2) $S = \frac{100 - 100(1.0025)^{12}}{1 - 1.0025}$
 3) $S = 100(1.0025)^{12}$
 4) $S = \frac{100 - 100(1.03)^{12}}{1 - 1.03}$
- 770 As x increases from 0 to $\frac{\pi}{2}$, the graph of the equation $y = 2\tan x$ will
 1) increase from 0 to 2
 2) decrease from 0 to -2
 3) increase without limit
 4) decrease without limit
- 771 The roots of the equation $x^2 + 2x + 5 = 0$ are
 1) -3 and 1
 2) -1 , only
 3) $-1 + 2i$ and $-1 - 2i$
 4) $-1 + 4i$ and $-1 - 4i$
- 772 The inverse of the function $f(x) = \frac{x + 1}{x - 2}$ is
 1) $f^{-1}(x) = \frac{x + 1}{x + 2}$
 2) $f^{-1}(x) = \frac{2x + 1}{x - 1}$
 3) $f^{-1}(x) = \frac{x + 1}{x - 2}$
 4) $f^{-1}(x) = \frac{x - 1}{x + 1}$

- 773 The loudness of sound is measured in units called decibels (dB). These units are measured by first assigning an intensity I_0 to a very soft sound that is called the threshold sound. The sound to be measured is assigned an intensity, I , and the decibel rating, d , of this sound is found using $d = 10 \log \frac{I}{I_0}$. The threshold sound audible to the average person is $1.0 \times 10^{-12} \text{ W/m}^2$ (watts per square meter). Consider the following sound level classifications:

Moderate	45-69 dB
Loud	70-89 dB
Very loud	90-109 dB
Deafening	>110 dB

How would a sound with intensity $6.3 \times 10^{-3} \text{ W/m}^2$ be classified?

- 1) moderate 3) very loud
2) loud 4) deafening

- 774 A student studying public policy created a model for the population of Detroit, where the population decreased 25% over a decade. He used the model $P = 714(0.75)^d$, where P is the population, in thousands, d decades after 2010. Another student, Suzanne, wants to use a model that would predict the population after y years. Suzanne's model is best represented by

- 1) $P = 714(0.6500)^y$
- 2) $P = 714(0.8500)^y$
- 3) $P = 714(0.9716)^y$
- 4) $P = 714(0.9750)^y$

- 775 If $f(x) = 3|x| - 1$ and $g(x) = 0.03x^3 - x + 1$, an approximate solution for the equation $f(x) = g(x)$ is

- 1) 1.96
- 2) 11.29
- 3) $(-0.99, 1.96)$
- 4) $(11.29, 32.87)$

- 776 Given i is the imaginary unit, $(2 - yi)^2$ in simplest form is

- 1) $y^2 - 4yi + 4$
- 2) $-y^2 - 4yi + 4$
- 3) $-y^2 + 4$
- 4) $y^2 + 4$

- 777 The distribution of the diameters of ball bearings made under a given manufacturing process is normally distributed with a mean of 4 cm and a standard deviation of 0.2 cm. What proportion of the ball bearings will have a diameter less than 3.7 cm?

- 1) 0.0668
- 2) 0.4332
- 3) 0.8664
- 4) 0.9500

- 778 The focal length, F , of a camera's lens is related to the distance of the object from the lens, J , and the distance to the image area in the camera, W , by the formula below.

$$\frac{1}{J} + \frac{1}{W} = \frac{1}{F}$$

When this equation is solved for J in terms of F and W , J equals

- 1) $F - W$
- 2) $\frac{FW}{F - W}$
- 3) $\frac{FW}{W - F}$
- 4) $\frac{1}{F} - \frac{1}{W}$

- 779 A sine function increasing through the origin can be used to model light waves. Violet light has a wavelength of 400 nanometers. Over which interval is the height of the wave *decreasing*, only?

1) (0,200)
 2) (100,300)
 3) (200,400)
 4) (300,400)

- 780 The function $M(t)$ represents the mass of radium over time, t , in years.

$$M(t) = 100e^{\frac{\left(\ln \frac{1}{2}\right)t}{1590}}$$

Determine if the function $M(t)$ represents growth or decay. Explain your reasoning.

- 781 Julie averaged 85 on the first three tests of the semester in her mathematics class. If she scores 93 on each of the remaining tests, her average will be 90. Which equation could be used to determine how many tests, T , are left in the semester?

1) $\frac{255 + 93T}{3T} = 90$
 2) $\frac{255 + 90T}{3T} = 93$
 3) $\frac{255 + 93T}{T + 3} = 90$
 4) $\frac{255 + 90T}{T + 3} = 93$

- 782 The solutions to the equation $-\frac{1}{2}x^2 = -6x + 20$ are

1) $-6 \pm 2i$
 2) $-6 \pm 2\sqrt{19}$
 3) $6 \pm 2i$
 4) $6 \pm 2\sqrt{19}$

- 783 Given: $h(x) = \frac{2}{9}x^3 + \frac{8}{9}x^2 - \frac{16}{13}x + 2$

$$k(x) = -|0.7x| + 5$$

State the solutions to the equation $h(x) = k(x)$, rounded to the *nearest hundredth*.

- 784 The sequence $a_1 = 6$, $a_n = 3a_{n-1}$ can also be written as

1) $a_n = 6 \cdot 3^n$
 2) $a_n = 6 \cdot 3^{n+1}$
 3) $a_n = 2 \cdot 3^n$
 4) $a_n = 2 \cdot 3^{n+1}$

- 785 The value of a new car depreciates over time. Greg purchased a new car in June 2011. The value, V , of his car after t years can be modeled by the equation

$$\log_{0.8} \left(\frac{V}{17000} \right) = t.$$

What is the average decreasing rate of change per year of the value of the car from June 2012 to June 2014, to the *nearest ten dollars per year*?

1) 1960
 2) 2180
 3) 2450
 4) 2770

- 786 Which scenario is best described as an observational study?

1) For a class project, students in Health class ask every tenth student entering the school if they eat breakfast in the morning.
 2) A social researcher wants to learn whether or not there is a link between attendance and grades. She gathers data from 15 school districts.
 3) A researcher wants to learn whether or not there is a link between children's daily amount of physical activity and their overall energy level. During lunch at the local high school, she distributed a short questionnaire to students in the cafeteria.
 4) Sixty seniors taking a course in Advanced Algebra Concepts are randomly divided into two classes. One class uses a graphing calculator all the time, and the other class never uses graphing calculators. A guidance counselor wants to determine whether there is a link between graphing calculator use and students' final exam grades.

787 A study of the annual population of the red-winged blackbird in Ft. Mill, South Carolina, shows the population, $B(t)$, can be represented by the function $B(t) = 750(1.16)^t$, where the t represents the number of years since the study began. In terms of the monthly rate of growth, the population of red-winged blackbirds can be best approximated by the function

- 1) $B(t) = 750(1.012)^t$
- 2) $B(t) = 750(1.012)^{12t}$
- 3) $B(t) = 750(1.16)^{12t}$
- 4) $B(t) = 750(1.16)^{\frac{t}{12}}$

788 In contract negotiations between a local government agency and its workers, it is estimated that there is a 50% chance that an agreement will be reached on the salaries of the workers. It is estimated that there is a 70% chance that there will be an agreement on the insurance benefits. There is a 20% chance that no agreement will be reached on either issue. Find the probability that an agreement will be reached on *both* issues. Based on this answer, determine whether the agreement on salaries and the agreement on insurance are independent events. Justify your answer.

789 Write an explicit formula for a_n , the n th term of the recursively defined sequence below.

$$a_1 = x + 1$$

$$a_n = x(a_{n-1})$$

For what values of x would $a_n = 0$ when $n > 1$?

790 Solve algebraically for all values of x :

$$\sqrt{x-5} + x = 7$$

791 Which equation represents an odd function?

- 1) $y = \sin x$
- 2) $y = \cos x$
- 3) $y = (x+1)^3$
- 4) $y = e^{5x}$

792 The solution set for the equation $\sqrt{56-x} = x$ is

- 1) $\{-8, 7\}$
- 2) $\{-7, 8\}$
- 3) $\{7\}$
- 4) $\{\}$


793 Algebraically determine the values of h and k to correctly complete the identity stated below.

$$2x^3 - 10x^2 + 11x - 7 = (x-4)(2x^2 + hx + 3) + k$$

794 Which value, to the *nearest tenth*, is *not* a solution of $p(x) = q(x)$ if $p(x) = x^3 + 3x^2 - 3x - 1$ and $q(x) = 3x + 8$?

- 1) -3.9
- 2) -1.1
- 3) 2.1
- 4) 4.7

795 The graph of the function $p(x)$ is sketched below.




Which equation could represent $p(x)$?

- 1) $p(x) = (x^2 - 9)(x - 2)$
- 2) $p(x) = x^3 - 2x^2 + 9x + 18$
- 3) $p(x) = (x^2 + 9)(x - 2)$
- 4) $p(x) = x^3 + 2x^2 - 9x - 18$

796 Which factorization is *incorrect*?

- 1) $4k^2 - 49 = (2k + 7)(2k - 7)$
- 2) $a^3 - 8b^3 = (a - 2b)(a^2 + 2ab + 4b^2)$
- 3) $m^3 + 3m^2 - 4m + 12 = (m - 2)^2(m + 3)$
- 4) $t^3 + 5t^2 + 6t + t^2 + 5t + 6 = (t + 1)(t + 2)(t + 3)$


- 797 Elaina has decided to run the Buffalo half-marathon in May. She researched training plans on the Internet and is looking at two possible plans: Jillian's 12-week plan and Josh's 14-week plan. The number of miles run per week for each plan is plotted below.




Which one of the plans follows an arithmetic pattern? Explain how you arrived at your answer. Write a recursive definition to represent the number of miles run each week for the duration of the plan you chose. Jillian's plan has an alternative if Elaina wanted to train instead for a full 26-mile marathon. Week one would start at 13 miles and follow the same pattern for the half-marathon, but it would continue for 14 weeks. Write an explicit formula, in *simplest form*, to represent the number of miles run each week for the full-marathon training plan.

- 798 Mrs. Jones had hundreds of jelly beans in a bag that contained equal numbers of six different flavors. Her student randomly selected four jelly beans and they were all black licorice. Her student complained and said "What are the odds I got all of that kind?" Mrs. Jones replied, "simulate rolling a die 250 times and tell me if four black licorice jelly beans is unusual." Explain how this simulation could be used to solve the problem.
- 800 The guidance department has reported that of the senior class, 2.3% are members of key club, K , 8.6% are enrolled in AP Physics, P , and 1.9% are in both. Determine the probability of P given K , to the *nearest tenth of a percent*. The principal would like a basic interpretation of these results. Write a statement relating your calculated probabilities to student enrollment in the given situation.
- 799 The equation $4x^2 - 24x + 4y^2 + 72y = 76$ is equivalent to
- 1) $4(x - 3)^2 + 4(y + 9)^2 = 76$
 - 2) $4(x - 3)^2 + 4(y + 9)^2 = 121$
 - 3) $4(x - 3)^2 + 4(y + 9)^2 = 166$
 - 4) $4(x - 3)^2 + 4(y + 9)^2 = 436$

- 801 Which equation is represented by the graph shown below?




- 1) $y = \frac{1}{2} \cos 2x$
 - 2) $y = \cos x$
 - 3) $y = \frac{1}{2} \cos x$
 - 4) $y = 2 \cos \frac{1}{2}x$
- 802 There was a study done on oxygen consumption of snails as a function of pH, and the result was a degree 4 polynomial function whose graph is shown below.



Which statement about this function is *incorrect*?

- 803 Stephen's Beverage Company is considering whether to produce a new brand of cola. The company will launch the product if at least 25% of cola drinkers will buy the product. Fifty cola drinkers are randomly selected to take a blind taste-test of products A, B, and the new product. Nine out of fifty participants preferred Stephen's new cola to products A and B. The company then devised a simulation based on the requirement that 25% of cola drinkers will buy the product. Each dot in the graph shown below represents the proportion of people who preferred Stephen's new product, each of sample size 50, simulated 100 times.




Assume the set of data is approximately normal and the company wants to be 95% confident of its results. Does the sample proportion obtained from the blind taste-test, nine out of fifty, fall within the margin of error developed from the simulation? Justify your answer. The company decides to continue developing the product even though only nine out of fifty participants preferred its brand of cola in the taste-test. Describe how the simulation data could be used to support this decision.

- 804 The volume of air in a person's lungs, as the person breathes in and out, can be modeled by a sine graph. A scientist is studying the differences in this volume for people at rest compared to people told to take a deep breath. When examining the graphs, should the scientist focus on the amplitude, period, or midline? Explain your choice.

- 1) The degree of the polynomial is even.
- 2) There is a positive leading coefficient.
- 3) At two pH values, there is a relative maximum value.
- 4) There are two intervals where the function is decreasing.

805 The graph of $p(x)$ is shown below.



What is the remainder when $p(x)$ is divided by $x + 4$?

- 1) $x - 4$
- 2) -4
- 3) 0
- 4) 4

806 Which value is *not* contained in the solution of the system shown below?

$$a + 5b - c = -20$$

$$4a - 5b + 4c = 19$$

$$-a - 5b - 5c = 2$$

- 1) -2
- 2) 2
- 3) 3
- 4) -3


807 Iridium-192 is an isotope of iridium and has a half-life of 73.83 days. If a laboratory experiment begins with 100 grams of Iridium-192, the number of grams, A , of Iridium-192 present after t days

would be $A = 100\left(\frac{1}{2}\right)^{\frac{t}{73.83}}$. Which equation

approximates the amount of Iridium-192 present after t days?

- 1) $A = 100\left(\frac{73.83}{2}\right)^t$
- 2) $A = 100\left(\frac{1}{147.66}\right)^t$
- 3) $A = 100(0.990656)^t$
- 4) $A = 100(0.116381)^t$

808 A cardboard box manufacturing company is building boxes with length represented by $x + 1$, width by $5 - x$, and height by $x - 1$. The volume of the box is modeled by the function below.



Over which interval is the volume of the box changing at the fastest average rate?

- 1) $[1, 2]$
- 2) $[1, 3.5]$
- 3) $[1, 5]$
- 4) $[0, 3.5]$

Algebra II Regents at Random Worksheets

Answer Section

1 ANS:

Based on these data, the two events do not appear to be independent. $P(J) = \frac{145}{277} = 0.52$, while

$P(J|D) = \frac{58}{139} = 0.42$. The probability of being a junior is not the same as the conditional probability of being a junior, given the junior drives to school.

PTS: 2 REF: 062431aii NAT: S.CP.A.4 TOP: Conditional Probability

2 ANS: 3

$$P = 210x^{\frac{4}{3}}y^{\frac{7}{3}} = 210x^{\frac{3}{3}}x^{\frac{1}{3}}y^{\frac{6}{3}}y^{\frac{1}{3}} = 210x \cdot x^{\frac{1}{3}}y^2y^{\frac{1}{3}} = 210xy^2\sqrt[3]{xy}$$

PTS: 2 REF: 012413aii NAT: N.RN.A.2 TOP: Radicals and Rational Exponents

3 ANS:

About 38% $\left(\frac{475}{1250}\right)$ of high school juniors in the population will choose a four-year college.

PTS: 2 REF: 012432aii NAT: S.IC.B.6 TOP: Analysis of Data

KEY: draw conclusions

4 ANS:



25

PTS: 2 REF: 012429aii NAT: S.ID.A.4 TOP: Normal Distributions

KEY: percent

5 ANS:

$$2x^4 - 10x^3 + 3x^2 - 15x = x(2x^3 - 10x^2 + 3x - 15) = x(2x^2(x - 5) + 3(x - 5)) = x(2x^2 + 3)(x - 5)$$

PTS: 2 REF: 082427aii NAT: A.SSE.A.2 TOP: Factoring Polynomials

6 ANS:

$$x = \frac{-3 \pm \sqrt{3^2 - 4(1)(11)}}{2(1)} = \frac{-3 \pm \sqrt{-35}}{2} = \frac{-3}{2} \pm \frac{i\sqrt{35}}{2}$$

PTS: 2 REF: 082432aii NAT: A.REI.B.4 TOP: Solving Quadratics

7 ANS:

$$\frac{8x - 3(x + 5)}{x(x + 5)} = 5$$

$$8x - 3x - 15 = 5x^2 + 25x$$

$$0 = 5x^2 + 20x + 15$$

$$0 = x^2 + 4x + 3$$

$$0 = (x + 3)(x + 1)$$

$$x = -3, -1$$

PTS: 2 REF: 062430aia NAT: A.REI.A.2 TOP: Solving Rationals

8 ANS: 4

$$S_{15} = \frac{10 - 10(1.09)^{15}}{1 - 1.09} \approx 293.609$$

PTS: 2 REF: 062424aia NAT: F.BF.B.7 TOP: Series

KEY: geometric

9 ANS: 2

$$2x \left(2x + \frac{5}{2} = \frac{3}{x} \right)$$

$$4x^2 + 5x = 6$$

$$4x^2 + 5x - 6 = 0$$

$$(4x - 3)(x + 2) = 0$$

$$x = \frac{3}{4}, -2$$

PTS: 2 REF: 012504aia NAT: A.REI.D.11 TOP: Other Systems

KEY: rational


10 ANS: 4

$$S_5 = \frac{350 - 350(1.15)^5}{1 - 1.15} \approx 2360$$

PTS: 2 REF: 012524aia NAT: F.BF.B.7 TOP: Series

KEY: geometric

11 ANS: 2



PTS: 2

REF: 062402aai

NAT: A.REI.D.11

TOP: Other Systems

KEY: rational

12 ANS:

$$C(t) = 130(0.5)^{\frac{t}{5.5}}$$

PTS: 2

REF: 082430aai

NAT: F.BF.A.1

TOP: Modeling Exponential Functions

13 ANS: 4

PTS: 2

REF: 012423aai

NAT: F.BF.B.7

TOP: Series

KEY: geometric

14 ANS: 3

$$6x + 8y - 10z = -54 \quad 6x + 8y - 10z = -54 \quad 6x + 9y - 3z = -9 \quad 10y - 7z = -12$$

$$6x + 9y - 3z = -9 \quad \underline{6x + 9y - 3z = -9} \quad \underline{6x - y + 4z = 3} \quad \underline{y + 7z = 45}$$

$$6x - y + 4z = 3 \quad y + 7z = 45 \quad 10y - 7z = -12 \quad 11y = 33$$

$$y = 3$$

PTS: 2

REF: 082421aai

NAT: A.REI.C.6

TOP: Solving Linear Systems

KEY: three variables

15 ANS: 2

$$(x - 1)^2 = 2x + 6 \quad -1 \text{ is extraneous.}$$

$$x^2 - 2x + 1 = 2x + 6$$

$$x^2 - 4x - 5 = 0$$

$$(x - 5)(x + 1) = 0$$

$$x = 5, -1$$


PTS: 2

REF: 082411aai

NAT: A.REI.A.2

TOP: Solving Radicals

16 ANS:



$$0 < x < \frac{\pi}{4}$$

PTS: 4

REF: 012436aai

NAT: F.IF.C.7

TOP: Graphing Trigonometric Functions

KEY: graph

17 ANS:

$$\frac{\sqrt[5]{a^{10}}}{\left(a^3\right)^{\frac{1}{2}}} = \frac{a^{\frac{10}{5}}}{a^{\frac{3}{2}}} = \frac{a^{\frac{20}{10}}}{a^{\frac{15}{10}}} = a^{\frac{5}{10}} \quad x = \frac{1}{2}$$

PTS: 2

REF: 012528aai

NAT: N.RN.A.2

TOP: Radicals and Rational Exponents

18 ANS: 1

$$8^{\frac{x}{2}} \bullet 8^{\frac{x}{3}} = 8^{\frac{5x}{6}} = \sqrt[6]{8^{5x}}$$

PTS: 2

REF: 082419aai

NAT: N.RN.A.2

TOP: Radicals and Rational Exponents

19 ANS:

$$\frac{3+42}{1500} = 3\% \quad \frac{3}{3+12} = 20\% \quad \text{No, because a person is more likely to be allergic milk if he is also allergic to nuts.}$$

PTS: 4

REF: 012433aai

NAT: S.CP.A.4

TOP: Conditional Probability

20 ANS:

$$a_1 = 12$$

$$a_n = a_{n-1} + 6$$

PTS: 2


REF: 012430aai

NAT: F.BF.A.2

TOP: Sequences

KEY: recursive

21 ANS:

 $x = 1, y = 1$

PTS: 4 REF: 062436aia NAT: F.IF.C.7 TOP: Graphing Logarithmic Functions

22 ANS:

$$x^3 + 4x^2 - 9x - 36 = x^2(x + 4) - 9(x + 4) = (x^2 - 9)(x + 4) = (x + 3)(x - 3)(x + 4)$$

PTS: 2 REF: 012425aia NAT: A.SSE.A.2 TOP: Factoring Polynomials

23 ANS:

Since there is no remainder when the cubic is divided by $x + 3$, this binomial is a factor.

$$\begin{array}{r|rrrr} -3 & 7 & 27 & 9 & -27 \\ & & -21 & -18 & 27 \\ \hline & 7 & 6 & -9 & 0 \end{array}$$

PTS: 2 REF: 082426aia NAT: A.APR.B.2 TOP: Remainder and Factor Theorems

24 ANS: 3

$$2.12 \pm 2(.05)$$

PTS: 2 REF: 012509aia NAT: S.ID.A.4 TOP: Normal Distributions

25 ANS: 2

$$\sqrt{4-x} = x + 8 \qquad -12 + 8 = -4$$

$$4 - x = x^2 + 16x + 64$$

$$0 = x^2 + 17x + 60$$

$$x = (x + 12)(x - 5) + x = -12, 5$$

PTS: 2 REF: 012521aia NAT: A.REI.A.2 TOP: Solving Radicals

26 ANS: 3

$$x^8 - y^8 = (x^4 + y^4)(x^4 - y^4) = (x^4 + y^4)(x^2 + y^2)(x^2 - y^2) = (x^4 + y^4)(x^2 + y^2)(x + y)(x - y)$$

PTS: 2 REF: 082423aia NAT: A.SSE.A.2 TOP: Factoring Polynomials

27 ANS: 2

$$i^2(5x - 2i)^2 = -(25x^2 - 20xi - 4)$$

PTS: 2 REF: 012512aia NAT: N.CN.A.2 TOP: Operations with Complex Numbers

28 ANS: 4 PTS: 2 REF: 012501aii NAT: F.TF.A.2
TOP: Determining Trigonometric Functions KEY: radians

29 ANS: 4

$$A(t) = 150\left((1.02)^{\frac{1}{7}}\right)^{7t} \approx 150(1.00283)^{7t}$$

PTS: 2 REF: 062415aii NAT: A.SSE.B.3 TOP: Modeling Exponential Functions

30 ANS: 1

Distance from the focus to the directrix is 2, so $p = 1$. Vertex is $(-3, 1)$. $y = \frac{1}{4(1)}(x + 3)^2 + 1$

PTS: 2 REF: 012409aii NAT: G.GPE.A.2 TOP: Graphing Quadratic Functions

31 ANS:
left 3, down 5

PTS: 2 REF: 012525aii NAT: F.IF.C.7 TOP: Graphing Logarithmic Functions

32 ANS: 2

$$4x^2 - 7x + 8 = 0 \quad x = \frac{7 \pm \sqrt{(-7)^2 - 4(4)(8)}}{2(4)} = \frac{7 \pm \sqrt{-79}}{8}$$

PTS: 2 REF: 012507aii NAT: A.REI.B.4 TOP: Solving Quadratics
KEY: complex solutions | quadratic formula

33 ANS:

$$\text{Julia: } V(x) = 33,400(0.85^{\frac{1}{12}})^{12x} \approx 33,400(0.9865)^{12x}$$

PTS: 2 REF: 012530aii NAT: A.SSE.B.3 TOP: Modeling Exponential Functions

34 ANS: 1

$$\cos \theta = -\frac{3}{5}; \sec \theta = -\frac{5}{3}$$

PTS: 2 REF: 012421aii NAT: F.TF.C.8 TOP: Determining Trigonometric Functions

35 ANS:

$$6x - 16y + 4z = -120 \quad 6x - 21y - 15z = -93 \quad 6x - 16y + 4z = -120 \quad 6 + z = 3 \quad -6x + 2(6) - 4(-3) = 36$$

$$6x - 21y - 15z = -93 \quad \underline{-6x + 2y - 4z = 36} \quad \underline{-6x + 2y - 4z = 36} \quad z = -3 \quad -6x + 24 = 36$$

$$-6x + 2y - 4z = 36 \quad \underline{-19y - 19z = -57} \quad \underline{-14y = -84} \quad -6x = 12$$

$$y + z = 3 \quad y = 6 \quad x = -2$$

PTS: 4 REF: 062433aii NAT: A.REI.C.6 TOP: Solving Linear Systems
KEY: three variables

36 ANS: 3

$$y = 40(1.2)^8 \approx 168$$

PTS: 2 REF: 062406aii NAT: S.ID.B.6 TOP: Regression
KEY: exponential

37 ANS:

$.795 \pm 2 \cdot .085 = .625 - .965$. Yes, as it is plausible at least .625 of the customers will purchase both.

PTS: 4

REF: 062435aai

NAT: S.IC.A.2

TOP: Analysis of Data

38 ANS: 1

$$2xy^2 \sqrt[3]{x^2 y} = 2x^{\frac{3}{3}} y^{\frac{6}{3}} x^{\frac{2}{3}} y^{\frac{1}{3}} = 2x^{\frac{5}{3}} y^{\frac{7}{3}}$$

PTS: 2

REF: 062413aai

NAT: N.RN.A.2

TOP: Radicals and Rational Exponents

39 ANS: 1

$$P = \frac{2\pi}{\frac{2\pi}{3}} = 3$$

PTS: 2

REF: 082413aai

NAT: F.IF.C.7

TOP: Graphing Trigonometric Functions

KEY: period

40 ANS: 4

$$p(2) = 4(2)^3 - 3(2) + 3 = 29$$

PTS: 2

REF: 062422aai

NAT: A.APR.B.2

TOP: Remainder and Factor Theorems

41 ANS: 4

$$6(2^{x+4}) = 36$$

$$\ln 2^{x+4} = \ln 6$$

$$(x+4)\ln 2 = \ln 6$$

$$x+4 = \frac{\ln 6}{\ln 2}$$

$$x = \frac{\ln 6}{\ln 2} - 4$$

PTS: 2

REF: 082408aai

NAT: F.LE.A.4

TOP: Exponential Equations

KEY: without common base

42 ANS:

Yes. Using a 95% confidence interval, values outside the interval 3.95 – 4.05 are unusual.

PTS: 2

REF: 012532aai

NAT: S.IC.A.2

TOP: Analysis of Data

43 ANS: 4

$$z = 7y - 31 \quad 5x + 2y - (7y - 31) = -14 \rightarrow 5x - 5y = -45 \rightarrow x - y = -9 \rightarrow y = x + 9$$

$$5y + 4(7y - 31) - 5x = -23 \rightarrow -5x + 33y = 101$$

$$-5x + 33(x + 9) = 101$$

$$28x = -196$$

$$x = -7$$

PTS: 2

REF: 012515aai

NAT: A.REI.C.6

TOP: Solving Linear Systems

KEY: three variables

44 ANS: 3

$$3 \mid 1 \ 1 \ -3 \ 9 \ -108 \quad x^3 + 4x^2 + 9x + 36 = 0$$

$$\mid \underline{3 \ 12 \ 27 \ 108} \quad x^2(x + 4) + 9(x + 4) = 0$$

$$1 \ 4 \ 9 \ 36 \quad 0 \quad (x^2 + 9)(x + 4) = 0$$

$$x = \pm 3i, -4$$


PTS: 2

REF: 062420aai

NAT: A.APR.D.6

TOP: Solving Polynomial Equations

45 ANS: 3



PTS: 2

REF: 012406aai

NAT: A.REI.D.11

TOP: Other Systems

KEY: exponential

46 ANS: 4

$$49 \times 16.7\% \approx 8$$

PTS: 2

REF: 062418aai

NAT: S.ID.A.4

TOP: Normal Distributions

KEY: predict

47 ANS: 2

$$x + 1 = \sqrt{4x + 25} \quad -4 + 1 < 0$$

$$x^2 + 2x + 1 = 4x + 25$$

$$x^2 - 2x - 24 = 0$$

$$(x - 6)(x + 4) = 0$$

$$x = 6, -4$$

PTS: 2

REF: 062408aai

NAT: A.REI.A.2

TOP: Solving Radicals

48 ANS:

The opinion sought is that of the entire student body, but the first period computer science class may not be representative of the entire student body.

PTS: 2

REF: 062427aai

NAT: S.IC.B.6

TOP: Analysis of Data

KEY: bias

49 ANS:

$$\frac{55}{t} = \frac{65}{t+3}$$

$$65t = 55t + 165$$

$$10t = 165$$

$$t = 16.5$$

$$t + 3 = 19.5$$

PTS: 2

REF: 082431aai

NAT: A.CED.A.1

TOP: Modeling Rationals

50 ANS: 4

$$y = \sqrt[3]{x+2}$$

$$x = \sqrt[3]{y+2}$$

$$x^3 = y + 2$$

$$y = x^3 - 2$$

PTS: 2

REF: 062419aai

NAT: F.BF.B.4

TOP: Inverse of Functions

KEY: cubic

51 ANS:

$$\frac{H(10) - H(2)}{10 - 2} \approx 11524 \text{ From 2014-2018, the median house price increased \$11524 per year on average.}$$

PTS: 4

REF: 062434aai

NAT: F.IF.B.6

TOP: Rate of Change

KEY: exponential

52 ANS: 3

PTS: 2

REF: 062409aai

NAT: A.REI.B.4

TOP: Using the Discriminant

KEY: determine nature of roots

53 ANS: 2

$$y = x^3 - 3$$

$$x = y^3 - 3$$

$$x + 3 = y^3$$

$$\sqrt[3]{x+3} = y$$

PTS: 2

REF: 012419aai

NAT: F.BF.B.4

TOP: Inverse of Functions

KEY: cubic

54 ANS:
 $2(-1) + 5 = 3$

PTS: 2 REF: 082429aai NAT: F.IF.B.4 TOP: Graphing Trigonometric Functions

55 ANS: 2

$$\left(\frac{1}{x^{-2}}\right)^{-\frac{3}{4}} = \frac{1}{x^{\frac{3}{2}}} = \frac{1}{x^{\frac{2}{2}} \cdot x^{\frac{1}{2}}} = \frac{1}{x\sqrt{x}}$$

PTS: 2 REF: 082412aai NAT: N.RN.A.2 TOP: Radicals and Rational Exponents
 KEY: variables

56 ANS: 1

$$(2x-3)^2 = 4x^2 - 12x + 9 \quad (x-2)^3 = (x-2)(x-2)^2 = (x-2)(x^2 - 4x + 4)$$

$$s = 4 \quad s = -4 \text{ and } 4$$

PTS: 2 REF: 062405aai NAT: A.APR.C.4 TOP: Polynomial Identities

57 ANS:

$$\ln e^{1.5t} = \ln \frac{16}{3.8}$$


$$1.5t = \ln \frac{16}{3.8}$$

$$t = \frac{\ln \frac{16}{3.8}}{1.5} \approx .96$$

PTS: 2 REF: 062426aai NAT: F.LE.A.4 TOP: Exponential Equations
 KEY: without common base

58 ANS:
 No, because a 180° rotation of f about the origin does not map f onto itself.

PTS: 2 REF: 062432aai NAT: F.BF.B.3 TOP: Even and Odd Functions



Since -4 is a zero, $x + 4$ is a factor.

PTS: 2 REF: 012426aai NAT: A.APR.B.2 TOP: Remainder and Factor Theorems

60 ANS:

$42.029 \pm 2 \cdot 3.105 \approx 35.82 - 48.24$. Yes, since 49.8 falls outside the 95% interval.


PTS: 4

REF: 082434aai

NAT: S.IC.A.2

TOP: Analysis of Data

61 ANS:



PTS: 2

REF: 062428aai

NAT: F.IF.C.7

TOP: Graphing Polynomial Functions

62 ANS: 1

$$\frac{4x^2 - 5}{x^2 - 1} = \frac{4(x^2 - 1)}{x^2 - 1} - \frac{1}{x^2 - 1}$$

PTS: 2

REF: 012510aai

NAT: A.APR.D.7

TOP: Addition and Subtraction of Rationals

63 ANS: 1

$$\frac{2\pi}{b} = 12$$

$$12b = 2\pi$$

$$b = \frac{\pi}{6}$$

PTS: 2

REF: 012520aai

NAT: F.IF.C.7

TOP: Graphing Trigonometric Functions

KEY: period

64 ANS: 1

The probability of being late given that a student walked is $\frac{4}{22}$. The probability that student walked given that the student was late is $\frac{4}{30}$.

PTS: 2

REF: 012518aai

NAT: S.CP.A.4

TOP: Conditional Probability

65 ANS: 4

$$P(B) \cdot P(P|B) = P(P \text{ and } B)$$

$$.68 \cdot P(P|B) = .49$$

$$P(P|B) = .72$$

PTS: 2

REF: 062416aai

NAT: S.CP.A.3

TOP: Conditional Probability

66 ANS: 2

$$y = \sqrt[3]{x} + 4$$

$$x = \sqrt[3]{y} + 4$$

$$x - 4 = \sqrt[3]{y}$$

$$(x - 4)^3 = y$$

PTS: 2

REF: 012519aai

NAT: F.BF.B.4

TOP: Inverse of Functions

KEY: cubic

67 ANS: 4

$$5i(2x + 3i) - x\sqrt{-9} = 10xi + 15i^2 - 3xi = -15 + 7xi$$

PTS: 2

REF: 082415aai

NAT: N.CN.A.2

TOP: Operations with Complex Numbers

68 ANS:

$$6x\left(\frac{1}{2x} - \frac{5}{6} = \frac{3}{x}\right)$$

$$3 - 5x = 18$$

$$-15 = 5x$$

$$-3 = x$$


PTS: 2

REF: 012526aai

NAT: A.REI.A.2

TOP: Solving Rationals

69 ANS: 4



$$x^2 + (-2x)^2 = 5 \quad y = -2(-1) = 2$$

$$5x^2 = 5$$

$$x^2 = 1$$

$$x = \pm 1$$


PTS: 2

REF: 012407aai

NAT: A.REI.C.7

TOP: Quadratic-Linear Systems

70 ANS: 2



PTS: 2 REF: 012414aai NAT: F.IF.B.4 TOP: Graphing Polynomial Functions

71 ANS: 1

$$\frac{f(x)}{g(x)} = \frac{2x^2 + 7x - 15}{3 - 2x} = \frac{(2x - 3)(x + 5)}{-(2x - 3)} = \frac{x + 5}{-1} = -x - 5$$

PTS: 2 REF: 012412aai NAT: F.BF.A.1 TOP: Operations with Functions

72 ANS: 2

$$V(x) = x(18 - 2x)(18 - 2x) = x(324 - 72x + 4x^2) = 324x - 72x^2 + 4x^3$$

PTS: 2 REF: 082418aai NAT: F.BF.A.1 TOP: Operations with Functions

73 ANS: 1

$$\text{amplitude} = \frac{8-2}{2} = 3, b = \frac{2\pi}{6} = \frac{\pi}{3}, c = \frac{8+2}{2} = 5$$

PTS: 2 REF: 062403aai NAT: F.TF.B.5 TOP: Modeling Trigonometric Functions

74 ANS:

$$(2xi^3 - 3y)^2 = 4x^2i^6 - 12xyi^3 + 9y^2 = -4x^2 + 12xyi + 9y^2$$

PTS: 2 REF: 012431aai NAT: N.CN.A.2 TOP: Operations with Complex Numbers

75 ANS:

$$\left(-\frac{2}{7}\right)^2 + \sin^2 \theta = 1 \quad \frac{3\sqrt{5}}{7} \text{ as sin is positive in Quadrant II.}$$

$$\frac{4}{49} + \sin^2 \theta = \frac{49}{49}$$

$$\sin^2 \theta = \frac{45}{49}$$

$$\sin \theta = \pm \frac{3\sqrt{5}}{7}$$

PTS: 2 REF: 012527aai NAT: F.TF.C.8 TOP: Determining Trigonometric Functions

76 ANS: 3

PTS: 2

REF: 012404aai

NAT: F.LE.A.4

TOP: Express Exponentials as Logarithms

77 ANS:

No, because $f(-x) = 2^{-x}$ $g(x) = f(x) + 5$ $y = 2^x + 5$

$$2^{-x} \neq 2^x \quad x = 2^y + 5$$

$$\log(x - 5) = \log 2^y$$

$$\frac{\log(x - 5)}{\log 2} = \frac{y \log 2}{\log 2}$$

$$\frac{\log(x - 5)}{\log 2} = h(x)$$

PTS: 4 REF: 082435aai NAT: F.BF.B.5 TOP: Inverse of Functions
KEY: exponential

78 ANS: 2

$$u = x - 2 \quad u^2 + 27u - 90$$

$$(u + 30)(u - 3)$$

$$(x - 2 + 30)(x - 2 - 3)$$

$$(x + 28)(x - 5)$$

PTS: 2 REF: 012503aai NAT: A.SSE.A.2 TOP: Factoring Polynomials

79 ANS: 4 PTS: 2 REF: 062411aai NAT: F.BF.A.1

TOP: Modeling Exponential Functions

80 ANS: 2

$$2^x - 4 > 0$$

$$2^x > 4$$

$$x > 2$$

PTS: 2 REF: 082402aai NAT: F.IF.C.7 TOP: Graphing Exponential Functions

81 ANS: 1

$$\frac{8+12}{120} \bullet \frac{8+40}{120} = \frac{8}{120}$$

$$\frac{1}{6} \bullet \frac{4}{10} = \frac{1}{15}$$

$$\frac{4}{60} = \frac{1}{15}$$

PTS: 2 REF: 082422aai NAT: S.CP.A.4 TOP: Conditional Probability

82 ANS: 3

$$10^{5x-2} = 3$$

$$\log 10^{5x-2} = \log 3$$

$$(5x-2)\log 10 = \log 3$$

$$5x-2 = \log 3$$

$$5x = \log 3 + 2$$

$$x = \frac{\log 3 + 2}{5}$$

PTS: 2


REF: 012517aai

NAT: F.LE.A.4

TOP: Exponential Equations

KEY: without common base

83 ANS:



PTS: 2

REF: 082425aai

NAT: F.IF.C.7

TOP: Graphing Exponential Functions

84 ANS: 4

$$S_{10} = \frac{90000 - 90000(1.02)^{10}}{1 - 1.02} \approx 985,475$$

PTS: 2

REF: 082424aai

NAT: F.BF.B.7

TOP: Series

KEY: geometric

85 ANS: 1

In vertex form, the parabola is $y = \frac{1}{4(2)}(x+5)^2 - 2$. The vertex is $(-5, -2)$ and $p = 2$. $2 + -2 = 0$

PTS: 2

REF: 082416aai

NAT: G.GPE.A.2

TOP: Graphing Quadratic Functions

86 ANS: 3

$$\frac{12(y+1)}{12} = \frac{(x-4)^2}{12}$$

The vertex is $(4, -1)$ and $p = 3$, so the focus is $(4, 2)$. $y = -1 - 3 = -4$

$$y = \frac{1}{4(3)}(x-4)^2 - 1$$

PTS: 2

REF: 062423aai

NAT: G.GPE.A.2

TOP: Graphing Quadratic Functions

87 ANS: 1

PTS: 2

REF: 012506aai

NAT: S.IC.B.3

TOP: Analysis of Data

88 ANS: 2

PTS: 2

REF: 012402aai

NAT: A.REI.B.4

TOP: Using the Discriminant

KEY: determine nature of roots

89 ANS: 3

$$r = \frac{-2\sqrt{3}}{\sqrt{6}} = \frac{-2}{\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}} = \frac{-2\sqrt{2}}{2} = -\sqrt{2} \quad a_7 = \sqrt{6}(-\sqrt{2})^{7-1} = \sqrt{6}(-\sqrt{2})^6 = \sqrt{6} \cdot 2^3 = 8\sqrt{6}$$

PTS: 2

REF: 012410aai

NAT: F.BF.A.2

TOP: Sequences

KEY: explicit

90 ANS: 2

$$x(x^3 + 4x^2 - 9x - 36)$$

$$x(x^2(x+4) - 9(x+4))$$

$$x(x^2 - 9)(x+4)$$

$$x(x+3)(x-3)(x+4)$$

PTS: 2

REF: 062407aai

NAT: A.SSE.A.2

TOP: Factoring Polynomials

91 ANS: 3



$\bar{x} + 2\sigma$ represents approximately 99.1% of the data.

PTS: 2

REF: 012514aai

NAT: S.ID.A.4

TOP: Normal Distributions

KEY: percent

92 ANS: 3

$$\sqrt{x} \bullet \sqrt[4]{x^{11}} = x^{\frac{1}{2}} \bullet x^{\frac{11}{4}} = x^{\frac{2}{4}} \bullet x^{\frac{11}{4}} = x^{\frac{13}{4}}$$

PTS: 2

REF: 012511aai

NAT: N.RN.A.2

TOP: Radicals and Rational Exponents

93 ANS:




PTS: 2

REF: 082428aai

NAT: S.ID.A.4

TOP: Normal Distributions

94 ANS: 2



PTS: 2 REF: 082417aia NAT: A.REI.D.11 TOP: Other Systems

KEY: logarithmic

95 ANS: 3 PTS: 2 REF: 082401aia NAT: S.IC.B.3

TOP: Analysis of Data

96 ANS:

$$\text{China: } \frac{P(120) - P(50)}{120 - 50} \approx 13.5 \quad \text{India: } \frac{1380 - 376.3}{120 - 50} \approx 14.3 \quad \text{India}$$

PTS: 4 REF: 082433aia NAT: F.IF.B.6 TOP: Rate of Change

KEY: exponential

97 ANS: 1

$$1 + \frac{0.027}{12} = 1.00225$$

PTS: 2 REF: 082403aia NAT: A.SSE.B.3 TOP: Modeling Exponential Functions

98 ANS:

$$\frac{20}{8} = 2.5 \quad a_1 = 8$$

$$a_n = 2.5 \cdot a_{n-1}$$

PTS: 2 REF: 012531aia NAT: F.BF.A.2 TOP: Sequences

KEY: recursive

99 ANS:

$$\left(\frac{1}{\sqrt[3]{y^2}} \right) y^4 = \frac{y^{\frac{12}{3}}}{y^{\frac{2}{3}}} = y^{\frac{10}{3}} \quad n = \frac{10}{3}$$

PTS: 2 REF: 012428aia NAT: N.RN.A.2 TOP: Radicals and Rational Exponents

100 ANS: 4

$$\frac{x(x-4)}{(x+3)(x-4)} + \frac{2(x+3)}{(x-4)(x+3)} = \frac{2x+27}{(x-4)(x+3)} \quad -3 \text{ is extraneous.}$$

$$x^2 - 4x + 2x + 6 = 2x + 27$$

$$x^2 - 2x + 6 = 2x + 27$$

$$x^2 - 4x - 21 = 0$$

$$(x-7)(x+3) = 0$$

$$x = 7, -3$$

PTS: 2

REF: 082405aai

NAT: A.REI.A.2

TOP: Solving Rationals

101 ANS: 3

$$\frac{1}{3} + \frac{1}{7} - \frac{9}{21} = \frac{7}{21} + \frac{3}{21} - \frac{9}{21} = \frac{1}{21}$$

PTS: 2

REF: 082410aai

NAT: S.CP.B.7

TOP: Addition Rule

102 ANS: 2

$$3x^2 - 4x + 2 = 2x - 3 \quad x = \frac{6 \pm \sqrt{(-6)^2 - 4(3)(5)}}{2(3)} = \frac{6 \pm \sqrt{-24}}{6} = \frac{6 \pm 2i\sqrt{6}}{6} = 1 \pm \frac{i\sqrt{6}}{3}$$

$$3x^2 - 6x + 5 = 0$$

PTS: 2

REF: 062410aai

NAT: A.REI.B.4

TOP: Solving Quadratics

103 ANS: 3

PTS: 2

REF: 012508aai

NAT: A.APR.B.2

TOP: Remainder and Factor Theorems

104 ANS:

$$.74 \cdot .24 = .1776 \quad .74 + .24 - .1776 = .8024$$

PTS: 4

REF: 012533aai

NAT: S.CP.B.7

TOP: Addition Rule

105 ANS: 1

$$50(1.19^{\frac{1}{12}})^{12t} \approx 50(1.015)^{12t}$$

PTS: 2

REF: 012424aai

NAT: A.SSE.B.3

TOP: Modeling Exponential Functions

106 ANS: 3

$$x^2 + 6x + 9 = -10 + 9$$

$$(x+3)^2 = -1$$

$$x+3 = \pm i$$

$$x = -3 \pm i$$

PTS: 2

REF: 012416aai

NAT: A.REI.B.4

TOP: Solving Quadratics

KEY: complex solutions | completing the square

107 ANS: 1

PTS: 2


REF: 012405aai

NAT: A.APR.B.3

TOP: Graphing Polynomial Functions

- 108 ANS: 3 PTS: 2 REF: 012418aii NAT: S.IC.B.6
TOP: Analysis of Data KEY: draw conclusions
- 109 ANS: 2
 $9.82 \pm 2(1.4)$
- PTS: 2 REF: 012411aii NAT: S.IC.B.6 TOP: Analysis of Data
KEY: draw conclusions
- 110 ANS: 1
 $x^4 + x$
 $x(x^3 + 1)$
 $x(x + 1)(x^2 - x + 1)$
 $(x + 1)(x^3 - x^2 + x)$
- PTS: 2 REF: 082404aii NAT: A.APR.C.4 TOP: Polynomial Identities
- 111 ANS: 1 PTS: 2 REF: 012502aii NAT: S.IC.B.3
TOP: Analysis of Data
- 112 ANS: 2
 $u = x + 3$ $u^2 + 4u - 5$
 $(u + 5)(u - 1)$
 $(x + 3 + 5)(x + 3 - 1)$
 $(x + 8)(x + 2)$
- PTS: 2 REF: 062401aii NAT: A.SSE.A.2 TOP: Factoring Polynomials

113 ANS:



$$A(t) = 10000e^{0.25t}$$

$$B(t) = 6000e^{0.45t}$$

2.6 months for same. $30000 = 10000e^{0.25t}$

$$\ln 3 = \ln e^{0.25t}$$

$$\ln 3 = 0.25t \ln e$$

$$\frac{\ln 3}{0.25} = t$$

$$4.4 \approx t$$

PTS: 6 REF: 012537aia NAT: A.REI.D.11 TOP: Other Systems

KEY: exponential

114 ANS: 2 PTS: 2 REF: 082409aia NAT: F.IF.C.7

TOP: Graphing Logarithmic Functions

115 ANS: 3


$$p(x) = r(x) - c(x)$$

$$-15x^2 + 600x + 60 = -0.4x^2 + 130x + 1200 - c(x)$$

$$c(x) = 14.6x^2 - 470x + 1140$$

PTS: 2 REF: 062421aia NAT: F.BF.A.1 TOP: Operations with Functions

116 ANS:



$$x^3 + 2x^2 - 16x - 32 = 0$$

$$x^2(x+2) - 16(x+2) = 0$$

$$(x^2 - 16)(x+2) = 0$$

$$(x+4)(x-4)(x+2) = 0$$

$$x = -4, 4, -2$$

PTS: 4

REF: 012536aai

NAT: F.IF.C.7

TOP: Graphing Polynomial Functions

117 ANS:

$$A(t) = 4000 \left(1 + \frac{2.4\%}{12} \right)^{12t} \quad B(t) = 3500 \left(1 + \frac{4\%}{4} \right)^{4t} \quad 8.4, \text{ the value of } t \text{ for which } A(t) = B(t)$$

PTS: 4

REF: 012435aai

NAT: A.REI.D.11

TOP: Other Systems

KEY: exponential

118 ANS:

$$(x-4)^2 + ((x-6)-1)^2 = 9 \quad 7-y = 6 \quad 4-y = 6 \quad (7,1), (4,-2)$$

$$x^2 - 8x + 16 + x^2 - 14x + 49 - 9 = 0 \quad 1 = y \quad -2 = y$$

$$2x^2 - 22x + 56 = 0$$

$$x^2 - 11x + 28 = 0$$

$$(x-7)(x-4) = 0$$

$$x = 7, 4$$

PTS: 4

REF: 082436aai

NAT: A.REI.C.7

TOP: Quadratic-Linear Systems

119 ANS: 1

Estimate (0, 50) and (1, 38) as points on the graph. $\frac{38}{50} = 76\%$ implies an estimated 24% rate of decay. Confirmed



with graph of $y = 50(.77)^x$:

PTS: 2

REF: 012516aai

NAT: F.LE.B.5

TOP: Modeling Exponential Functions

120 ANS:

$$112 = 73 + (237 - 73)e^{-1.5k} \quad T(2.5) = 73 + (237 - 73)e^{(-.958)(2.5)} \approx 88 \quad 80 = 73 + (237 - 73)e^{-.958t}$$

$$k \approx .958 \quad t \approx 3.3$$

PTS: 6 REF: 062437aia NAT: F.LE.A.4 TOP: Exponential Decay

121 ANS:

$$3x - 2x^2i + 6i - 4xi^2 + 2x^2i = 3x + 6i + 4x = 7x + 6i$$

PTS: 2 REF: 062425aia NAT: N.CN.A.2 TOP: Operations with Complex Numbers

122 ANS:

$$2x - 6 = 2\sqrt{x - 1} \quad 2 \text{ is extraneous.}$$

$$4x^2 - 24x + 36 = 4(x - 1)$$

$$x^2 - 6x + 9 = x - 1$$

$$x^2 - 7x + 10 = 0$$


$$(x - 5)(x - 2) = 0$$

$$x = 2, 5$$

PTS: 4 REF: 012434aia NAT: A.REI.A.2 TOP: Solving Radicals
KEY: extraneous solutions

123 ANS:

$$\frac{10 - 4}{2} = 7, \text{ so the vertex is } (-2, 7) \text{ and } p = 3. \quad y = -\frac{1}{4(3)}(x + 2)^2 + 7 = y = -\frac{1}{12}(x + 2)^2 + 7$$



PTS: 4 REF: 012535aia NAT: G.GPE.A.2 TOP: Graphing Quadratic Functions

124 ANS:

$$A(t) = 1200 \left(1 + \frac{6.4\%}{4} \right)^{4t} \quad \text{Barnyard because } A(10) \approx 2264.28$$

$$3 = e^{6.35\% t}$$

$$B(18) = 2264.43$$

$$\ln 3 = \ln e^{6.35\% t}$$

$$B(t) = 1200e^{6.35\% t}$$

$$\ln 3 = 0.635t$$

$$\frac{\ln 3}{0.635} = \frac{0.635t}{0.635}$$

$$t \approx 17.3$$

PTS: 6 REF: 082437aia NAT: F.LE.A.4 TOP: Exponential Growth

125 ANS:

$$\frac{V(7) - V(2)}{7 - 2} \approx 48$$

PTS: 2

REF: 012427aai

NAT: F.IF.B.6

TOP: Rate of Change

KEY: exponential

126 ANS: 1

PTS: 2

REF: 082406aai

NAT: S.ID.B.6

TOP: Regression

KEY: choose model

127 ANS: 2

$$3x^2 - 7x + 25 - (7x^2 - 10x + 22) = -4x^2 + 3x + 3$$

PTS: 2

REF: 012513aai

NAT: F.BF.A.1

TOP: Operations with Functions

128 ANS: 4

$$F = 325 - 185e^{-0.4(0)} = 325 - 185 = 140$$

PTS: 2

REF: 012415aai

NAT: F.IF.B.4

TOP: Evaluating Exponential Expressions

129 ANS: 2

$$x^2 - 24 = x - 12 \quad y = -3 - 12 = -15$$

$$x^2 - x - 12 = 0$$

$$(x - 4)(x + 3) = 0$$

$$x = 4, -3$$

PTS: 2

REF: 062404aai

NAT: A.REI.C.7

TOP: Quadratic-Linear Systems

130 ANS: 4

$$(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$$


PTS: 2

REF: 012417aai

NAT: A.APR.C.4

TOP: Polynomial Identities

134 ANS:



(7, 78) If 7000 sweatshirts are sold, the profit is \$78,000. 3,549, because that is when $p(x)$ is first greater than 0.

PTS: 6

REF: 012437aii

NAT: F.IF.C.7

TOP: Graphing Polynomial Functions

135 ANS: 2

$$\begin{array}{r} 3 \quad 1 \quad -1 \quad -21 \quad 45 \quad 0 \\ \quad 3 \quad 6 \quad -45 \quad 0 \\ \quad 1 \quad 2 \quad -15 \quad 0 \quad 0 \end{array}$$

$$x^3 + 2x^2 - 15x = 0$$

$$x(x^2 + 2x - 15) = 0$$

$$x(x^2 + 2x - 15) = 0$$

$$x(x + 5)(x - 3) = 0$$

$$x = 0, -5, 3$$


PTS: 2

REF: 012403aii

NAT: A.APR.B.3

TOP: Solving Polynomial Equations

136 ANS:



PTS: 2

REF: 012529aii

NAT: F.IF.C.7

TOP: Graphing Trigonometric Functions

KEY: graph

137 ANS: 3

$$(x^2 - 49) \left(\frac{7}{x+7} + \frac{4x}{x-7} = \frac{3x+7}{x-7} \right)$$

$$7(x-7) + 4x(x+7) = (3x+7)(x+7)$$

$$7x - 49 + 4x^2 + 28x = 3x^2 + 21x + 7x + 49$$

$$4x^2 + 35x - 49 = 3x^2 + 28x + 49$$

$$x^2 + 7x - 98 = 0$$

$$(x+14)(x-7) = 0$$

$$x = -14, 7$$

PTS: 2 REF: 012422aai NAT: A.REI.A.2 TOP: Solving Rationals

138 ANS: 4

Since the terminal side of θ passes through $(-3, -4)$, $\cos \theta < 0$ and $\sin \theta < 0$. $\cos \theta < 0 \rightarrow \sec \theta < 0$

$$\tan \theta = \frac{\sin \theta}{\cos \theta} \rightarrow \frac{-}{-} = +$$

PTS: 2 REF: 082420aai NAT: F.TF.A.2 TOP: Determining Trigonometric Functions
KEY: extension to reals

139 ANS:

$$\frac{m(4) - m(-1)}{4 - (-1)} = \frac{81 - 1}{5} = 16 \quad p(x) \text{ has a greater rate of change}$$

$$\frac{p(4) - p(-1)}{4 - (-1)} = 16.1\bar{3}$$

PTS: 4 REF: 012534aai NAT: F.IF.B.6 TOP: Rate of Change
KEY: exponential

140 ANS: 4

I. Minimum does not change, only period; II. $-16 + 8 = -8$; III. $\frac{1}{2}(-16) = -8$

PTS: 2 REF: 012523aai NAT: F.BF.B.3 TOP: Transformations with Functions
141 ANS: 4 PTS: 2 REF: 062412aai NAT: F.BF.A.2
TOP: Sequences KEY: recursive

142 ANS: 2

$$i = \frac{6.24\%}{12} = .52\% \quad R = \frac{(18000)(.52\%)}{1 - (1 + .52\%)^{-12 \cdot 6}} \approx 300.36$$

PTS: 2 REF: 012420aai NAT: F.IF.B.4 TOP: Evaluating Exponential Expressions
143 ANS: 1 PTS: 2 REF: 082414aai NAT: F.IF.B.4
TOP: Graphing Polynomial Functions

144 ANS: 2

$$\begin{array}{r}
 2x^3 + 6x^2 + 13x + 42 \\
 x - 3 \overline{) 2x^4 + 0x^3 - 5x^2 + 3x - 2} \\
 \underline{2x^4 - 6x^3} \\
 6x^3 - 5x^2 \\
 \underline{6x^3 - 18x^2} \\
 13x^2 + 3x - 2 \\
 \underline{13x^2 - 39x} \\
 42x - 2 \\
 \underline{42x - 126} \\
 124
 \end{array}$$

PTS: 2 REF: 012408aai NAT: A.APR.D.6 TOP: Rational Expressions
 KEY: division

145 ANS: 3 PTS: 2 REF: 012401aai NAT: S.IC.B.3
 TOP: Analysis of Data

146 ANS: 1

If $\sin \theta = \frac{7}{25}$, $\cos \theta = -\frac{24}{25}$ in QII, and $\tan \theta = \frac{\sin \theta}{\cos \theta} = \frac{\frac{7}{25}}{-\frac{24}{25}} = -\frac{7}{24}$

PTS: 2 REF: 062417aai NAT: F.TF.C.8 TOP: Determining Trigonometric Functions
 147 ANS:

$$\frac{2\pi}{\frac{2\pi}{5}} = 5 \text{ The wheel rotates every 5 minutes.}$$

PTS: 2 REF: 062429aai NAT: F.IF.C.7 TOP: Graphing Trigonometric Functions
 KEY: period

Algebra II Regents at Random Worksheets

Answer Section

148 ANS:

$$y = 2.459(1.616)^x$$

PTS: 2

REF: 012329aai

NAT: S.ID.B.6

TOP: Regression

KEY: exponential

149 ANS: 4

PTS: 2

REF: 012303aai

NAT: F.LE.B.5

TOP: Modeling Exponential Functions

150 ANS: 2

PTS: 2

REF: 082324aai

NAT: A.APR.B.3

TOP: Graphing Polynomial Functions

151 ANS:

$$\sqrt[3]{81} = \sqrt[3]{3^4} = 3^{\frac{4}{3}} \quad a = \frac{4}{3}$$

PTS: 2

REF: 062230aai

NAT: N.RN.A.2

TOP: Radicals and Rational Exponents

KEY: variables

152 ANS: 2

Since the distance from the focus to the directrix is 2, $p = 1$ and the vertex of the parabola is (0,5).

$$y = \frac{1}{4p}(x-h)^2 + k$$

$$y = \frac{1}{4(1)}(x-0)^2 + 5$$

$$y = \frac{1}{4}x^2 + 5$$

$$y - 5 = \frac{1}{4}x^2$$

$$4(y - 5) = x^2$$

PTS: 2

REF: 062323aai

NAT: G.GPE.A.2

TOP: Graphing Quadratic Functions

153 ANS:

$$3x + 7 = x^2 - 2x + 1 \quad -1 \text{ is extraneous.}$$

$$0 = x^2 - 5x - 6$$

$$0 = (x - 6)(x + 1)$$

$$x = 6, -1$$

PTS: 2

REF: 062326aai

NAT: A.REI.A.2

TOP: Solving Radicals

KEY: extraneous solutions

154 ANS:

$\frac{B(10) - B(6)}{10 - 6} \approx -3.88$. The average monthly high temperature decreases about 4° each month from June and October.

PTS: 4

REF: 012336aai

NAT: F.IF.B.6

TOP: Rate of Change

KEY: trigonometric

155 ANS: 3

$$\frac{x+2}{x} + \frac{x}{3} = \frac{2x^2+6}{3x} \quad 0 \text{ is extraneous.}$$

$$\frac{x^2+3x+6}{3x} = \frac{2x^2+6}{3x}$$

$$x^2+3x+6 = 2x^2+6$$

$$x^2-3x = 0$$

$$x(x-3) = 0$$

$$x = 0, 3$$

PTS: 2

REF: 012309aai

NAT: A.REI.A.2

TOP: Solving Rationals

156 ANS:

$$2x^2 - 7x + 4 = 11 - 2x \quad y = 11 - 2\left(\frac{7}{2}\right) = 4 \quad \left\{\left(\frac{7}{2}, 4\right), (-1, 13)\right\}$$

$$2x^2 - 5x - 7 = 0$$

$$y = 11 - 2(-1) = 13$$

$$(2x-7)(x+1) = 0$$

$$x = \frac{7}{2}, -1$$

PTS: 2

REF: 082232aai

NAT: A.REI.C.7

TOP: Quadratic-Linear Systems

157 ANS: 1

$$\frac{20}{14+20+6} = \frac{1}{2}$$

PTS: 2

REF: 082303aai

NAT: S.CP.A.4

TOP: Conditional Probability

158 ANS: 4

$$y = -(x-1)^2 + 5 \quad 3 + y = 4$$

$$4 - x = -x^2 + 2x - 1 + 5 \quad y = 1$$

$$x^2 - 3x = 0$$

$$x(x-3) = 0$$

$$x = 0, 3$$

PTS: 2

REF: 082305aai

NAT: A.REI.C.7

TOP: Quadratic-Linear Systems

159 ANS:
 $0.133696 \times 9256 \approx 1237$

PTS: 2 REF: 082230aai NAT: S.ID.A.4 TOP: Normal Distributions

KEY: predict

160 ANS: 4

$$\text{I. } \left(\frac{y}{x^3}\right)^{-1} = \frac{x^3}{y}; \text{ II. } \sqrt[3]{x^9}(y^{-1}) = \frac{x^{\frac{9}{3}}}{y} = \frac{x^3}{y}; \text{ III. } \frac{x^{6.4}\sqrt[4]{y^8}}{x^3y^3} = \frac{x^3y^{\frac{8}{4}}}{y^3} = \frac{x^3}{y}$$


PTS: 2 REF: 062320aai NAT: N.RN.A.2 TOP: Radicals and Rational Exponents

161 ANS: 4

1) -1 is also a zero. 2) $x^2(x-a) + 16(x-a) = (x^2+16)(x-a)$ a is the only zero. 3) $-a$ is the only zero. 4) $x^2(x-a) - 9(x-a) = (x^2-9)(x-a)$.

PTS: 2 REF: 012019aai NAT: A.APR.B.3 TOP: Solving Polynomial Equations

162 ANS: 4



PTS: 2 REF: 062316aai NAT: S.ID.A.4 TOP: Normal Distributions

KEY: percent

163 ANS: 1

$$\frac{-12}{16} = \frac{9}{-12} = \frac{-6.75}{9}$$

PTS: 2 REF: 012017aai NAT: F.IF.A.3 TOP: Sequences


KEY: difference or ratio

164 ANS: 2

$.43 \pm 2(0.05)$ contains about 95% of the data.

PTS: 2 REF: 062317aai NAT: S.IC.B.4 TOP: Analysis of Data

165 ANS: 2



PTS: 2 REF: 012316aai NAT: F.IF.B.4 TOP: Graphing Polynomial Functions

166 ANS: 3
 $a = 105, 0 < b < 1$

PTS: 2 REF: 082314aai NAT: F.BF.A.1 TOP: Modeling Exponential Functions

167 ANS:
 $m(3) = 3^3 - 3^2 - 5(3) - 3 = 27 - 9 - 15 - 3 = 0$ Since $m(3) = 0$, there is no remainder when $m(x)$ is divided by $x - 3$, and so $x - 3$ is a factor.

PTS: 2 REF: 012026aai NAT: A.APR.B.2 TOP: Remainder and Factor Theorems

168 ANS: 2
 $2x^3 + x^2 - 18x - 9$
 $x^2(2x + 1) - 9(2x + 1)$
 $(x^2 - 9)(2x + 1)$
 $(x + 3)(x - 3)(2x + 1)$

PTS: 2 REF: 082206aai NAT: A.APR.B.2 TOP: Remainder and Factor Theorems

169 ANS: 3 PTS: 2 REF: 012002aai NAT: F.BF.A.1
 TOP: Operations with Functions

170 ANS:

$$\frac{x \cdot x^{\frac{3}{2}}}{x^{\frac{5}{3}}} = \frac{x^{\frac{6}{6}} \cdot x^{\frac{9}{6}}}{x^{\frac{10}{6}}} = x^{\frac{5}{6}}$$

PTS: 2 REF: 082331aai NAT: N.RN.A.2 TOP: Radicals and Rational Exponents

171 ANS:
 $F(t) = 169.136(.971)^t$

PTS: 2 REF: 062232aai NAT: S.ID.B.6 TOP: Regression
 KEY: exponential

172 ANS:
 $(x^3 + 2x - 1)(x^2 + 7) - 3(x^4 - 5x)$
 $x^5 + 7x^3 + 2x^3 + 14x - x^2 - 7 - 3x^4 + 15x$
 $x^5 - 3x^4 + 9x^3 - x^2 + 29x - 7$

PTS: 2 REF: 012330aai NAT: F.BF.A.1 TOP: Operations with Functions

173 ANS: 2 PTS: 2 REF: 082313aai NAT: S.ID.A.4
 TOP: Normal Distributions KEY: percent

174 ANS: 4
 1) $d(2) = 2$; 2) $d(1) = 12$; 3) $d(9) \approx 11$; 4) $d(-1) = 2$

PTS: 2 REF: 062220aai NAT: F.IF.B.4 TOP: Graphing Trigonometric Functions

175 ANS: 3

$$S_{20} = \frac{-2 - (-2)(-3)^{20}}{1 - (-3)} = 1,743,392,200$$

PTS: 2 REF: 012306aai NAT: F.BF.B.7 TOP: Series
KEY: geometric

176 ANS: 4

$$M = \frac{45000 \left(\frac{6.75\%}{12} \right) \left(1 + \frac{6.75\%}{12} \right)^{5 \times 12}}{\left(1 + \frac{6.75\%}{12} \right)^{5 \times 12} - 1} \approx 885.76$$

PTS: 2 REF: 082316aai NAT: F.IF.B.4 TOP: Evaluating Exponential Expressions
177 ANS: 2 PTS: 2 REF: 012311aai NAT: A.APR.C.4
TOP: Polynomial Identities

178 ANS: 4


$$(x - 2i)(x - 2i) = x^2 - 4xi + 4i^2 = x^2 - 4xi - 4$$

PTS: 2 REF: 082202aai NAT: N.CN.A.2 TOP: Operations with Complex Numbers
179 ANS: 2

$$\begin{array}{r} x^2 + 2x + 4 \\ x - 2 \overline{) x^3 - 0x^2 + 0x - 2} \\ \underline{x^3 - 2x^2} \\ 2x^2 + 0x \\ \underline{2x^2 - 4x} \\ 4x - 2 \\ \underline{4x - 8} \\ 6 \end{array}$$

PTS: 2 REF: 082217aai NAT: A.APR.D.6 TOP: Rational Expressions
KEY: division

180 ANS:



PTS: 4

REF: 082234aii

NAT: F.IF.C.7

TOP: Graphing Polynomial Functions

181 ANS:

$$a_n = 100(.8)^{n-1} \quad S_{20} = \frac{100 - 100(.8)^{20}}{1 - .8} \approx 494 \quad \text{No, because } 494 > 40 \times 12.$$

PTS: 4

REF: 012033aii

NAT: F.BF.B.7

TOP: Series

KEY: geometric

182 ANS: 1

$$\frac{x(x^2 - 9)}{-(x^2 - 9)} = -x$$

PTS: 2

REF: 012023aii

NAT: A.APR.D.6

TOP: Rational Expressions

KEY: factoring

183 ANS: 3

To determine student opinion, survey the widest range of students.

PTS: 2

REF: 062202aii

NAT: S.IC.B.6

TOP: Analysis of Data

KEY: bias

184 ANS: 1

PTS: 2

REF: 062318aii

NAT: F.BF.B.3

TOP: Even and Odd Functions

185 ANS:

$$\frac{60 - 20}{4 - 2} = \frac{40}{2} = 20$$

PTS: 2


REF: 082225aii

NAT: F.IF.B.6

TOP: Rate of Change

KEY: graph

186 ANS:



$$N(t) = 400 \sin\left(\frac{2\pi}{5}t\right) + 2400.$$

4 times.

PTS: 6 REF: 062337aai NAT: A.REI.D.11 TOP: Other Systems
KEY: trigonometric

187 ANS: 2

$$\frac{x^2 + 3x}{x^2 + 5x + 6} = \frac{x(x+3)}{(x+2)(x+3)}$$

PTS: 2 REF: 082215aai NAT: A.APR.D.6 TOP: Rational Expressions
KEY: factoring

188 ANS: 2 PTS: 2 REF: 062219aai NAT: F.TF.A.1
TOP: Unit Circle

189 ANS: 4

$$\frac{1}{2}x^2 + 2x = \frac{1}{4}x - 8 \quad b^2 - 4ac$$

$$2x^2 + 8x = x - 32 \quad 7^2 - 4(2)(32) < 0$$

$$2x^2 + 7x + 32 = 0$$

PTS: 2 REF: 012310aai NAT: A.REI.C.7 TOP: Quadratic-Linear Systems

190 ANS:

$$\frac{1200}{1200 + 2016} \approx .373. \text{ Yes, because } \frac{1600}{4288} \approx .373 \text{ also.}$$

PTS: 4 REF: 062334aai NAT: S.CP.A.4 TOP: Conditional Probability

191 ANS: 2

$$2x^4 - x^3 - 16x + 8 = 0$$

$$x^3(2x - 1) - 8(2x - 1) = 0$$

$$(x^3 - 8)(2x - 1) = 0$$

$$x = 2, \frac{1}{2}$$

PTS: 2 REF: 012307aai NAT: A.APR.B.2 TOP: Remainder and Factor Theorems

192 ANS: 4

$$x^3 - x^2yi - xy^2 + x^2yi - xy^2i^2 - y^3i = x^3 - xy^2 - xy^2(-1) - y^3i = x^3 - y^3i$$

PTS: 2

REF: 062223aai

NAT: N.CN.A.2

TOP: Operations with Complex Numbers

193 ANS: 2

$$\sqrt{(-2)^2 + (-3)^2} = \sqrt{13}; \tan \theta = \frac{\sin \theta}{\cos \theta} = \frac{\frac{-3}{\sqrt{13}}}{\frac{-2}{\sqrt{13}}} = \frac{3}{2}$$

PTS: 2

REF: 062304aai

NAT: F.TF.A.2

TOP: Determining Trigonometric Functions

KEY: extension to reals

194 ANS: 4

PTS: 2

REF: 012314aai

NAT: S.IC.B.3

TOP: Analysis of Data

195 ANS:

$$(5xi^3 - 4i)^2 = (-5xi - 4i)^2 = 25x^2i^2 + 40xi^2 + 16i^2 = -25x^2 - 40x - 16$$

PTS: 2

REF: 082329aai

NAT: N.CN.A.2

TOP: Operations with Complex Numbers

196 ANS: 4

PTS: 2

REF: 062309aai

NAT: F.IF.C.9

TOP: Comparing Functions

197 ANS:

$$\left(\frac{y^{\frac{17}{8}}}{y^{\frac{10}{8}}} \right)^{-4} = y^n \quad n = -\frac{7}{2}$$

$$\left(y^{\frac{7}{8}} \right)^{-4} = y^n$$

$$y^{-\frac{7}{2}} = y^n$$

PTS: 2

REF: 082228aai

NAT: N.RN.A.2

TOP: Radicals and Rational Exponents

198 ANS:

$$x^2 + 8x - 5 = 8x - 4$$

$$x^2 - 1 = 0$$

$$x = \pm 1$$

PTS: 2

REF: 082326aai

NAT: A.REI.C.7

TOP: Quadratic-Linear Systems

199 ANS:

Translation 3 units right and 4 units up

PTS: 2

REF: 012027aai

NAT: F.IF.C.7

TOP: Graphing Exponential Functions

200 ANS:

$$\frac{3}{n} = \frac{2}{n^2} \quad 0 \text{ is an extraneous solution.}$$

$$3n^2 = 2n$$

$$3n^2 - 2n = 0$$

$$n(3n - 2) = 0$$

$$n = 0, \frac{2}{3}$$

PTS: 2

REF: 062227aai

NAT: A.REI.A.2

TOP: Solving Rationals

201 ANS: 1

$$2) (x^4 - x^2y^2 + y^4) \neq (x^2 - y^2)(x^2 - y^2); \quad 3) x^6 + y^6 \neq (x^3 + y^3)^2; \quad 4) \frac{x^6 + y^6}{x^2 + y^2} \neq x^6 + y^6 - (x^2 + y^2)$$

PTS: 2

REF: 082219aai

NAT: A.APR.C.4

TOP: Polynomial Identities

202 ANS: 2

PTS: 2

REF: 082222aai

NAT: A.CED.A.1

TOP: Modeling Rationals

203 ANS: 3

PTS: 2

REF: 082214aai

NAT: F.IF.C.7

TOP: Graphing Exponential Functions

204 ANS: 2

$$f(x) = f(-x)$$

$$x^2 + 1 = (-x)^2 + 1$$

$$x^2 + 1 = x^2 + 1$$

PTS: 2

REF: 082323aai

NAT: F.BF.B.3

TOP: Even and Odd Functions

205 ANS: 1

$$\begin{array}{r}
 x^3 - 2x^2 - x + 6 \\
 x + 2 \overline{) x^4 + 0x^3 - 5x^2 + 4x + 14} \\
 \underline{x^4 + 2x^3} \\
 -2x^3 - 5x^2 \\
 \underline{-2x^3 - 4x^2} \\
 -x^2 + 4x + 14 \\
 \underline{-x^2 - 2x} \\
 6x + 14 \\
 \underline{6x + 12} \\
 2
 \end{array}$$

PTS: 2 REF: 012305aia NAT: A.APR.D.6 TOP: Rational Expressions
 KEY: division

206 ANS:

$$\begin{aligned}
 & -x(2x^3 - x^2 - 18x + 9) \\
 & -x(x^2(2x - 1) - 9(2x - 1)) \\
 & \quad -x(x^2 - 9)(2x - 1) \\
 & -x(x + 3)(x - 3)(2x - 1)
 \end{aligned}$$

PTS: 2 REF: 062228aia NAT: A.SSE.A.2 TOP: Factoring Polynomials
 KEY: factoring by grouping

207 ANS:

$$\begin{aligned}
 t &= 2\pi\sqrt{\frac{67}{9.81}} \approx 16.4 \quad 9.6 = 2\pi\sqrt{\frac{L}{9.81}} \\
 L &\approx 22.9
 \end{aligned}$$

PTS: 4 REF: 062234aia NAT: A.REI.A.2 TOP: Solving Radicals
 KEY: context

208 ANS: 1

1) $A(20) > 0$; 2) $.5 \times .5 = .25$; 3) true; 4) $A(7) \approx 9.9$

PTS: 2 REF: 082211aia NAT: F.LE.B.5 TOP: Modeling Exponential Functions

209 ANS:

$$x^2 + (2x - 5)^2 = 25 \quad y + 5 = 2(0) \quad y + 5 = 2(4) \quad (0, -5), (4, 3)$$

$$x^2 + 4x^2 - 20x + 25 = 25 \quad y = -5 \quad y = 3$$

$$5x^2 - 20x = 0$$

$$5x(x - 4) = 0$$

$$x = 0, 4$$

PTS: 4

REF: 062236aai

NAT: A.REI.C.7

TOP: Quadratic-Linear Systems

210 ANS: 2

$$\frac{x^2 + 12}{x^2 + 3} = \frac{x^2 + 3}{x^2 + 3} + \frac{9}{x^2 + 3} = 1 + \frac{9}{x^2 + 3}$$


PTS: 2

REF: 062218aai

NAT: A.APR.D.7

TOP: Addition and Subtraction of Rationals

211 ANS:

vertex (3, 6), focus (3, 1), $p = 5$, directrix $y = 6 + 5 = 11$

PTS: 2

REF: 012028aai

NAT: G.GPE.A.2

TOP: Graphing Quadratic Functions

212 ANS: 2

PTS: 2

REF: 082203aai

NAT: F.IF.C.7

TOP: Graphing Trigonometric Functions KEY: amplitude

213 ANS:

$$\sqrt{49 - 10x} = 2x - 5 \quad -\frac{3}{2} \text{ is extraneous.}$$

$$49 - 10x = 4x^2 - 20x + 25$$

$$0 = 4x^2 - 10x - 24$$

$$0 = 2x^2 - 5x - 12$$

$$0 = (2x + 3)(x - 4)$$

$$x = -\frac{3}{2}, 4$$

PTS: 4

REF: 012333aai

NAT: A.REI.A.2

TOP: Solving Radicals

KEY: extraneous solutions

214 ANS: 3

PTS: 2

REF: 062205aai

NAT: F.BF.B.3

TOP: Transformations with Functions

215 ANS: 2

$$\begin{array}{r}
 2x^2 - 3x + 5 \\
 x + 3 \overline{) 2x^3 + 3x^2 - 4x + 5} \\
 \underline{2x^3 + 6x^2} \\
 -3x^2 - 4x \\
 \underline{-3x^2 - 9x} \\
 5x + 5 \\
 \underline{5x + 15} \\
 -10
 \end{array}$$

PTS: 2 REF: 082302aia NAT: A.APR.D.6 TOP: Rational Expressions
KEY: division

216 ANS:

No. $0.852 \pm 2(0.029) \rightarrow 0.794 - 0.91$. 0.88 falls within this interval.

PTS: 2 REF: 062332aia NAT: S.IC.A.2 TOP: Analysis of Data

217 ANS: 2 PTS: 2 REF: 082308aia NAT: A.REI.B.4
TOP: Using the Discriminant KEY: determine nature of roots

218 ANS: 3

The distance from the vertex to the focus, p , is 4. Since the focus is below the vertex, p is negative.

$$y = -\frac{1}{4(4)}(x - 2)^2 + 1$$

PTS: 2 REF: 082212aia NAT: G.GPE.A.2 TOP: Graphing Quadratic Functions

219 ANS:

$$\frac{P(10.5) - P(0)}{10.5 - 0} \approx 10.76 \text{ fruit flies per day}$$

PTS: 2 REF: 082332aia NAT: F.IF.B.6 TOP: Rate of Change
KEY: exponential

220 ANS:

No, because $P(F / CR) \neq P(F)$

$$\begin{aligned}
 \frac{36}{42 + 36} &\neq \frac{17 + 37 + 36 + 15}{39 + 17 + 42 + 12 + 17 + 37 + 36 + 15} \\
 \frac{36}{78} &\neq \frac{105}{215} \\
 \frac{6}{13} &\neq \frac{21}{43}
 \end{aligned}$$

PTS: 2 REF: 082231aia NAT: S.CP.A.4 TOP: Conditional Probability

221 ANS: 2 PTS: 2 REF: 082204aia NAT: S.IC.B.3
TOP: Analysis of Data

222 ANS: 4

$$f(0) = 4 \sin(2(0)) = 0; \quad g(0) = 3(0)^4 + 2(0)^3 + 7 = 7; \quad h(0) = 5e^{2(0)} + 3 = 8; \quad j(0) = 6 \log_2(3(0) + 4) = 12$$

PTS: 2

REF: 082310aai

NAT: F.IF.C.9

TOP: Comparing Functions

223 ANS:

$$\ln e^{0.49x} = \ln 7.5$$

$$0.49x = \ln 7.5$$

$$x = \frac{\ln 7.5}{0.49} \approx 4.112$$

PTS: 2

REF: 062330aai

NAT: F.LE.A.4

TOP: Exponential Equations

KEY: without common base

224 ANS:

$$\text{a) } p(t) = 11000(2)^{\frac{t}{20}}; \quad \text{b) } \frac{1000000}{11000} = \frac{11000(2)^{\frac{t}{20}}}{11000}$$

$$\log \frac{1000}{11} = \log 2^{\frac{t}{20}}$$

$$\log \frac{1000}{11} = \frac{t \cdot \log 2}{20}$$

$$\frac{20 \log \frac{1000}{11}}{\log 2} = t$$

$$t \approx 130.13$$

PTS: 4

REF: 082233aai

NAT: F.LE.A.4

TOP: Exponential Growth

225 ANS:

$\frac{1}{10}, \frac{1}{5}$, and no, since 0.10 clearly falls within 95% of 0.20.

PTS: 4

REF: 012334aai

NAT: S.IC.A.2

TOP: Analysis of Data

226 ANS:

$.651 \pm 2 \cdot .034 = .58 - .72$. No, since .61 (122/200) falls within the 95% interval.

PTS: 4

REF: 062235aai

NAT: S.IC.A.2

TOP: Analysis of Data

227 ANS: 2

$$1) \frac{29860 - 629}{1910 - 1850} \approx 487; \quad 2) \frac{790390 - 494290}{2010 - 1990} \approx 14805; \quad 3) \frac{251808 - 132459}{1970 - 1950} \approx 5967; \quad 4) \frac{251808 - 14575}{1970 - 1890} \approx 2965$$

PTS: 2

REF: 062301aai

NAT: F.IF.B.6

TOP: Rate of Change

KEY: table

228 ANS: 4

1) is a correct formula, but not recursive

PTS: 2

REF: 082216aaii

NAT: F.BF.A.2

TOP: Sequences

KEY: recursive

229 ANS:

$$(x-2)^2 + (-2x+7-3)^2 = 20 \quad y = -2(0) + 7 = 7 \quad (0, 7), (4, -1)$$

$$(x-2)^2 + (-2x+4)^2 = 20 \quad y = -2(4) + 7 = -1$$

$$x^2 - 4x + 4 + 4x^2 - 16x + 16 = 20$$

$$5x^2 - 20x = 0$$

$$5x(x-4) = 0$$

$$x = 0, 4$$

PTS: 4

REF: 062335aaii

NAT: A.REI.C.7

TOP: Quadratic-Linear Systems

230 ANS:

Pick random names from a list of all students and ask each one his method.

PTS: 2

REF: 062325aaii

NAT: S.IC.B.6

TOP: Analysis of Data

KEY: bias

231 ANS:

$$B(t) = 100(2)^{\frac{t}{30}}$$

PTS: 2

REF: 012031aaii

NAT: F.BF.A.1

TOP: Modeling Exponential Functions

232 ANS:

$$T = (400 - 75)e^{-0.0735t} + 75, \quad 325e^{-0.0735(5)} + 75 \approx 300, \quad 270 = (450 - 75)e^{-8r} + 75, \quad 325e^{-0.0735t} + 75 = 375e^{-0.0817t} + 75$$

$$r \approx 0.0817 \qquad t \approx 17$$

PTS: 6

REF: 012337aaii

NAT: A.REI.D.11

TOP: Other Systems

KEY: exponential

233 ANS: 1

$$\begin{array}{r}
 2x^2 + x - 6 \\
 x + 3 \overline{) 2x^3 + 7x^2 - 3x - 25} \\
 \underline{2x^3 + 6x^2} \\
 x^2 - 3x \\
 \underline{x^2 + 3x} \\
 -6x - 25 \\
 \underline{-6x - 18} \\
 -7
 \end{array}$$

PTS: 2

REF: 062203aai

NAT: A.APR.D.6

TOP: Rational Expressions

KEY: division

234 ANS: 3

$$x^2 - 6x + 9 - (x^2 + 6x + 9) = -12x$$

PTS: 2

REF: 062210aai

NAT: F.BF.A.1

TOP: Operations with Functions

235 ANS: 4

$$\frac{x^2 + 6}{x^2 + 4} = \frac{x^2 + 4}{x^2 + 4} + \frac{2}{x^2 + 4} = 1 + \frac{2}{x^2 + 4}$$


PTS: 2

REF: 082321aai

NAT: A.APR.D.7

TOP: Addition and Subtraction of Rationals

236 ANS:

Domain: $x < 2$, Asymptote $x = 2$

PTS: 4

REF: 012034aai

NAT: F.IF.C.7

TOP: Graphing Logarithmic Functions

237 ANS: 3

$$\frac{4}{k^2 - 8k + 12} = \frac{k(k-6) + (k-2)}{k^2 - 8k + 12} \quad k = 6 \text{ is extraneous}$$

$$4 = k^2 - 6k + k - 2$$

$$0 = k^2 - 5k - 6$$

$$0 = (k-6)(k+1)$$

$$k = 6, -1$$

PTS: 2

REF: 082218aai

NAT: A.REI.A.2

TOP: Solving Rationals

238 ANS:

$$x^3 - 2x^2 - 9x + 18 = x^2(x-2) - 9(x-2) = (x^2 - 9)(x-2) = (x+3)(x-3)(x-2)$$

PTS: 2

REF: 082226aai

NAT: A.SSE.A.2

TOP: Factoring Polynomials

KEY: factoring by grouping

239 ANS: 3

$$\begin{array}{r} 2x+1 \\ x+2 \overline{) 2x^2+5x+8} \end{array}$$

$$\underline{2x^2+4x}$$

$$x+8$$

$$\underline{x+2}$$

$$6$$

PTS: 2


REF: 012007aai

NAT: A.APR.D.6

TOP: Rational Expressions

KEY: division

240 ANS:



PTS: 2

REF: 062231aai

NAT: F.IF.C.7

TOP: Graphing Trigonometric Functions

KEY: graph

241 ANS: 1

The vertical distance from the directrix to the vertex, p , is 2. The vertical distance from the vertex to the focus must also be 2.

PTS: 2 REF: 062213aai NAT: G.GPE.A.2 TOP: Graphing Quadratic Functions

242 ANS:

$$\frac{x-2}{(x-6)(x-2)} + \frac{x(x-6)}{(x-6)(x-2)} = \frac{4}{(x-6)(x-2)}. \quad 6 \text{ is extraneous.}$$

$$x-2+x^2-6x=4$$

$$x^2-5x-6=0$$

$$(x-6)(x+1)=0$$

$$x=6, -1$$

PTS: 4 REF: 082334aai NAT: A.REI.A.2 TOP: Solving Rationals

243 ANS: 2 PTS: 2 REF: 012321aai NAT: F.BF.A.2

TOP: Sequences KEY: recursive

244 ANS: 3

$$\frac{-2}{\sqrt{5^2-2^2}} = \frac{-2}{\sqrt{21}}$$

PTS: 2 REF: 082312aai NAT: F.TF.C.8 TOP: Determining Trigonometric Functions

245 ANS: 3

$$x+y+z=2 \quad x-2y-z=-4 \quad 2x-y=-2 \quad x+2+z=2 \quad x+z=0 \quad 0+2+z=2$$

$$\underline{x-2y-z=-4} \quad \underline{x-9y+z=-18} \quad \underline{2x-11y=-22} \quad x-2(2)-z=-4 \quad \underline{x-z=0} \quad z=0$$

$$2x-y=-2 \quad 2x-11y=-22 \quad 10y=20 \quad 2x=0$$

$$y=2 \quad x=0$$

PTS: 2 REF: 062311aai NAT: A.REI.C.6 TOP: Solving Linear Systems

KEY: three variables

246 ANS:

$$\begin{array}{r}
 3x^2 + 8x + 34 \\
 x - 4 \overline{) 3x^3 - 4x^2 + 2x - 1} \quad 3x^2 + 8x + 34 + \frac{135}{x-4} \quad x = 4 \text{ is not a root of } f(x) \text{ because } \frac{f(x)}{g(x)} \text{ has a remainder.} \\
 \underline{3x^3 - 12x^2} \\
 8x^2 + 2x - 1 \\
 \underline{8x^2 - 32x} \\
 34x - 1 \\
 \underline{34x - 136} \\
 135
 \end{array}$$

PTS: 4 REF: 082235aai NAT: A.APR.D.6 TOP: Rational Expressions
KEY: division

247 ANS: 4

$$g(x): \frac{10-6}{4-2} = 2 \quad t(x): \frac{3-5}{4-2} = 4$$

PTS: 2 REF: 062212aai NAT: F.IF.B.6 TOP: Rate of Change
KEY: graph | table

248 ANS: 3 PTS: 2 REF: 012005aai NAT: F.IF.B.4
TOP: Graphing Polynomial Functions

249 ANS: 2
.962¹⁰ ≈ .679


PTS: 2 REF: 082311aai NAT: A.SSE.B.3 TOP: Modeling Exponential Functions
250 ANS: 2

$$\begin{aligned}
 (x^2 + 3)^2 - 2(x^2 + 3) - 24 \quad \text{let } u = x^2 + 3 \\
 u^2 - 2u - 24 \\
 (u - 6)(u + 4) \\
 (x^2 + 3 - 6)(x^2 + 3 + 4)
 \end{aligned}$$

PTS: 2 REF: 062310aai NAT: A.SSE.A.2 TOP: Factoring Polynomials
251 ANS: 4
 $2 \times 0.035 = 0.07$

PTS: 2 REF: 012319aai NAT: S.IC.B.4 TOP: Analysis of Data

252 ANS:



$$p(x) = (x-2)(x-3)(x+6)$$

PTS: 4

REF: 062333aai

NAT: F.IF.C.7

TOP: Graphing Polynomial Functions

253 ANS:

$$g(3) = 0; \quad 0 = 3^3 + a(3)^2 - 5(3) + 6$$

$$0 = 27 + 9a - 15 + 6$$

$$-18 = 9a$$

$$a = -2$$

PTS: 2

REF: 062328aai

NAT: A.APR.B.2

TOP: Remainder and Factor Theorems

254 ANS:

$$e^{0.0532} > 1, \text{ so } P(t) \text{ is increasing.}$$

PTS: 2

REF: 062327aai

NAT: F.IF.C.7

TOP: Graphing Exponential Functions

255 ANS: 1

$$\begin{array}{r} x^2 - 2x + 5 \\ 2x + 4 \overline{) 2x^3 + 0x^2 + 2x - 7} \end{array}$$

$$\underline{2x^3 + 4x^2}$$

$$-4x^2 + 2x$$

$$\underline{-4x^2 - 8x}$$

$$10x - 7$$

$$\underline{10x + 20}$$

$$-27$$

PTS: 2

REF: 062313aai

NAT: A.APR.D.6

TOP: Rational Expressions

KEY: division

256 ANS:

$$b^2 - 4ac = (-4)^2 - 4(1)(13) = 16 - 52 = -36 \text{ imaginary}$$

PTS: 2

REF: 062225aai

NAT: A.REI.B.4

TOP: Using the Discriminant

KEY: determine nature of roots

257 ANS: 1

$$\left(1.03^{\frac{1}{12}}\right)^{12t} \approx 1.00247^{12t}$$

PTS: 2

REF: 062224aai

NAT: A.SSE.B.3

TOP: Modeling Exponential Functions

258 ANS: 2

PTS: 2

REF: 062222aai

NAT: F.IF.C.9

TOP: Comparing Functions

259 ANS: 3

$$x = \frac{2}{3}y + \frac{1}{6}$$

$$6x = 4y + 1$$

$$4y = 6x - 1$$

$$y = \frac{6}{4}x - \frac{1}{4}$$

PTS: 2

REF: 062321aai

NAT: F.BF.B.4

TOP: Inverse of Functions

KEY: linear

260 ANS: 1

$$\left(a^3\sqrt[3]{2b^2}\right)\left(\sqrt[3]{4a^2b}\right) = a^3\sqrt[3]{8a^2b^3} = 2ab^3\sqrt[3]{a^2}$$

PTS: 2

REF: 082213aai

NAT: N.RN.A.2

TOP: Radicals and Rational Exponents

261 ANS: 2

$$a^5\sqrt[5]{a^4} = a^{\frac{5}{5}} \cdot a^{\frac{4}{5}} = a^{\frac{9}{5}}$$

PTS: 2

REF: 062306aai

NAT: N.RN.A.2

TOP: Radicals and Rational Exponents

262 ANS:

$P(x) = 500(0.97)^x$; 18; The number of palm trees and flamingos will be equal in 18 years.

$$F(x) = 200e^{0.02x}$$

PTS: 4

REF: 062336aai

NAT: A.REI.D.11

TOP: Other Systems

KEY: exponential

263 ANS: 1

The product of the roots equals $(3+i)(3-i) = 9 - i^2 = 10 = \frac{c}{a}$. OR

$$(x - (3+i))(x - (3-i)) = 0$$

$$(x - 3 - i)(x - 3 + i) = 0$$

$$((x - 3) - i)((x - 3) + i) = 0$$

$$(x - 3)^2 - i^2 = 0$$

$$x^2 - 6x + 9 + 1 = 0$$

$$x^2 - 6x + 10 = 0$$

PTS: 2

REF: 082208aai

NAT: A.REI.B.4

TOP: Complex Conjugate Root Theorem

264 ANS: 2

1) $x \rightarrow \infty, f(x) \rightarrow \infty$; 3) quartic polynomial; 4) three real roots

PTS: 2

REF: 012318aai

NAT: F.IF.B.4

TOP: Graphing Polynomial Functions

265 ANS:

$$y = 101.523(.883)^x \quad 29 = 101.523(.883)^x$$

$$\frac{29}{101.523} = (.883)^x$$

$$\log \frac{29}{101.523} = x \log(.883)$$

$$\frac{\log \frac{29}{101.523}}{\log(.883)} = x$$

$$x \approx 10.07$$

PTS: 4

REF: 012036aai

NAT: S.ID.B.6

TOP: Regression

KEY: exponential

266 ANS:

$0.01 \pm 2 \cdot 0.38 = -0.75 - 0.77$. No, since 0.6 falls within the 95% interval.

PTS: 4

REF: 082336aai

NAT: S.IC.B.5

TOP: Analysis of Data

267 ANS: 3

between 000 and 449, inclusive $\rightarrow \frac{450}{1000} = 45\%$

PTS: 2

REF: 012024aai

NAT: S.IC.B.3

TOP: Analysis of Data

268 ANS: 4

(1) and (3) are not recursive

PTS: 2

REF: 012013aai

NAT: F.BF.A.2

TOP: Sequences

KEY: recursive

269 ANS:



43

PTS: 2 REF: 012328aai NAT: S.ID.A.4 TOP: Normal Distributions

KEY: percent

270 ANS: 2

$$4300e^{0.07x} = 5123$$

$$\ln e^{0.07x} = \ln \frac{5123}{4300}$$

$$0.07x = \ln \frac{5123}{4300}$$

$$x = \frac{\ln \frac{5123}{4300}}{0.07}$$

$$x \approx 2.5$$

PTS: 2 REF: 012302aai NAT: F.LE.A.4 TOP: Exponential Equations

KEY: without common base

271 ANS: 4

$$(x^2 - y^2) + (2xy)^2 = x^2 + 4x^2y^2 - y^2$$

$$(x - y) + (x^2 - xy + y^2) = x^2 + x - y - xy + y^2$$

$$(x - y)(x - y)(x^2 + y^2) = (x^2 - 2xy + y^2)(x^2 + y^2) = x^4 - 2x^3y + x^2y^2 + x^2y^2 - 2xy^3 + y^4$$

PTS: 2 REF: 062322aai NAT: A.APR.C.4 TOP: Polynomial Identities

272 ANS:

Yes. $P(B|I) = P(B|G|I)$

$$0.14 + 0.26 = \frac{.14}{.35}$$

$$.4 = .4$$

PTS: 2 REF: 062229aai NAT: S.CP.A.4 TOP: Conditional Probability

273 ANS: 2

The mass of the carbon-14 is decreasing by half every 5715 years.

PTS: 2 REF: 062211aai NAT: F.LE.B.5 TOP: Modeling Exponential Functions

274 ANS: 4

$$\log 2^t = \log \sqrt{10} \quad 2) \frac{\log \sqrt{10}}{\log 2} = \log_2 \sqrt{10}, \quad 1) \log_2 \sqrt{10} = \log_2 10^{\frac{1}{2}} = \frac{1}{2} \log_2 10, \quad 3) \log_4 10 = \frac{\log_2 10}{\log_2 4} = \frac{1}{2} \log_2 10$$

$$t \log 2 = \log \sqrt{10}$$

$$t = \frac{\log \sqrt{10}}{\log 2}$$

PTS: 2 REF: 012009aia NAT: F.LE.A.4 TOP: Exponential Equations

KEY: without common base

275 ANS:

$$a^{x+1} = a^{\frac{2}{3}}$$

$$x+1 = \frac{2}{3}$$

$$x = -\frac{1}{3}$$

PTS: 2 REF: 012326aia NAT: F.LE.A.4 TOP: Exponential Equations

KEY: common base shown

276 ANS: 3

$$\sin^2 A + \left(\frac{\sqrt{5}}{3} \right)^2 = 1 \quad \text{Since } \tan A < 0, \sin A = -\frac{2}{3}$$

$$\sin^2 A + \frac{5}{9} = \frac{9}{9}$$

$$\sin^2 A = \frac{4}{9}$$

$$\sin A = \pm \frac{2}{3}$$

PTS: 2 REF: 012320aia NAT: F.TF.C.8 TOP: Determining Trigonometric Functions

277 ANS: 1

$$1.0325^{\frac{1}{12}} \approx 1.0027$$

PTS: 2 REF: 012323aia NAT: A.SSE.B.3 TOP: Modeling Exponential Functions

278 ANS: 3 PTS: 2 REF: 062302aia NAT: A.SSE.A.2

TOP: Factoring Polynomials

279 ANS: 2

$$2x + 4y - 2z = 2 \quad -x - 3y + 2z = 0 \quad x + y = 2 \quad 3 + 2y - z = 1 \quad 2y - z = -2$$

$$\underline{-x - 3y + 2z = 0} \quad \underline{4x - 8y + 2z = 20} \quad \underline{x - y = 4} \quad \underline{6 - 4y + z = 10} \quad \underline{2(-1) - z = -2}$$

$$x + y = 2 \quad 5x - 5y = 20 \quad 2x = 6 \quad 2y - z = -2 \quad z = 0$$

$$x - y = 4 \quad x = 3 \quad \underline{-4y + z = 4}$$

$$-2y = 2$$

$$y = -1$$

PTS: 2

REF: 062208aai

NAT: A.REI.C.6

TOP: Solving Linear Systems

KEY: three variables

280 ANS:

Light wave C. The periods for A, B, and C are 280, 220 and 320.

PTS: 2

REF: 012030aai

NAT: F.IF.C.7

TOP: Graphing Trigonometric Functions

KEY: period

281 ANS: 2

$$1 = \frac{2\pi}{k}$$

$$k = 2\pi$$


PTS: 2

REF: 012313aai

NAT: F.TF.B.5

TOP: Modeling Trigonometric Functions

282 ANS:



$$y = -x + 1 \quad y = -2 + 1 = -1 \quad (2, -1)$$

$$(x - 2)^2 + (-x + 1 - 3)^2 = 16 \quad y = 2 + 1 = 3 \quad (-2, 3)$$

$$x^2 - 4x + 4 + x^2 + 4x + 4 = 16$$

$$2x^2 = 8$$

$$x = -2, 2$$

PTS: 4

REF: 012035aai

NAT: A.REI.C.7

TOP: Quadratic-Linear Systems

283 ANS: 3

PTS: 2

REF: 012003aai

NAT: A.APR.C.4

TOP: Polynomial Identities

284 ANS: 3

$$(6 - ki)^2 = 27 - 36i$$

$$36 - 12ki + k^2 i^2 = 27 - 36i$$

$$9 - k^2 - 12ki = -36i$$

Set real part equal to real part: $9 - k^2 = 0$ Set imaginary part equal to imaginary part: $-12ki = -36i$

$$k = \pm 3$$

$$\frac{-12ki}{-12i} = \frac{-36i}{-12i}$$

$$k = 3$$

PTS: 2

REF: 012308aai

NAT: N.CN.A.2

TOP: Operations with Complex Numbers

285 ANS: 4

PTS: 2

REF: 012014aai

NAT: S.IC.B.5

TOP: Analysis of Data

286 ANS: 4

$$\frac{15000}{12000} = \frac{12000e^{.025t}}{12000}$$

$$1.25 = e^{.025t}$$

$$\ln 1.25 = \ln e^{.025t}$$

$$\ln 1.25 = .025t$$

$$\frac{\ln 1.25}{.025} = t$$

PTS: 2

REF: 082209aai

NAT: F.LE.A.4

TOP: Exponential Growth

287 ANS: 3

$$M = \frac{240000 \left(\frac{4.5\%}{12} \right) \left(1 + \frac{4.5\%}{12} \right)^{15 \times 12}}{\left(1 + \frac{4.5\%}{12} \right)^{15 \times 12} - 1} \approx 1835.98$$

PTS: 2

REF: 062209aai

NAT: F.IF.B.4

TOP: Evaluating Exponential Expressions

288 ANS:

 p is the distance from the focus to the vertex: $8 - 7 = 1$. p is the distance from the directrix to the vertex:

$$1 = 7 - d. \quad y = 6$$

$$d = 6$$

PTS: 2

REF: 082330aai

NAT: G.GPE.A.2

TOP: Graphing Quadratic Functions

289 ANS: 3

PTS: 2

REF: 012015aai

NAT: S.IC.B.3

TOP: Analysis of Data

290 ANS: 1

PTS: 2

REF: 062214aai

NAT: S.ID.A.4

TOP: Normal Distributions

KEY: predict

291 ANS:

$$\sqrt{4x+1} = 11-x \quad 20 \text{ is extraneous.}$$

$$4x+1 = 121-22x+x^2$$

$$0 = x^2 - 26x + 120$$

$$0 = (x-6)(x-20)$$

$$x = 6, 20$$

PTS: 2


REF: 082227aai

NAT: A.REI.A.2

TOP: Solving Radicals

KEY: extraneous solutions

292 ANS:



35; $B(100) - A(100) \approx 2$, which represents the difference of the average wait time when there are 100 CSRs between the plans.

PTS: 6


REF: 082237aai

NAT: A.REI.D.11

TOP: Other Systems

KEY: exponential

293 ANS: 4



$$s(x) = x^4 - 9x^2 + 3x^3 - 27x - 10x^2 + 90$$

$$= x^2(x^2 - 9) + 3x(x^2 - 9) - 10(x^2 - 9)$$

$$= (x^2 + 3x - 10)(x^2 - 9)$$

$$= (x+5)(x-2)(x+3)(x-3)$$

PTS: 2

REF: 062303aai

NAT: A.APR.B.3

TOP: Solving Polynomial Equations

294 ANS: 1

$$7 - 3i + x^2 - 4xi + 4i^2 - 4i - 2x^2 = 7 - 7i - x^2 - 4xi - 4 = 3 - x^2 - 4xi - 7i = (3 - x^2) - (4x + 7)i$$

PTS: 2

REF: 012022aai

NAT: N.CN.A.2

TOP: Operations with Complex Numbers

295 ANS: 4

Translate the parent log function 2 to the right and reflect over the x -axis.

PTS: 2 REF: 082207aia NAT: F.IF.C.7 TOP: Graphing Logarithmic Functions

296 ANS: 2

$$a_2 = 8 + \log_{2+1} 1 = 8 + 0 = 8$$

$$a_3 = 8 + \log_{3+1} 2 = 8 + \frac{1}{2} = 8.5$$

PTS: 2 REF: 062221aia NAT: F.IF.A.3 TOP: Sequences

KEY: recursive

297 ANS: 1

$$x + y + z = 9 \quad 4 - y - z = -1 \quad 4 - 6 + z = 9$$

$$\underline{x - y - z = -1} \quad 4 - y + z = 21 \quad z = 11$$

$$2x = 8 \quad -y - z = -5$$

$$x = 4 \quad \underline{-y + z = 17}$$

$$-2y = 12$$

$$y = -6$$

PTS: 2 REF: 012018aia NAT: A.REI.C.6 TOP: Solving Linear Systems

KEY: three variables

298 ANS: 4 PTS: 2 REF: 082205aia NAT: F.TF.A.2

TOP: Unit Circle

299 ANS: 1

$$2000 \left(1 + \frac{.032}{12} \right)^{12t} \approx 2000(1.003)^{12t}$$

PTS: 2 REF: 012004aia NAT: F.BF.A.1 TOP: Modeling Exponential Functions

300 ANS: 1

$$\ln e^{x+2} = \ln \frac{7}{5}$$


$$(x+2) \ln e = \ln \frac{7}{5}$$

$$x = -2 + \ln \frac{7}{5}$$

PTS: 2 REF: 062207aia NAT: F.LE.A.4 TOP: Exponential Equations

KEY: without common base

301 ANS:



PTS: 2

REF: 082328aai

NAT: F.IF.C.7

TOP: Graphing Trigonometric Functions

KEY: graph

302 ANS:

$$\text{antibiotic } n(0) = \frac{0+1}{0+5} + \frac{18}{0^2 + 8(0) + 15} = \frac{3}{15} + \frac{18}{15} = \frac{21}{15}$$

$$\frac{t+1}{t+5} + \frac{18}{t^2 + 8t + 15} = \frac{9}{t+3}$$

$$a(0) = \frac{9}{0+3} = 3$$

$$\frac{(t+1)(t+3)}{(t+5)(t+3)} + \frac{18}{(t+3)(t+5)} = \frac{9(t+5)}{(t+3)(t+5)}$$

$$t^2 + 4t + 3 + 18 = 9t + 45$$

$$t^2 - 5t - 24 = 0$$

$$(t-8)(t+3) = 0$$

$$t = 8$$

PTS: 6

REF: 012037aai

NAT: A.REI.D.11

TOP: Other Systems

KEY: rational

303 ANS: 4

PTS: 2

REF: 082220aai

NAT: F.IF.B.4

TOP: Graphing Trigonometric Functions

304 ANS: 1

$$50(.9)^t = 25$$

$$t \approx 6.57$$

PTS: 2

REF: 082317aai

NAT: F.LE.A.4

TOP: Exponential Decay

305 ANS:

$$\cos A = \frac{\cos A}{\sin A}$$

$$-3 = \frac{\frac{3}{\sqrt{10}}}{\sin A}$$

$$\sin A = \frac{3}{-3\sqrt{10}} = -\frac{1}{\sqrt{10}}$$


PTS: 2

REF: 082229aai

NAT: F.TF.C.8

TOP: Determining Trigonometric Functions

306 ANS: 3



PTS: 2 REF: 012317aia NAT: A.REI.D.11 TOP: Other Systems

KEY: logarithmic

307 ANS: 1 PTS: 2 REF: 062201aia NAT: N.RN.A.2

TOP: Radicals and Rational Exponents

308 ANS: 3

$$(x+a)^2 + 5(x+a) + 4 \text{ let } u = x+a$$

$$u^2 + 5u + 4$$

$$(u+4)(u+1)$$

$$(x+a+4)(x+a+1)$$

PTS: 2 REF: 012006aia NAT: A.SSE.A.2 TOP: Factoring Polynomials

KEY: multivariable

309 ANS: 4

The distance between the focus and directrix is $1 - (-3) = 4$. p is half this distance, or 2. The vertex of the parabola is $(4, -1)$. Since the directrix is above the focus, the parabola faces downward. $y = -\frac{1}{4p}(x-h)^2 + k$

$$y = -\frac{1}{4(2)}(x-4)^2 - 1$$

$$y+1 = -\frac{1}{8}(x-4)^2$$

PTS: 2 REF: 012322aia NAT: G.GPE.A.2 TOP: Graphing Quadratic Functions

310 ANS: 1

$$u = x+2 \quad u^2 - 5u + 6$$

$$(u-3)(u-2)$$

$$(x+2-3)(x+2-2)$$

$$(x-1)x$$

PTS: 2 REF: 012301aia NAT: A.SSE.A.2 TOP: Factoring Polynomials

KEY: higher power

311 ANS: 3

$$x = 12y - 4$$

$$x + 4 = 12y$$

$$\frac{x+4}{12} = y$$

PTS: 2

REF: 082304aai

NAT: F.BF.B.4

TOP: Inverse of Functions

KEY: linear

312 ANS:

 $\pi < \theta < 2\pi \rightarrow$ Quadrant III or IV θ must be in Quadrant IV, where $\tan \theta$ is negative.

$$\cos \theta = \frac{\sqrt{3}}{4} \rightarrow \text{Quadrant I or IV}$$

PTS: 2

REF: 012332aai

NAT: F.TF.A.2

TOP: Finding the Terminal Side of an Angle

313 ANS: 4

$$45\% + 31\% - 58\% = 18\%$$

PTS: 2

REF: 082307aai

NAT: S.CP.B.7

TOP: Addition Rule

314 ANS:

$$2x^3 - 3x^2 - 18x + 27$$

$$x^2(2x - 3) - 9(2x - 3)$$

$$(x^2 - 9)(2x - 3)$$

$$(x + 3)(x - 3)(2x - 3)$$

PTS: 2

REF: 082325aai

NAT: A.SSE.A.2

TOP: Factoring Polynomials

315 ANS:

$$x^4 - 5x^2 + 4$$

$$(x^2 - 4)(x^2 - 1)$$

$$(x + 2)(x - 2)(x + 1)(x - 1)$$

PTS: 2

REF: 012331aai

NAT: A.SSE.A.2

TOP: Factoring Polynomials

316 ANS: 2

PTS: 2

REF: 062206aai

NAT: A.APR.B.2

TOP: Remainder and Factor Theorems

317 ANS:

$$\frac{x^2(2x+1) - 9(2x+1)}{x(3-x)} = \frac{(x^2-9)(2x+1)}{x(3-x)} = \frac{(x+3)(x-3)(2x+1)}{x(3-x)} = \frac{(x+3)(2x+1)}{-x}$$

PTS: 2


REF: 062331ai

NAT: A.APR.D.6

TOP: Rational Expressions

KEY: factoring

318 ANS:



3

PTS: 4 REF: 062233aai NAT: A.REI.D.11 TOP: Other Systems
KEY: polynomial

319 ANS: 4

$$\frac{13}{13+11} = \frac{13}{24}$$

PTS: 2 REF: 012011aai NAT: S.CP.A.4 TOP: Conditional Probability
320 ANS: 4 PTS: 2 REF: 082318aai NAT: F.IF.B.4
TOP: Graphing Polynomial Functions

321 ANS: 4 PTS: 2 REF: 082301aai NAT: S.IC.B.6
TOP: Analysis of Data KEY: bias

322 ANS: 3 PTS: 2 REF: 082201aai NAT: S.IC.B.6
TOP: Analysis of Data KEY: draw conclusions

323 ANS: 1

$$\frac{N(6) - N(0)}{6 - 0} \approx -8.93$$

PTS: 2 REF: 012012aai NAT: F.IF.B.6 TOP: Rate of Change
KEY: exponential


324 ANS: 1

$$\begin{array}{r|rrrrrr} -2 & 1 & -1 & -11 & 5 & 30 \\ & & -2 & 6 & 10 & -30 \\ \hline & 1 & -3 & -5 & 15 & 0 \end{array}$$

Since there is no remainder when the quartic is divided by $x + 2$, this binomial is a factor.

PTS: 2 REF: 082320aai NAT: A.APR.B.2 TOP: Remainder and Factor Theorems

325 ANS:

As $x \rightarrow -3, y \rightarrow \infty$. As $x \rightarrow \infty, y \rightarrow -\infty$.

PTS: 4

REF: 082333aii

NAT: F.IF.C.7

TOP: Graphing Logarithmic Functions

326 ANS: 4

PTS: 2

REF: 062216aii

NAT: S.IC.B.3

TOP: Analysis of Data

327 ANS:

$$A(t) = 8000 \left(1 + \frac{.042}{4} \right)^{4t} \quad A(18) = 16970.900 \quad 24000 = 8000e^{.039t}$$

$$B(t) = 8000e^{.039t} \quad B(18) = \underline{16142.274} \quad \ln 3 = \ln e^{.039t}$$

$$828.63 \quad \ln 3 = .039t$$

$$t \approx 28.2$$

PTS: 6

REF: 082337aii

NAT: F.LE.A.4

TOP: Exponential Growth

328 ANS:

$$3(x^3 + 4x^2 - x - 4) = 0$$

$$(x^2(x+4) - (x+4)) = 0$$

$$(x^2 - 1)(x+4) = 0$$

$$x = \pm 1, -4$$


PTS: 2

REF: 012325aii

NAT: A.APR.B.3

TOP: Solving Polynomial Equations

329 ANS: 2



PTS: 2

REF: 082319aii

NAT: A.REI.D.11

TOP: Other Systems

KEY: polynomial

330 ANS: 4

PTS: 2

REF: 012016aii

NAT: F.IF.B.4

TOP: Graphing Trigonometric Functions KEY: increasing/decreasing

331 ANS: 3

$$e^{\left(-\frac{3}{0.6}\right)} \approx 0.006738$$

PTS: 2

REF: 062315aai

NAT: A.SSE.B.3

TOP: Modeling Exponential Functions

332 ANS:

$$\begin{array}{rclclclcl} 2x + 4y - 3z = 12 & 2x + 4y - 3z = 12 & 8x + z = -6 & 32x + 4z = -24 & 8(-1) + z = -6 & -(-1) + y - 3(2) = 0 \\ 2(3x - 2y + 2z = -9) & 6x - 4y + 4z = -18 & 2x - 8z = -18 & \underline{x - 4z = -9} & z = 2 & y = 5 \\ 4(-x + y - 3z = 0) & -4x + 4y - 12z = 0 & & 33x = -33 & & \\ & & & x = -1 & & \end{array}$$

PTS: 4

REF: 082335aai

NAT: A.REI.C.6

TOP: Solving Linear Systems

KEY: three variables

333 ANS: 3

$$x = \frac{1}{2}y + 2$$

$$2x = y + 4$$

$$y = 2x - 4$$

PTS: 2

REF: 012315aai

NAT: F.BF.B.4

TOP: Inverse of Functions

KEY: linear

334 ANS: 1


PTS: 2

REF: 062308aai

NAT: F.IF.C.7

TOP: Graphing Logarithmic Functions

335 ANS: 1



PTS: 2

REF: 082210aai

NAT: A.REI.D.11

TOP: Other Systems

KEY: rational

336 ANS: 1

$$\frac{(x+3)(x+2)}{(x-5)(x+2)} + \frac{6(x-5)}{(x+2)(x-5)} = \frac{6+10x}{(x-5)(x+2)} \quad 5 \text{ is extraneous.}$$

$$x^2 + 5x + 6 + 6x - 30 = 10x + 6$$

$$x^2 + x - 30 = 0$$

$$(x+6)(x-5) = 0$$

$$x = -6, 5$$

PTS: 2

REF: 062319aai

NAT: A.REI.A.2

TOP: Solving Rationals

337 ANS: 1


PTS: 2

REF: 082309aai

NAT: F.BF.A.1

TOP: Modeling Exponential Functions

338 ANS: 2



PTS: 2

REF: 012021aai

NAT: A.REI.D.11

TOP: Other Systems

KEY: trigonometric

339 ANS:

$$\left(p^2 n^{\frac{1}{2}}\right)^8 \sqrt{p^5 n^4} = \left(p^{16} n^4\right) p^2 n^2 \sqrt{p} = p^{18} n^6 \sqrt{p}$$

PTS: 2

REF: 012025aai

NAT: N.RN.A.2

TOP: Radicals and Rational Exponents

340 ANS:

$.819 \pm 2 \cdot .053 = .713 - .925$. Since $.70$ does not fall within the 95% interval.

PTS: 4

REF: 082236aai

NAT: S.IC.A.2

TOP: Analysis of Data

341 ANS: 3

$$y = 1.77(1.18)^x \quad y(41) \approx 1,850,950$$

PTS: 2

REF: 062314aai

NAT: S.ID.B.6

TOP: Regression

KEY: exponential

342 ANS:

$$\frac{63}{189} = \frac{1}{3} a_1 = 189$$

$$a_n = \frac{1}{3} a_{n-1}$$

PTS: 2 REF: 062329aai NAT: F.BF.A.2 TOP: Sequences
KEY: recursive

343 ANS: 2

$$x^2 = 3x + 40. \quad x = -5 \text{ is an extraneous solution.}$$

$$x^2 - 3x - 40 = 0$$

$$(x - 8)(x + 5) = 0$$

$$x = 8, -5$$

PTS: 2 REF: 012010aai NAT: A.REI.A.2 TOP: Solving Radicals
KEY: extraneous solutions

344 ANS: 1

$$x^2 - 4x + 4 = -13 + 4$$

$$(x - 2)^2 = -9$$

$$x - 2 = \pm 3i$$

$$x = 2 \pm 3i$$

PTS: 2 REF: 062312aai NAT: A.REI.B.4 TOP: Solving Quadratics
KEY: complex solutions | completing the square

345 ANS: 4 PTS: 2 REF: 012008aai NAT: S.CP.A.3
TOP: Conditional Probability

346 ANS: 3

$$3i(ai - 6i^2) = 3ai^2 - 18i^3 = -3a + 18i$$

PTS: 2 REF: 062307aai NAT: N.CN.A.2 TOP: Operations with Complex Numbers
347 ANS: 2 PTS: 2 REF: 062324aai NAT: F.BF.B.7
TOP: Series KEY: geometric

348 ANS:

$$S_5 = \frac{6 - 6(.8)^5}{1 - .8} \approx 20.17$$

PTS: 2 REF: 062226aai NAT: F.BF.B.7 TOP: Series
KEY: geometric

349 ANS: 2

$$p(x) = 4^x, q(x) = \left(\frac{5}{9}\right)^x, r(x) = 5.29^x, s(x) = 2^x$$

PTS: 2 REF: 012304aai NAT: F.IF.C.7 TOP: Graphing Exponential Functions

350 ANS: 2

1) 1 real, mult. 2; 3) not a quadratic; 4) not a function.

PTS: 2

REF: 012324aai

NAT: A.REI.B.4

TOP: Using the Discriminant

KEY: determine nature of roots

351 ANS: 3

$$y = -6x + \frac{1}{2}$$

$$x = -6y + \frac{1}{2}$$

$$x - \frac{1}{2} = -6y$$

$$-\frac{1}{6}\left(x - \frac{1}{2}\right) = y$$

PTS: 2

REF: 062217aai

NAT: F.BF.B.4

TOP: Inverse of Functions

KEY: linear

352 ANS: 1

$$0.5^{\frac{1}{0.0803}} \approx 0.000178$$

PTS: 2

REF: 082224aai

NAT: A.SSE.B.3

TOP: Modeling Exponential Functions

353 ANS:

$$x = \frac{-5 \pm \sqrt{5^2 - 4(3)(8)}}{2(3)} = -\frac{5}{6} \pm \frac{i\sqrt{71}}{6}$$

PTS: 2

REF: 082327aai

NAT: A.REI.B.4

TOP: Solving Quadratics

KEY: complex solutions | quadratic formula

354 ANS: 2

$$-23(1) + 56 = 33; -23(-1) + 56 = 79$$

PTS: 2

REF: 062305aai

NAT: F.IF.B.4

TOP: Graphing Trigonometric Functions

355 ANS: 3

$$x = -\frac{2y}{5} + 4 \quad y = -\frac{5}{2}(6) + 10 = -5$$

$$5x = -2y + 20$$

$$2y = -5x + 20$$

$$y = -\frac{5}{2}x + 10$$

PTS: 2

REF: 082223aai

NAT: F.BF.B.4

TOP: Inverse of Functions

KEY: linear

356 ANS: 4

PTS: 2

REF: 062215aai

NAT: F.IF.C.7

TOP: Graphing Logarithmic Functions

357 ANS: 1 PTS: 2 REF: 082221aii NAT: F.BF.B.6
TOP: Sigma Notation KEY: represent

358 ANS: 2

$$5x^2 - 4x + 2 = 0 \quad \frac{4 \pm \sqrt{(-4)^2 - 4(5)(2)}}{2(5)} = \frac{4 \pm \sqrt{-24}}{10} = \frac{4 \pm 2i\sqrt{6}}{10} = \frac{2 \pm i\sqrt{6}}{5}$$

PTS: 2 REF: 012020aii NAT: A.REI.B.4 TOP: Solving Quadratics
KEY: complex solutions | quadratic formula

359 ANS:

$$1.5\%; P(t) = 92.2(1.015)^t; \quad \frac{300}{92.2} = (1.015)^t$$


$$\log \frac{300}{92.2} = t \log(1.015)$$

$$\frac{\log \frac{300}{92.2}}{\log(1.015)} = t$$

$$t \approx 79$$

PTS: 6 REF: 062237aii NAT: F.BF.A.2 TOP: Sequences
KEY: recursive

360 ANS:



As $x \rightarrow \infty, c(x) \rightarrow -\infty$. As $x \rightarrow -\infty, c(x) \rightarrow 2$.

PTS: 4 REF: 012335aii NAT: F.IF.C.7 TOP: Graphing Exponential Functions
361 ANS:

$$\frac{1}{3} \times \frac{5}{12} = \frac{5}{36}$$

PTS: 2 REF: 012327aii NAT: S.CP.A.2 TOP: Probability of Compound Events

362 ANS: 1

$$\log 3^{x+4} = \log 28$$

$$\frac{(x+4)\log 3}{\log 3} = \frac{\log 28}{\log 3}$$

$$x+4 = \frac{\log 28}{\log 3}$$

$$x = \log_3 28 - 4$$

PTS: 2 REF: 082306aai NAT: F.LE.A.4 TOP: Exponential Equations
KEY: without common base

363 ANS:

$$r = \frac{360}{300} = 1.2 \quad S_n = \frac{300 - 300(1.2)^n}{1 - 1.2} \quad S_{10} = \frac{300 - 300(1.2)^{10}}{1 - 1.2} \approx 7787.6$$

PTS: 2 REF: 012029aai NAT: F.BF.B.7 TOP: Series
KEY: geometric

364 ANS: 3

$$\frac{x^{\frac{1}{5}}}{x^{\frac{1}{2}}} = x^{\frac{1}{5} - \frac{1}{2}} = x^{-\frac{3}{10}} = \frac{1}{x^{\frac{3}{10}}} = \frac{1}{\sqrt[10]{x^3}}$$

PTS: 2 REF: 012312aai NAT: N.RN.A.2 TOP: Radicals and Rational Exponents

365 ANS: 2

$$40 - (20 + 22 - 15) = 13$$

PTS: 2 REF: 062204aai NAT: S.CP.B.7 TOP: Addition Rule

366 ANS: 1

$$\sqrt[4]{81x^8y^6} = 81^{\frac{1}{4}} x^{\frac{8}{4}} y^{\frac{6}{4}} = 3x^2y^{\frac{3}{2}}$$

PTS: 2 REF: 012001aai NAT: N.RN.A.2 TOP: Radicals and Rational Exponents
KEY: variables

367 ANS:

$$3x^3 + x^2 + 3xy + y = x^2(3x + 1) + y(3x + 1) = (x^2 + y)(3x + 1)$$

PTS: 2 REF: 011828aai NAT: A.SSE.A.2 TOP: Factoring Polynomials
KEY: factoring by grouping

368 ANS: 3

$$95.4x - 6x^2 - (0.18x^3 + 0.02x^2 + 4x + 180)$$

PTS: 2 REF: 082322aai NAT: F.BF.A.1 TOP: Operations with Functions

369 ANS: 3

$$\sqrt{3x+18} = x \quad -3 \text{ is extraneous.}$$

$$3x+18 = x^2$$

$$x^2 - 3x - 18 = 0$$

$$(x-6)(x+3) = 0$$

$$x = 6, -3$$

PTS: 2


REF: 082315aai

NAT: A.REI.A.2

TOP: Solving Radicals

KEY: extraneous solutions

370 ANS:



PTS: 2

REF: 012032aai

NAT: F.IF.C.7

TOP: Graphing Polynomial Functions

371 ANS: 1

PTS: 2

REF: 011814aai

NAT: A.REI.D.11

TOP: Other Systems

KEY: logarithmic

Algebra II Regents at Random Worksheets

Answer Section

372 $2(0.042) = 0.084 \approx 0.08$ The percent of users making in-app purchases will be within 8% of 35%.

373 1

$$x - \frac{20}{x} = 8$$

$$x^2 - 8x - 20 = 0$$

$$(x - 10)(x + 2) = 0$$

$$x = 10, -2$$

374 1

$$1.025^{\frac{1}{12}} \approx 1.00206$$

$$375 \quad M = \frac{(152500 - 15250) \left(\frac{.036}{12} \right) \left(1 + \frac{.036}{12} \right)^{360}}{\left(1 + \frac{.036}{12} \right)^{360} - 1} \approx 624$$

376 2

$$x + y - z = 6 \quad 2x + 2y - 2z = 12 \quad 5y - 4z = 31 \quad 5y - 2(-4) = 23 \quad x + 3 - (-4) = 6$$

$$\frac{-x + 4y - z = 17}{5y - 2z = 23} \quad \frac{2x - 3y + 2z = -19}{5y - 4z = 31} \quad \frac{5y - 2z = 23}{-2z = 8} \quad 5y = 15 \quad x = -1$$

$$5y - 2z = 23 \quad 5y - 4z = 31 \quad -2z = 8 \quad y = 3$$

$$z = -4$$

$$377 \quad \frac{7}{2x} - \frac{2}{x+1} = \frac{1}{4}$$

$$\frac{7x + 7 - 4x}{2x^2 + 2x} = \frac{1}{4}$$

$$2x^2 + 2x = 12x + 28$$

$$x^2 - 5x - 14 = 0$$

$$(x - 7)(x + 2) = 0$$

$$x = 7, -2$$

378 2

379 4

$$380 \quad x^2 + (x-28)^2 = 400 \quad y = 12 - 28 = -16 \quad y = 16 - 28 = -12$$

$$x^2 + x^2 - 56x + 784 = 400$$

$$2x^2 - 56x + 384 = 0$$

$$x^2 - 28x + 192 = 0$$

$$(x-16)(x-12) = 0$$

$$x = 12, 16$$

$$381 \quad S_{10} = \frac{15 - 15(1.03)^{10}}{1 - 1.03} \approx 171.958$$

$$382 \quad 1$$

2) linear, 3) quadratic, 4) cubic

$$383 \quad 3$$

$$384 \quad D = 1.223(2.652)^A$$

$$385 \quad 2$$

$$386 \quad 4x + 6y - 8z = -2 \quad 4x + 6y - 8z = -2 \quad 4x - 8y + 20z = 12 \quad z + 2 = 3z - 4 \quad y = 3 + 2 \quad -4x + 5 + 3 = 16$$


$$4x - 8y + 20z = 12 \quad \underline{-4x + y + z = 16} \quad \underline{-4x + y + z = 16} \quad 6 = 2z \quad = 5 \quad -4x = 8$$

$$-4x + y + z = 16 \quad 7y - 7z = 14 \quad -7y + 21z = 28 \quad z = 3 \quad x = -2$$

$$y - z = 2 \quad y - 3z = -4$$

$$y = z + 2 \quad y = 3z - 4$$

$$387 \quad 3$$



$$388 \quad 2$$

$$389 \quad 4$$

$$\sqrt{3x^2y} \cdot \sqrt[3]{27x^3y^2} = 3^{\frac{1}{2}}x^{\frac{1}{2}}y^{\frac{1}{2}} \cdot 3^{\frac{2}{3}}x^{\frac{2}{3}}y^{\frac{2}{3}} = 3^{\frac{3}{2}}x^2y^{\frac{7}{6}}$$

$$390 \quad 4$$

$$a_1 = 2.5 + 0.5(1) = 3$$

391 3

$$x^2 - 4x - 5 = 4x^2 - 40x + 100$$

$$3x^2 - 36x + 105 = 0$$

$$x^2 - 12x + 35 = 0$$

$$(x - 7)(x - 5) = 0$$

$$x = 5, 7$$

392 4

$$400 \cdot .954 \approx 380$$

393 4

$$\frac{n}{m} = \frac{\sqrt{a^5}}{a} = \frac{a^{\frac{5}{2}}}{a^{\frac{2}{2}}} = a^{\frac{3}{2}} = \sqrt{a^3}$$

394 2

$$P(B) \cdot P(A|B) = P(A \text{ and } B)$$

$$P(B) \cdot 0.8 = 0.2$$

$$P(B) = 0.25$$

395 2

$$b^2 = 2b^2 - 64 \quad -8 \text{ is extraneous.}$$

$$-b^2 = -64$$

$$b = \pm 8$$

396 2

$$\frac{85}{210 + 85}$$

397 3

$$x^2 + (2x)^2 = 5 \quad y = 2x = \pm 2$$

$$x^2 + 4x^2 = 5$$

$$5x^2 = 5$$

$$x = \pm 1$$

398 2

$$(1) 0.4 \cdot 0.3 \neq 0.2, (2) 0.8 \cdot 0.25 = 0.2, (3) P(A|B) = P(A) = 0.2, (4) 0.2 \neq 0.15 \cdot 0.05$$

$$0.2 \neq 0.2 \cdot 0.2$$

399 4

$$400 \quad \frac{13.9 - 9.4}{4 - 1} = 1.5 \quad \text{The average rate of change in the number of hours of daylight from January 1-April 1 is 1.5.}$$

401 1

402 $\sqrt{6-2x} + x = 2x + 30 - 9$ $\sqrt{6-2(-29)} \neq -29 + 21$, so -29 is extraneous.

$$\sqrt{6-2x} = x + 21 \quad \sqrt{64} \neq -8$$

$$6-2x = x^2 + 42x + 441$$

$$x^2 + 44x + 435 = 0$$

$$(x+29)(x+15) = 0$$

$$x = -29, -15$$

403 1

$$\begin{array}{r} 3x-1 \\ 3x+1 \overline{) 9x^2+0x-2} \end{array}$$

$$\underline{9x^2+3x}$$

$$-3x-2$$

$$\underline{-3x-1}$$

$$-1$$

404 $2 = e^{0.0375t}$


$$t \approx 18.5$$

405 $\frac{47}{108} = \frac{1}{4} + \frac{116}{459} - P(M \text{ and } J)$; No, because $\frac{31}{459} \neq \frac{1}{4} \cdot \frac{116}{459}$

$$P(M \text{ and } J) = \frac{31}{459}$$

406 4

There is no x -intercept.



407 period = $\frac{2\pi}{0.8\pi} = 2.5$. The wheel rotates once every 2.5 seconds.
of $f(t) = 26$.

408 3

$$\frac{x^2(x+2)-9(x+2)}{x(x^2-x-6)} = \frac{(x^2-9)(x+2)}{x(x-3)(x+2)} = \frac{(x+3)(x-3)}{x(x-3)} = \frac{x+3}{x}$$

409 2



No, because the maximum

410 1

$$100\left(\frac{1}{2}\right)^{\frac{d}{8}} = 100e^{kd}$$

$$\left(\frac{1}{2}\right)^{\frac{1}{8}} = e^k$$

$$k \approx -0.087$$



411

412 3

$$-3 + 5i - (4 + 24i - 2i - 12i^2) = -3 + 5i - (16 + 22i) = -19 - 17i$$

413 1


$$(2x - i)^2 - (2x - i)(2x + 3i)$$

$$(2x - i)[(2x - i) - (2x + 3i)]$$

$$(2x - i)(-4i)$$

$$-8xi + 4i^2$$

$$-8xi - 4$$



$$y = \frac{1}{4(2)}(x - 4)^2 - 3 \quad y = \frac{-1 + -5}{2} = -3. \text{ The vertex is } (4, -3) \text{ and } p = 2.$$

415 1

The time of the next high tide will be the midpoint of consecutive low tides.

416 3

417 4

$$\frac{2}{x} = \frac{4x}{x+3}$$

$$2x+6=4x^2$$

$$4x^2-2x-6=0$$

$$2(2x^2-x-3)=0$$

$$(2x-3)(x+1)=0$$

$$x = \frac{3}{2}, -1$$

418 3

$$\frac{2}{3x+1} = \frac{1}{x} - \frac{6x}{3x+1} - \frac{1}{3} \text{ is extraneous.}$$

$$\frac{6x+2}{3x+1} = \frac{1}{x}$$

$$6x^2+2x=3x+1$$

$$6x^2-x-1=0$$

$$(2x-1)(3x+1)=0$$

$$x = \frac{1}{2}, -\frac{1}{3}$$

419 2

$$x=4y+5$$

$$x-5=4y$$

$$\frac{1}{4}x - \frac{5}{4} = y$$

420 3

$$y = 278(0.5)^{\frac{18}{1.8}} \approx 0.271$$

421 $P(-2) = 60$ $Q(-2) = 0$ $(x+2)$ is a factor of $Q(x)$ since $Q(-2) = 0$.

$$422 \quad \frac{10.1-2}{2} - \frac{2.5-0.1}{2} = 6.05 - 1.3 = 4.75$$

423 The denominator of the rational exponent represents the index of a root, and the numerator of the rational exponent represents the power of the base. $(\sqrt{9})^5 = 243$

424 2


$$P = \frac{2\pi}{\frac{\pi}{45}} = 90$$

425 3

$$\begin{array}{r}
 2x^3 - 4x^2 - x + \frac{14}{x+6} \\
 x+6 \overline{) 2x^4 + 8x^3 - 25x^2 - 6x + 14} \\
 \underline{2x^4 + 12x^3} \\
 -4x^3 - 25x^2 \\
 \underline{-4x^3 - 24x^2} \\
 -x^2 - 6x + 14 \\
 \underline{-x^2 - 6x} \\
 14
 \end{array}$$

426 4

427 2



$$428 \quad P(16) = \log(16 - 4) \approx 1.1,$$

, 14000

429 3

$$e^{bt} = \frac{c}{a}$$

$$\ln e^{bt} = \ln \frac{c}{a}$$

$$bt \ln e = \ln \frac{c}{a}$$

$$t = \frac{\ln \frac{c}{a}}{b}$$

$$\begin{array}{r}
 2a^2 + 5a + 2 \\
 3a - 2 \overline{) 6a^3 + 11a^2 - 4a - 9} \\
 \underline{6a^3 - 4a^2} \\
 15a^2 - 4a - 9 \\
 \underline{15a^2 - 10a} \\
 6a - 9 \\
 \underline{6a - 4} \\
 -5
 \end{array}$$

431 2

432 The denominator of the rational exponent represents the index of a root, and the 4th root of 81 is 3 and 3^3 is 27.

433 2

434 4

The vertex is (2,2) and $p = 3$. $3 + 2 = 5$

435 4

$$S_7 = \frac{85000 - 85000(1.06)^7}{1 - 1.06} \approx 713476.20$$

436 $20e^{.05t} = 30e^{.03t}$

$$\frac{\frac{2}{3}e^{.05t}}{e^{.05t}} = \frac{e^{.03t}}{e^{.05t}}$$

$$\ln \frac{2}{3} = \ln e^{-.02t}$$

$$\ln \frac{2}{3} = -.02t \ln e$$

$$\frac{\ln \frac{2}{3}}{-.02} = t$$

$$20.3 \approx t$$

437 $\frac{165 + 66 - 33}{825} = \frac{198}{825}$

438 1

439 $(x^2 - 6)(x^2 + 2)$

440 1

441 3

442 $N(t) = 950e^{0.0475t}$ The base is e because growth is continuous. $N\left(\frac{36}{24}\right) \approx 1020$

443 2

444 4

$$\frac{5+9}{2} = 7, \text{ vertex: } (-2, 7); p = 7 - 9 = -2, y = \frac{1}{4(-2)}(x+2)^2 + 7$$

$$y - 7 = \frac{1}{-8}(x+2)^2$$

$$-8(y - 7) = (x+2)^2$$

445 $\frac{B(11) - B(8)}{11 - 8} \approx -10.1$ The average monthly high temperature decreases 10.1° each month from August to November.

$$446 \quad \frac{p(x)}{x-1} = x^2 + 7 + \frac{5}{x-1}$$

$$p(x) = x^3 - x^2 + 7x - 7 + 5$$

$$p(x) = x^3 - x^2 + 7x - 2$$

$$447 \quad j(-1) = 2(-1)^4 - (-1)^3 - 35(-1)^2 + 16(-1) + 48 = 2 + 1 - 35 - 16 + 48 = 0; \quad x + 1 \text{ is a factor of } j(x);$$

$$2x^3 - 3x^2 - 32x + 48 = 0$$

$$x^2(2x - 3) - 16(2x - 3) = 0$$

$$(x^2 - 16)(2x - 3) = 0$$

$$x = \pm 4, \frac{3}{2}$$

$$448 \quad x + 2 \overline{) \begin{array}{r} x^3 \\ + 4 \\ \hline x^4 + 2x^3 + 4x - 10 \end{array}} \quad x^3 + 4 - \frac{18}{x+2}. \quad \text{No, because there is a remainder.}$$

$$\underline{x^4 + 2x^3}$$

$$4x - 10$$

$$\underline{4x + 8}$$

$$-18$$

$$449 \quad \frac{1}{8} + \frac{1}{6} = \frac{1}{t_b}; \quad \frac{24t_b}{8} + \frac{24t_b}{6} = \frac{24t_b}{t_b}$$

$$3t_b + 4t_b = 24$$

$$t_b = \frac{24}{7} \approx 3.4$$

$$450 \quad -6(x+3) \left(\frac{-3}{x+3} - \frac{x}{6} + 1 = 0 \right)$$

$$18 + x(x+3) - 6(x+3) = 0$$


$$18 + x^2 + 3x - 6x - 18 = 0$$

$$x^2 - 3x = 0$$

$$x(x-3) = 0$$

$$x = 0, 3$$

451 4



$$452 \quad C(t) = 63000 \left(1 + \frac{0.0255}{12} \right)^{12t} = 100000$$

$$12t \log(1.002125) = \log \frac{100}{63}$$

$$t \approx 18.14$$

453 4

$$(x-y)^2 = x^2 - 2xy + y^2 \quad (x+y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$$

454 2



455 3

$$\sqrt{x+1} = x+1$$

$$x+1 = x^2 + 2x + 1$$

$$0 = x^2 + x$$

$$0 = x(x+1)$$


$$x = -1, 0$$

456 2

457 1

$$1240(1.06)^x = 890(1.11)^x$$

$$x \approx 7$$



458

459 3

The vertex is $(-3, 5)$ and $p = 2$. $y = \frac{-1}{4(2)}(x+3)^2 + 5$

$$460 \quad (2x^2 + x - 3) \bullet (x - 1) - \left[(2x^2 + x - 3) + (x - 1) \right]$$

$$(2x^3 - 2x^2 + x^2 - x - 3x + 3) - (2x^2 + 2x - 4)$$

$$2x^3 - 3x^2 - 6x + 7$$

461 4

$$f(x) = (x+1)(x-1)(x-2) = (x^2 - 1)(x-2) = x^3 - 2x^2 - x + 2$$

462 4

$$\frac{x^2 - 4x}{2x} = \frac{x(x-4)}{2x} = \frac{x-4}{2} = \frac{x}{2} - 2 \quad \frac{x-1}{2} - \frac{3}{2} = \frac{x-1-3}{2} = \frac{x-4}{2}$$

463 4

464 4


$$0.48 \cdot 0.25 = 0.12$$

465 4

The maximum of p is 5. The minimum of f is $-\frac{21}{4}$ ($x = \frac{-6}{2(4)} = -\frac{3}{4}$)

$$f\left(-\frac{3}{4}\right) = 4\left(-\frac{3}{4}\right)^2 + 6\left(-\frac{3}{4}\right) - 3 = 4\left(\frac{9}{16}\right) - \frac{18}{4} - \frac{12}{4} = -\frac{21}{4}. \quad \frac{20}{4} - \left(-\frac{21}{4}\right) = \frac{41}{4} = 10.25$$

466 4



467 The period of P is $\frac{2}{3}$, which means the patient's blood pressure reaches a high every $\frac{2}{3}$ second and a low every $\frac{2}{3}$ second. The patient's blood pressure is high because 144 over 96 is greater than 120 over 80.

468 4

$$wx^2 + w = 0$$

$$w(x^2 + 1) = 0$$

$$x^2 = -1$$

$$x = \pm i$$

469 $0.301 \pm 2(0.058) \rightarrow 0.185 - 0.417 \frac{14}{60} \approx 0.23$. It is not unusual because 0.23 falls within this interval.

470 $\frac{103}{110 + 103} = \frac{103}{213}$

471 2

$$121(b)^2 = 64 \quad 64\left(\frac{8}{11}\right)^2 \approx 34$$

$$b = \frac{8}{11}$$

472 1

$$-4(-1) - 3 = 1 \quad 8 = \frac{2\pi}{b}$$

$$b = \frac{\pi}{4}$$

473 2

$$0.254 \pm 2(0.060) \rightarrow (0.134, 0.374)$$

474 1

$$p(x) = r(x) - c(x)$$

$$-0.5x^2 + 250x - 300 = -0.3x^2 + 150x - c(x)$$

$$c(x) = 0.2x^2 - 100x + 300$$

475 4

$$\log_2(x - 1) - 1 = 0$$

$$\log_2(x - 1) = 1$$

$$x - 1 = 2^1$$

$$x = 3$$

476 $250(1) + 2450 = 2700$ The maximum lung capacity of a person is 2700 mL.

477 2

$$n^2(n^2 - 9) + 4n(n^2 - 9) - 12(n^2 - 9)$$

$$(n^2 + 4n - 12)(n^2 - 9)$$

$$(n + 6)(n - 2)(n + 3)(n - 3)$$

478 4


$$1 + \frac{.009}{12} = 1.00075$$

479 4

$$p(5) = 2(5)^3 - 3(5) + 5 = 240$$

480 3

$$T(19) = 8 \sin(0.3(19) - 3) + 74 \approx 77$$



481 $A(t) = 318000(1.07)^t$ $318000(1.07)^t = 1000000$ The graph of $A(t)$ nearly intersects

$$1.07^t = \frac{1000}{318}$$

$$t \log 1.07 = \log \frac{1000}{318}$$

$$t = \frac{\log \frac{1000}{318}}{\log 1.07}$$

$$t \approx 17$$

the point (17, 1000000).

482 4

$$a = \frac{14-4}{2} = 5, d = \frac{14+4}{2} = 9$$

483 4

484 John found the means of the scores of the two rooms and subtracted the means. The mean score for the classical room was 7 higher than the rap room (82-75). Yes, there is less than a 5% chance this difference occurring due to random chance. It is likely the difference was due to the music.

485 q has the smaller minimum value for the domain $[-2, 2]$. h 's minimum is $-1 (2(-1) + 1)$ and q 's minimum is -8 .

486 No. $0.499 \pm 2(0.049) \rightarrow 0.401 - 0.597$. Since 0.43 falls within this interval, Robin's coin is likely not unfair.

487 2

$$u = x + 2 \quad u^2 + 4u + 3$$

$$(u + 3)(u + 1)$$

$$(x + 2 + 3)(x + 2 + 1)$$

$$(x + 5)(x + 3)$$

488 $\frac{h(2) - h(1)}{2 - 1} = -12$, $h(t) = 0$ at $t \approx 2.2, 3.8$, using a graphing calculator to find where $h(t) = 0$.

489 3

$$1^3 - k(1)^2 + 2(1) = 0$$

$$k = 3$$

490 1

$$x - \frac{4}{x-1} = 2 \quad x = \frac{3 \pm \sqrt{(-3)^2 - 4(1)(-2)}}{2(1)} = \frac{3 \pm \sqrt{17}}{2}$$

$$x(x-1) - 4 = 2(x-1)$$

$$x^2 - x - 4 = 2x - 2$$

$$x^2 - 3x - 2 = 0$$

491 4

$$1.06^{\frac{1}{52}}$$

492 4

493 1

$$x^2 + 2x + 1 = (x+1)^2$$

$$494 \quad t^2 + \left(\frac{4}{7}\right)^2 = 1 \quad -\frac{\sqrt{33}}{7}$$

$$t^2 + \frac{16}{49} = \frac{49}{49}$$

$$t^2 = \frac{33}{49}$$

$$t = \frac{\pm\sqrt{33}}{7}$$

495 3

$$\frac{x^{\frac{2}{3}} \bullet x^{\frac{5}{2}}}{x^{\frac{1}{6}}} = \frac{x^{\frac{4}{6}} \bullet x^{\frac{15}{6}}}{x^{\frac{1}{6}}} = x^{\frac{18}{6}} = x^3$$

$$496 \quad \frac{p(8) - p(4)}{8 - 4} \approx 48.78$$

497 4

498 $(a+b)^3 = a^3 + b^3$ No. Erin's shortcut only works if $a = 0$, $b = 0$ or $a = -b$.

$$a^3 + 3a^2b + 3ab^2 + b^3 = a^3 + b^3$$

$$3ab^2 + 3a^2b = 0$$

$$3ab(b+a) = 0$$

$$a = 0, b = 0, a = -b$$

499 $3\sqrt{x} - 2x = -5$ 1 is extraneous.

$$3\sqrt{x} = 2x - 5$$

$$9x = 4x^2 - 20x + 25$$

$$4x^2 - 29x + 25 = 0$$

$$(4x - 25)(x - 1) = 0$$

$$x = \frac{25}{4}, 1$$

500 2

$$y = \frac{1}{2}x + 8 \quad x = \frac{1}{2}y + 8$$

$$2x = y + 16$$

$$y = 2x - 16$$

501 3

$$440 \times 2.3\% \approx 10$$


502 3

503 4

$$m^3 - 2m^2 + 4m - 8 = 0$$

$$m^2(m - 2) + 4(m - 2) = 0$$

$$(m^2 + 4)(m - 2) = 0$$



504

505 4

(1) quadratic has two roots and both are real $(-2, 0)$ and $(-0.5, 0)$, (2) $x = \pm\sqrt{32} - 3$, (3) the real root is 3, with a multiplicity of 2, (4) $x = \pm 4i$


506 $f(x) = x^2(x + 4)(x - 3)$; $g(x) = (x + 2)^2(x + 6)(x - 1)$

507 $\frac{-1}{\sqrt{2^2 + (-1)^2}} = -\frac{1}{\sqrt{5}}$

508 4

509 $\tan \theta = \frac{\sin \theta}{\cos \theta} = \frac{-7/25}{-24/25} \quad \cos \theta = \frac{-24}{25}$


510 Self selection is a cause of bias because people with more free time are more likely to respond.



511

512 3

$$(x + 3i)^2 - (2x - 3i)^2 = x^2 + 6xi + 9i^2 - (4x^2 - 12xi + 9i^2) = -3x^2 + 18xi$$



513

514 $a_1 = 4$

$$a_n = 3a_{n-1}$$

515 1



516 $16x^4 - 81 = (4x^2 + 9)(4x^2 - 9) = (4x^2 + 9)(2x + 3)(2x - 3)$. No, because $\pm \frac{3i}{2}$ are roots.

517 1

518 2

$$\text{If } \cos \theta = \frac{7}{25}, \sin \theta = \pm \frac{24}{25}, \text{ and } \tan \theta = \frac{\sin \theta}{\cos \theta} = \frac{\pm \frac{24}{25}}{\frac{7}{25}} = \pm \frac{24}{7}$$

519 2

$$S_{20} = \frac{.01 - .01(3)^{20}}{1 - 3} = 17,433,922$$

520 4

$$(a+b+c)^2 = a^2 + ab + ac + ab + b^2 + bc + ac + ab + c^2$$

$$x = a^2 + b^2 + c^2 + 2(ab + bc + ac)$$

$$x = y + 2z$$

521 3

$$(x+4)^2 - 10 = 3x + 6 \quad y = 3(-5) + 6 = -9$$

$$x^2 + 8x + 16 - 10 = 3x + 6 \quad y = 3(0) + 6 = 6$$

$$x^2 + 5x = 0$$

$$x(x+5) = 0$$

$$x = -5, 0$$

522 3

$$\frac{c^2 - d^2}{d^2 + cd - 2c^2} = \frac{(c+d)(c-d)}{(d+2c)(d-c)} = \frac{-(c+d)}{d+2c} = \frac{-c-d}{d+2c}$$

523 4

$$1 \text{ year} = 365 \text{ days}$$

$$524 \quad P(A+B) = P(A) \cdot P(B|A) = 0.8 \cdot 0.85 = 0.68$$


525 4

$$3x - (-2x + 14) = 16 \quad 3(6) - 4z = 2$$

$$5x = 30 \quad -4z = -16$$

$$x = 6 \quad z = 4$$

526 1



527 2

$$x^2 + 4x - 1 = x - 3 \quad y + 3 = -1$$

$$x^2 + 3x + 2 = 0 \quad y = -4$$

$$(x+2)(x+1) = 0$$

$$x = -2, -1$$

528 2

$$529 \quad \frac{2x^{\frac{3}{2}}}{2x^{\frac{2}{2}}} = x^{\frac{1}{2}} = \sqrt{x}$$

530 2


531 1

$$9110 = 5000e^{30r}$$

$$\ln \frac{911}{500} = \ln e^{30r}$$

$$\frac{\ln \frac{911}{500}}{30} = r$$

$$r \approx .02$$



532

533 4


$$\begin{array}{r} 5x^2 + x - 3 \\ 2x - 1 \overline{) 10x^3 - 3x^2 - 7x + 3} \\ \underline{10x^3 - 5x^2} \\ 2x^2 - 7x \\ \underline{2x^2 - x} \\ -6x + 3 \\ \underline{-6x + 3} \end{array}$$

534 1

$$84.1\% \times 750 \approx 631$$

535 1

$$\frac{N(10) - N(1)}{10 - 1} \approx -2.03, \frac{N(20) - N(10)}{20 - 10} \approx -1.63, \frac{N(25) - N(15)}{25 - 15} \approx -1.46, \frac{N(30) - N(1)}{30 - 1} \approx -1.64$$



536

537 2

$$2 - \frac{x-1}{x+2} = 1 + \frac{x+2}{x+2} - \frac{x-1}{x+2} = 1 + \frac{x+2-(x-1)}{x+2} = 1 + \frac{3}{x+2}$$

$$538 \quad s(t) = 200(0.5)^{\frac{t}{15}} \quad \frac{1}{10} = (0.5)^{\frac{t}{15}}$$

$$\log \frac{1}{10} = \log(0.5)^{\frac{t}{15}}$$

$$-1 = \frac{t \cdot \log(0.5)}{15}$$

$$t = \frac{-15}{\log(0.5)} \approx 50$$

$$539 \quad 3$$

$$540 \quad 4$$

$$120 = 68 + (195 - 68)e^{-0.05t}$$

$$52 = 127e^{-0.05t}$$

$$\ln \frac{52}{127} = \ln e^{-0.05t}$$

$$\ln \frac{52}{127} = -0.05t$$

$$\frac{\ln \frac{52}{127}}{-0.05} = t$$

$$18 \approx t$$

$$541 \quad 4$$

$$542 \quad 138.905 \pm 2 \cdot 7.95 = 123 - 155. \text{ No, since } 125 \text{ (50\% of 250) falls within the 95\% interval.}$$


$$543 \quad 4$$

$$x(x-2) \left(\frac{10}{x^2-2x} + \frac{4}{x} = \frac{5}{x-2} \right) \quad 2 \text{ is extraneous.}$$

$$10 + 4(x-2) = 5x$$

$$10 + 4x - 8 = 5x$$

$$2 = x$$



$$544 \quad y = 2 \sin 4x$$

$$545 \quad a + 4b + 6c = 23 \quad a + 2b + c = 2 \quad 8b + 3c = 16 \quad 2b + 5(4) = 21 \quad a + 4\left(\frac{1}{2}\right) + 6(4) = 23$$

$$\underline{a + 2b + c = 2} \quad \underline{-a + 6b + 2c = 14} \quad \underline{8b + 20c = 84} \quad 2b = 1 \quad a + 2 + 24 = 23$$

$$2b + 5c = 21 \quad 8b + 3c = 16 \quad 17c = 68 \quad b = \frac{1}{2} \quad a = -3$$

$$c = 4$$

$$546 \quad B = 1.69\sqrt{30 + 4.45} - 3.49 \approx 6, \text{ which is a steady breeze.} \quad 15 = 1.69\sqrt{s + 4.45} - 3.49$$

$$18.49 = 1.69\sqrt{s + 4.45}$$

$$\frac{18.49}{1.69} = \sqrt{s + 4.45}$$

$$\left(\frac{18.49}{1.69}\right)^2 = s + 4.45$$

$$s = \left(\frac{18.49}{1.69}\right)^2 - 4.45$$

$$s \approx 115$$

$$9.5 = 1.69\sqrt{s + 4.45} - 3.49 \quad 10.49 = 1.69\sqrt{s + 4.45} - 3.49 \quad 55-64$$

$$12.99 = 1.69\sqrt{s + 4.45} \quad 13.98 = 1.69\sqrt{s + 4.45}$$

$$\frac{12.99}{1.69} = \sqrt{s + 4.45} \quad \frac{13.98}{1.69} = \sqrt{s + 4.45}$$

$$\left(\frac{12.99}{1.69}\right)^2 = s + 4.45 \quad \left(\frac{13.98}{1.69}\right)^2 = s + 4.45$$

$$s = \left(\frac{12.99}{1.69}\right)^2 - 4.45 \quad s = \left(\frac{13.98}{1.69}\right)^2 - 4.45$$

$$s \approx 55$$

$$s \approx 64$$

$$547 \quad 2$$

$$1.00643^{12} \approx 1.08$$

$$548 \quad 4$$

$$5000\left(1 + \frac{.035}{12}\right)^{12 \cdot 6} \approx 6166.50$$

549 1

$$1) (x+3)^2 - 16 = x^2 + 6x + 9 - 16 = x^2 + 6x - 7 = (x+7)(x-1); \quad 2) \quad u = x+3 \quad ; \quad 3)$$

$$u^2 - 10u - 2u + 20$$

$$u(u-10) - 2(u-10)$$

$$(u-2)(u-10)$$

$$(x+3-2)(x+3-10)$$

$$(x+1)(x-7)$$

$$\frac{(x-1)(x-7)(x+1)}{(x+1)} = (x-1)(x-7); \quad 4) \quad \frac{(x+7)(x+1)(x+3)}{(x+3)} = (x+7)(x+1)$$

550 $23 - 18 = 5$, $\bar{x} \pm 2\sigma = -3.07 - 3.13$, Yes, a difference of 5 or more occurred three times out of a thousand, which is statistically significant.

$$551 \quad \frac{9}{6} = 1.5 \quad a_1 = 6$$

$$a_n = 1.5 \cdot a_{n-1}$$

$$552 \quad 1200 \cdot 0.784 \approx 941$$

$$553 \quad 4$$

$$554 \quad 1$$

In vertex form, the parabola is $y = -\frac{1}{4(2)}(x+4)^2 + 3$. The vertex is $(-4, 3)$ and $p = 2$. $3 + 2 = 5$

555 3

$$8r^3 = 216 \quad S_{12} = \frac{8 - 8(3)^{12}}{1 - 3} = 2125760$$

$$r^3 = 27$$

$$r = 3$$

$$556 \quad a_1 = 3 \quad a_2 = 7 \quad a_3 = 15 \quad a_4 = 31; \quad \text{No, because there is no common ratio: } \frac{7}{3} \neq \frac{15}{7}$$

557 3

558 3

$$x = \frac{-2 \pm \sqrt{2^2 - 4(3)(7)}}{2(3)} = \frac{-2 \pm \sqrt{-80}}{6} = \frac{-2 \pm i\sqrt{16}\sqrt{5}}{6} = -\frac{1}{3} \pm \frac{2i\sqrt{5}}{3}$$

559 1

$$6 - (3x - 2i)(3x - 2i) = 6 - (9x^2 - 12xi + 4i^2) = 6 - 9x^2 + 12xi + 4 = -9x^2 + 12xi + 10$$

560 1

$$x^3 + 2x^2 - 9x - 18 = 0 \quad x^3 - 9x + 2x^2 - 18 = 0 \quad x^3 - 9x + 2x^2 - 18 = 0$$

$$x^2(x+2) - 9(x+2) = 0 \quad x(x^2 - 9) + 2(x^2 - 9) = 0 \quad x(x^2 - 9) + 2(x^2 - 9) = 0$$

$$(x+2)(x^2 - 9) = 0$$

561 2

$$f(x) = f(-x)$$

$$x^2 - 4 = (-x)^2 - 4$$

$$x^2 - 4 = x^2 - 4$$

$$562 \quad \frac{\sqrt[3]{x^2 y^5}}{\sqrt[4]{x^3 y^4}} = \frac{x^{\frac{2}{3}} y^{\frac{5}{3}}}{x^{\frac{3}{4}} y^{\frac{1}{2}}} = \frac{x^{\frac{8}{12}} y^{\frac{20}{12}}}{x^{\frac{9}{12}} y^{\frac{12}{12}}} = x^{-\frac{1}{12}} y^{\frac{2}{3}}$$

$$563 \quad P(F|L) = \frac{12}{27} \quad P(F) = \frac{22}{45} \quad \text{Since } P(F|L) \neq P(F), \text{ the events are not independent.}$$

564 2

$$x = -6(y - 2)$$

$$-\frac{x}{6} = y - 2$$

$$-\frac{x}{6} + 2 = y$$

$$565 \quad i^2 = -1, \text{ and not } 1; \quad 10 + 10i$$

$$566 \quad \text{No. } \left(\sqrt[7]{x^2} \right) \left(\sqrt[5]{x^3} \right) = x^{\frac{2}{7}} \cdot x^{\frac{3}{5}} = x^{\frac{31}{35}} = \sqrt[35]{x^{31}}$$

567 3

568 1

569 1

$$570 \quad 29.101 \pm 2 \cdot 0.934 = 27.23 - 30.97. \quad \text{Yes, since 30 falls within the 95\% interval.}$$

571 2

$$1) x^4 - 2x^2 y^2 + y^4 + 4x^2 y^2; \quad 3) x^4 + 2x^2 y^2 + y^4; \quad 4) 4x^4 + 4x^2 y^2 + y^4 - 3x^4 - 2x^2 y^2$$

572 1

$$-\sqrt{1 - \left(-\frac{3}{4}\right)^2} = -\sqrt{\frac{16}{16} - \frac{9}{16}} = -\sqrt{\frac{7}{16}} = -\frac{\sqrt{7}}{4}$$

$$573 \quad x^2 - 6x = -17 \quad \text{The solution is imaginary because the parabola and line do not intersect.}$$

$$x^2 - 6x + 9 = -17 + 9$$

$$(x - 3)^2 = -8$$

$$x - 3 = \pm 2i\sqrt{2}$$

$$x = 3 \pm 2i\sqrt{2}$$

574 2

$$x = \frac{2 \pm \sqrt{(-2)^2 - 4(5)(4)}}{2(5)} = \frac{2 \pm \sqrt{-76}}{10} = \frac{2 \pm i\sqrt{4}\sqrt{19}}{10} = \frac{1}{5} \pm \frac{i\sqrt{19}}{5}$$

575 3

$$1.04^{\frac{1}{12}} \approx 1.0032737$$

576 2

$$4x \bullet x^{\frac{2}{3}} + 2x^{\frac{5}{3}} = 4x^{\frac{5}{3}} + 2x^{\frac{5}{3}} = 6x^{\frac{5}{3}} = 6\sqrt[3]{x^5}$$

$$577 \quad -\frac{1}{2}i^3(3i-4) - 3i^2 = -\frac{3}{2}i^4 + 2i^3 - 3i^2 = -\frac{3}{2} - 2i + 3 = \frac{3}{2} - 2i$$

578 3

579 2

$$580 \quad x = \frac{-5 \pm \sqrt{5^2 - 4(2)(8)}}{2(2)} = -\frac{5}{4} \pm \frac{i\sqrt{39}}{4}$$

581 4

$$\ln e^{0.3x} = \ln \frac{5918}{87}$$

$$x = \frac{\ln \frac{5918}{87}}{0.3}$$

582 4

583 3

$$y = x^3 - 2$$

$$x = y^3 - 2$$

$$x + 2 = y^3$$

$$\sqrt[3]{x+2} = y$$

584 2

585 1

The cosine function has been translated +3. Since the maximum is 5 and the minimum is 1, the amplitude is 2.

$$\frac{\pi}{3} = \frac{2\pi}{b}.$$

$$b = 6$$

586 2

$$x = \frac{y}{y+2}$$

$$xy + 2x = y$$

$$xy - y = -2x$$

$$y(x-1) = -2x$$



$$y = \frac{-2x}{x-1}$$

587 1

$$(x^{\frac{3}{2}})^2 = x^3$$

588 2

589 2



590 $P(x) = R(x) - C(x) = -330x^3 + 9000x^2 - 67000x + 167000$

5 because there is a minimum in $P(x)$. Most profitable at year 13 because there is a maximum in $P(x)$.

591 4

$$(x^6y^4 - 9)(x^4 - 16)$$

$$(x^3y^2 + 3)(x^3y^2 - 3)(x^2 + 4)(x^2 - 4)$$

Least profitable at year

Algebra II Regents at Random Worksheets

Answer Section

592 ANS:

period is $\frac{2}{3}$. The wheel rotates once every $\frac{2}{3}$ second.

PTS: 2

REF: 061728aai

NAT: F.IF.C.7

TOP: Graphing Trigonometric Functions

KEY: period

593 ANS: 3

PTS: 2

REF: 061720aai

NAT: F.BF.A.2

TOP: Sequences

KEY: function notation

594 ANS:

Rewrite $\frac{4}{3}$ as $\frac{1}{3} \cdot \frac{4}{1}$, using the power of a power rule.

PTS: 2

REF: 081725aai

NAT: N.RN.A.1

TOP: Radicals and Rational Exponents

595 ANS: 4

$$k^4 - 4k^2 + 8k^3 - 32k + 12k^2 - 48$$

$$k^2(k^2 - 4) + 8k(k^2 - 4) + 12(k^2 - 4)$$

$$(k^2 - 4)(k^2 + 8k + 12)$$

$$(k + 2)(k - 2)(k + 6)(k + 2)$$

PTS: 2

REF: fall1505aai

NAT: A.SSE.A.2

TOP: Factoring Polynomials

KEY: factoring by grouping

596 ANS:

$$A = Pe^{rt}$$

$$135000 = 100000e^{5r}$$

$$1.35 = e^{5r}$$

$$\ln 1.35 = \ln e^{5r}$$

$$\ln 1.35 = 5r$$

$$.06 \approx r \text{ or } 6\%$$

PTS: 2

REF: 061632aai

NAT: F.LE.A.4

TOP: Exponential Growth

597 ANS:

$r(2) = -6$. Since there is a remainder when the cubic is divided by $x - 2$, this binomial is not a factor.

$$\begin{array}{r|rrrr} 2 & 1 & -4 & 4 & 6 \\ & & 2 & -4 & 0 \\ \hline & 1 & -2 & 0 & -6 \end{array}$$

PTS: 2

REF: 061725aai

NAT: A.APR.B.2


TOP: Remainder and Factor Theorems

598 ANS:

$$\begin{aligned} \left(\sqrt{2x-7}\right)^2 &= (5-x)^2 & \sqrt{2(4)-7}+4=5 & \sqrt{2(8)-7}+8=5 \\ 2x-7 &= 25-10x+x^2 & \sqrt{1} &= 1 & \sqrt{9} &\neq -3 \\ 0 &= x^2-12x+32 \\ 0 &= (x-8)(x-4) \\ x &= 4, 8 \end{aligned}$$

PTS: 4 REF: 081635aia NAT: A.REI.A.2 TOP: Solving Radicals
KEY: extraneous solutions

599 ANS:



PTS: 2 REF: 081732aia NAT: F.IF.C.7 TOP: Graphing Polynomial Functions

600 ANS:

No, because $P(M/R) \neq P(M)$

$$\frac{70}{180} \neq \frac{230}{490}$$

$$0.38 \neq 0.47$$

PTS: 2 REF: 011731aia NAT: S.CP.A.4 TOP: Conditional Probability

601 ANS: 1

$$d = 18; r = \pm \frac{5}{4}$$

PTS: 2 REF: 011714aia NAT: F.BF.A.2 TOP: Sequences
KEY: explicit

602 ANS: 4

If $1-i$ is one solution, the other is $1+i$. $(x-(1-i))(x-(1+i))=0$

$$x^2 - x - ix - x + ix + (1-i^2) = 0$$

$$x^2 - 2x + 2 = 0$$

PTS: 2 REF: 081601aia NAT: A.REI.B.4 TOP: Complex Conjugate Root Theorem

603 ANS:

$$\begin{array}{r}
 \overline{3x+13} \\
 x-2 \overline{) 3x^2+7x-20} \frac{6}{x-2} \\
 \underline{3x^2-6x} \\
 13x-20 \\
 \underline{13x-26} \\
 6
 \end{array}$$

PTS: 2

REF: 011732aai

NAT: A.APR.D.6

TOP: Rational Expressions

KEY: division

604 ANS: 1

$$\begin{aligned}
 \frac{2x}{x-2} \left(\frac{x}{x} \right) - \frac{11}{x} \left(\frac{x-2}{x-2} \right) &= \frac{8}{x^2-2x} \\
 2x^2 - 11x + 22 &= 8 \\
 2x^2 - 11x + 14 &= 0 \\
 (2x-7)(x-2) &= 0 \\
 x &= \frac{7}{2}, 2
 \end{aligned}$$

PTS: 2

REF: 061719aai

NAT: A.REI.A.2

TOP: Solving Rationals

605 ANS:

$$\begin{aligned}
 A &= 5000(1.045)^n & 5000 \left(1 + \frac{.046}{4} \right)^{4(6)} - 5000(1.045)^6 &\approx 6578.87 - 6511.30 \approx 67.57 & 10000 &= 5000 \left(1 + \frac{.046}{4} \right)^{4n} \\
 B &= 5000 \left(1 + \frac{.046}{4} \right)^{4n} & 2 &= 1.0115^{4n} \\
 & & \log 2 &= 4n \cdot \log 1.0115 \\
 & & n &= \frac{\log 2}{4 \log 1.0115} \\
 & & n &\approx 15.2
 \end{aligned}$$

PTS: 6

REF: 081637aai

NAT: F.LE.A.4

TOP: Exponential Growth

606 ANS: 1

$$\begin{array}{r}
 3x^2 + 4x - 1 \\
 2x + 3 \overline{) 6x^3 + 17x^2 + 10x + 2} \\
 \underline{6x^3 + 9x^2} \\
 8x^2 + 10x \\
 \underline{8x^2 + 12x} \\
 -2x + 2 \\
 \underline{-2x - 3} \\
 5
 \end{array}$$

PTS: 2 REF: fall1503aii NAT: A.APR.D.6 TOP: Rational Expressions
 KEY: division

607 ANS: 2

$$\begin{array}{r}
 x^2 + 0x + 1 \\
 x + 2 \overline{) x^3 + 2x^2 + x + 6} \\
 \underline{x^3 + 2x^2} \\
 0x^2 + x \\
 \underline{0x^2 + 0x} \\
 x + 6 \\
 \underline{x + 2} \\
 4
 \end{array}$$

PTS: 2 REF: 081611aii NAT: A.APR.D.6 TOP: Rational Expressions
 KEY: division

608 ANS: 4

The maximum volume of $p(x) = -(x+2)(x-10)(x-14)$ is about 56, at $x = 12.1$

PTS: 2 REF: 081712aii NAT: F.IF.B.4 TOP: Graphing Polynomial Functions

609 ANS: 2

$$\frac{212}{1334} \approx .16 \quad ME = \left(z \sqrt{\frac{p(1-p)}{n}} \right) = \left(1.96 \sqrt{\frac{(0.16)(0.84)}{1334}} \right) \approx 0.02 \quad \text{or} \quad \frac{1}{\sqrt{1334}} \approx .027$$

PTS: 2 REF: 081716aii NAT: S.IC.B.4 TOP: Analysis of Data

610 ANS:

Let x equal the first integer and $x + 1$ equal the next. $(x + 1)^2 - x^2 = x^2 + 2x + 1 - x^2 = 2x + 1$. $2x + 1$ is an odd integer.

PTS: 2 REF: fall1511aii NAT: A.APR.C.4 TOP: Polynomial Identities

611 ANS:

$$\sqrt[3]{x} \bullet \sqrt{x} = x^{\frac{1}{3}} \bullet x^{\frac{1}{2}} = x^{\frac{2}{6}} \bullet x^{\frac{3}{6}} = x^{\frac{5}{6}}$$

PTS: 2 REF: 061731aii NAT: N.RN.A.2 TOP: Radicals and Rational Exponents

612 ANS: 2

$$\cos \theta = \pm \sqrt{1 - \left(\frac{-\sqrt{2}}{5} \right)^2} = \pm \sqrt{\frac{25}{25} - \frac{2}{25}} = \pm \frac{\sqrt{23}}{5}$$

PTS: 2

REF: 061712aai

NAT: F.TF.C.8

TOP: Determining Trigonometric Functions

613 ANS: 1

The car lost approximately 19% of its value each year.

PTS: 2

REF: 081613aai

NAT: F.LE.B.5

TOP: Modeling Exponential Functions

614 ANS: 4

$$x(x+7) \left[\frac{3x+25}{x+7} - 5 = \frac{3}{x} \right]$$

$$x(3x+25) - 5x(x+7) = 3(x+7)$$

$$3x^2 + 25x - 5x^2 - 35x = 3x + 21$$

$$2x^2 + 13x + 21 = 0$$

$$(2x+7)(x+3) = 0$$

$$x = -\frac{7}{2}, -3$$

PTS: 2

REF: fall1501aai

NAT: A.REI.A.2

TOP: Solving Rationals

KEY: rational solutions

615 ANS: 1

1) let $y = x + 2$, then $y^2 + 2y - 8$

$$(y+4)(y-2)$$

$$(x+2+4)(x+2-2)$$

$$(x+6)x$$

PTS: 2

REF: 081715aai

NAT: A.SSE.A.2

TOP: Factoring Polynomials

KEY: multivariable

616 ANS: 2

The vertex of the parabola is (0,0). The distance, p , between the vertex and the focus or the vertex and the directrix is 1. $y = \frac{-1}{4p}(x-h)^2 + k$

$$y = \frac{-1}{4(1)}(x-0)^2 + 0$$

$$y = -\frac{1}{4}x^2$$

PTS: 2


REF: 081706aai

NAT: G.GPE.A.2

TOP: Graphing Quadratic Functions

617 ANS: 4 PTS: 2 REF: 081718aia NAT: F.IF.C.7
 TOP: Graphing Trigonometric Functions KEY: amplitude

618 ANS: 2



PTS: 2 REF: 011716aia NAT: A.REI.D.11 TOP: Other Systems
 KEY: exponential

619 ANS:

$$x = (y - 3)^3 + 1$$

$$x - 1 = (y - 3)^3$$


$$\sqrt[3]{x - 1} = y - 3$$

$$\sqrt[3]{x - 1} + 3 = y$$

$$f^{-1}(x) = \sqrt[3]{x - 1} + 3$$

PTS: 2 REF: fall1509aia NAT: F.BF.B.4 TOP: Inverse of Functions
 KEY: cubic

620 ANS:



$$A(t) = 800e^{-0.347t}$$

$$800e^{-0.347t} = 400e^{-0.231t} \quad 0.15 = e^{-0.347t}$$

$$B(t) = 400e^{-0.231t}$$



$$\ln 2e^{-0.347t} = \ln e^{-0.231t} \quad \ln 0.15 = \ln e^{-0.347t}$$

$$\ln 2 + \ln e^{-0.347t} = \ln e^{-0.231t} \quad \ln 0.15 = -0.347t \cdot \ln e$$

$$\ln 2 - 0.347t = -0.231t \quad 5.5 \approx t$$

$$\ln 2 = 0.116t$$

$$6 \approx t$$



PTS: 6

REF: 061637aai

NAT: A.REI.D.11

TOP: Other Systems

KEY: exponential

621 ANS:

$0 = \log_{10}(x - 4)$ The x -intercept of h is $(2, 0)$. f has the larger value.

$$10^0 = x - 4$$

$$1 = x - 4$$

$$x = 5$$

PTS: 2

REF: 081630aai

NAT: F.IF.C.9

TOP: Comparing Functions

622 ANS:

$$0 = \sqrt{t} - 2t + 6 \quad 2\left(\frac{9}{4}\right) - 6 < 0, \text{ so } \frac{9}{4} \text{ is extraneous.}$$

$$2t - 6 = \sqrt{t}$$

$$4t^2 - 24t + 36 = t$$

$$4t^2 - 25t + 36 = 0$$

$$(4t - 9)(t - 4) = 0$$

$$t = \frac{9}{4}, 4$$

$$(\sqrt{1} - 2(1) + 6) - (\sqrt{3} - 2(3) + 6) = 5 - \sqrt{3} \approx 3.268 \quad 327 \text{ mph}$$

PTS: 6 REF: 011737aia NAT: A.REI.A.2 TOP: Solving Radicals
KEY: context

623 ANS: 4 PTS: 2 REF: 081624aia NAT: F.BF.A.2
TOP: Sequences KEY: recursive

624 ANS: 2 PTS: 2 REF: 011701aia NAT: F.IF.B.4
TOP: Graphing Trigonometric Functions

625 ANS: 4

$$\frac{m(c)}{g(c)} = \frac{c+1}{1-c^2} = \frac{c+1}{(1+c)(1-c)} = \frac{1}{1-c}$$

PTS: 2 REF: 061608aia NAT: F.BF.A.1 TOP: Operations with Functions

626 ANS: 4

$$\frac{-3x^2 - 5x + 2}{x^3 + 2x^2} = \frac{(-3x+1)(x+2)}{x^2(x+2)} = \frac{-3x}{x^2} + \frac{1}{x^2} = -3x^{-1} + x^{-2}$$

PTS: 2 REF: 061723aia NAT: A.APR.D.6 TOP: Rational Expressions
KEY: factoring

627 ANS: 3

$$(3k - 2i)^2 = 9k^2 - 12ki + 4i^2 = 9k^2 - 12ki - 4$$

PTS: 2 REF: 081702aia NAT: N.CN.A.2 TOP: Operations with Complex Numbers
628 ANS: 2 PTS: 2 REF: 011709aia NAT: S.IC.B.5
TOP: Analysis of Data

629 ANS: 3 PTS: 2 REF: 081724aia NAT: F.BF.A.2
TOP: Sequences KEY: recursive

630 ANS: 3 PTS: 2 REF: 011710aia NAT: F.BF.A.1
TOP: Operations with Functions

631 ANS: 1

$$\begin{array}{r|rrrrrr}
 2 & 1 & 0 & -4 & -4 & 8 \\
 & & 2 & 4 & 0 & -8 \\
 \hline
 & 1 & 2 & 0 & -4 & 0
 \end{array}$$

Since there is no remainder when the quartic is divided by $x - 2$, this binomial is a factor.

PTS: 2

REF: 061711aii

NAT: A.APR.B.2

TOP: Remainder and Factor Theorems

632 ANS:

$$y = 4.168(3.981)^x \quad 100 = 4.168(3.981)^x$$

$$\log \frac{100}{4.168} = \log(3.981)^x$$

$$\log \frac{100}{4.168} = x \log(3.981)$$

$$\frac{\log \frac{100}{4.168}}{\log(3.981)} = x$$

$$x \approx 2.25$$

PTS: 4

REF: 081736aii

NAT: S.ID.B.6

TOP: Regression

KEY: exponential

633 ANS: 4

The vertex is $(2, -1)$ and $p = 2$. $y = -\frac{1}{4(2)}(x - 2)^2 - 1$

PTS: 2

REF: 081619aii

NAT: G.GPE.A.2

TOP: Graphing Quadratic Functions

634 ANS: 3

The pattern suggests an exponential pattern, not linear or sinusoidal. A 4% growth rate is accurate, while a 43% growth rate is not.

PTS: 2

REF: 011713aii

NAT: S.ID.B.6

TOP: Regression

KEY: choose model

635 ANS:

$0.506 \pm 2 \cdot 0.078 = 0.35 - 0.66$. The 32.5% value falls below the 95% confidence level.


PTS: 4

REF: 061736aii

NAT: S.IC.B.5

TOP: Analysis of Data

636 ANS:



PTS: 2

REF: 061628aai

NAT: F.IF.C.7

TOP: Graphing Trigonometric Functions

KEY: graph

637 ANS: 2

$$\begin{aligned}
 x(30 - 0.01x) - (0.15x^3 + 0.01x^2 + 2x + 120) &= 30x - 0.01x^2 - 0.15x^3 - 0.01x^2 - 2x - 120 \\
 &= -0.15x^3 - 0.02x^2 + 28x - 120
 \end{aligned}$$

PTS: 2

REF: 061709aai

NAT: F.BF.A.1

TOP: Operations with Functions

638 ANS: 4

$$496 \pm 2(115)$$

PTS: 2


REF: 011718aai

NAT: S.ID.A.4


TOP: Normal Distributions

KEY: interval

639 ANS:



The amplitude, 12, can be interpreted from the situation, since the water level has a minimum of -12 and a maximum of 12 . The value of A is -12 since at 8:30 it is low tide. The period of the function is 13 hours, and is expressed in the function through the parameter B . By experimentation with technology or using the relation $P = \frac{2\pi}{B}$ (where P is the period), it is determined that $B = \frac{2\pi}{13}$.



$$f(t) = -12 \cos\left(\frac{2\pi}{13} t\right)$$

In order to answer the question about when to fish, the student must interpret the function and determine which choice, 7:30 pm or 10:30 pm, is on an increasing interval. Since the function is increasing from $t = 13$ to $t = 19.5$ (which corresponds to 9:30 pm to 4:00 am), 10:30 is the appropriate choice.

PTS: 6

REF: spr1514aii

NAT: F.IF.C.7

TOP: Graphing Trigonometric Functions

KEY: graph

640 ANS:

$$4\% \quad 8.75 = 1.25(1+r)^{49} \text{ or } 8.75 = 1.25e^{49r}$$

$$7 = (1+r)^{49} \quad \ln 7 = \ln e^{49r}$$

$$r+1 = \sqrt[49]{7} \quad \ln 7 = 49r$$

$$r \approx .04 \quad r = \frac{\ln 7}{49}$$

$$r \approx .04$$

PTS: 2

REF: 081730aii

NAT: F.LE.A.4

TOP: Exponential Growth

641 ANS:

$$\frac{6.25 - 2.25}{21 - 5} = \frac{4}{16} = \$0.25 \text{ fine per day. } 2.25 - 5(.25) = \$1 \text{ replacement fee. } a_n = 1.25 + (n-1)(.25). \quad a_{60} = \$16$$

PTS: 4

REF: 081734aii

NAT: F.BF.A.2

TOP: Sequences

KEY: explicit

642 ANS:

$$(x^2 + y^2)^2 = (x^2 - y^2)^2 + (2xy)^2$$

$$x^4 + 2x^2y^2 + y^4 = x^4 - 2x^2y^2 + y^4 + 4x^2y^2$$

$$x^4 + 2x^2y^2 + y^4 = x^4 + 2x^2y^2 + y^4$$

PTS: 2

REF: 081727aai

NAT: A.APR.C.4

TOP: Polynomial Identities

643 ANS:

Applying the commutative property, $\left(3^{\frac{1}{5}}\right)^2$ can be rewritten as $\left(3^2\right)^{\frac{1}{5}}$ or $9^{\frac{1}{5}}$. A fractional exponent can be

rewritten as a radical with the denominator as the index, or $9^{\frac{1}{5}} = \sqrt[5]{9}$.

PTS: 2

REF: 081626aai

NAT: N.RN.A.1

TOP: Radicals and Rational Exponents

644 ANS: 1

$$8(2^{x+3}) = 48$$

$$2^{x+3} = 6$$

$$(x+3)\ln 2 = \ln 6$$

$$x+3 = \frac{\ln 6}{\ln 2}$$

$$x = \frac{\ln 6}{\ln 2} - 3$$

PTS: 2

REF: 061702aai

NAT: F.LE.A.4

TOP: Exponential Equations

KEY: without common base

645 ANS: 1

$$\frac{A}{P} = e^{rt}$$

$$0.42 = e^{rt}$$

$$\ln 0.42 = \ln e^{rt}$$

$$-0.87 \approx rt$$

PTS: 2

REF: 011723aai

NAT: F.BF.A.1

TOP: Modeling Exponential Functions

646 ANS:

The expression is of the form $y^2 - 5y - 6$ or $(y - 6)(y + 1)$. Let $y = 4x^2 + 5x$:

$$(4x^2 + 5x - 6)(4x^2 + 5x + 1)$$

$$(4x - 3)(x + 2)(4x + 1)(x + 1)$$

PTS: 2

REF: fall1512aai

NAT: A.SSE.A.2

TOP: Factoring Polynomials

KEY: a>1

647 ANS: 2

$$\begin{array}{r}
 2x^2 - 3x + 7 \\
 2x + 3 \overline{) 4x^3 + 0x^2 + 5x + 10} \\
 \underline{4x^3 + 6x^2} \\
 -6x^2 + 5x \\
 \underline{-6x^2 - 9x} \\
 14x + 10 \\
 \underline{14x + 21} \\
 -11
 \end{array}$$

PTS: 2


REF: 061614aai

NAT: A.APR.D.6

TOP: Rational Expressions

KEY: division

648 ANS:



$$y = -x + 5 \quad y = -7 + 5 = -2$$

$$(x - 3)^2 + (-x + 5 + 2)^2 = 16 \quad y = -3 + 5 = 2$$

$$x^2 - 6x + 9 + x^2 - 14x + 49 = 16$$

$$2x^2 - 20x + 42 = 0$$

$$x^2 - 10x + 21 = 0$$

$$(x - 7)(x - 3) = 0$$

$$x = 7, 3$$

PTS: 4

REF: 061633aai

NAT: A.REI.C.7

TOP: Quadratic-Linear Systems

649 ANS:

$$(4 - 3i)(5 + 2yi - 5 + 2yi)$$

$$(4 - 3i)(4yi)$$

$$16yi - 12yi^2$$

$$12y + 16yi$$

PTS: 2

REF: spr1506aai

NAT: N.CN.A.2

TOP: Operations with Complex Numbers

650 ANS:

$$20000 = PMT \left(\frac{1 - (1 + 0.00625)^{-60}}{0.00625} \right) \quad 21000 - x = 300 \left(\frac{1 - (1 + 0.00625)^{-60}}{0.00625} \right)$$

$$PMT \approx 400.76$$

$$x \approx 6028$$


PTS: 4

REF: 011736aai

NAT: F.IF.B.4

TOP: Evaluating Exponential Expressions

651 ANS: 2



PTS: 2

REF: 011712aai

NAT: A.REI.D.11

TOP: Other Systems

KEY: rational

652 ANS:

$0.602 \pm 2 \cdot 0.066 = 0.47 - 0.73$. Since 0.50 falls within the 95% interval, this supports the concern there may be an even split.

PTS: 4

REF: 061635aai

NAT: S.IC.B.5

TOP: Analysis of Data

653 ANS: 4

$$4x^2 = -98$$

$$x^2 = -\frac{98}{4}$$

$$x^2 = -\frac{49}{2}$$

$$x = \pm \sqrt{-\frac{49}{2}} = \pm \frac{7i}{\sqrt{2}} \cdot \frac{\sqrt{2}}{\sqrt{2}} = \pm \frac{7i\sqrt{2}}{2}$$

PTS: 2

REF: 061707aai

NAT: A.REI.B.4

TOP: Solving Quadratics

KEY: complex solutions | taking square roots

654 ANS: 4

PTS: 2

REF: 081708aai

NAT: A.APR.B.3

TOP: Solving Polynomial Equations

655 ANS: 4

PTS: 2

REF: 081707aai

NAT: F.TF.A.2

TOP: Reference Angles

KEY: bimodalgraph

656 ANS: 1

$$\frac{2(x-4)}{(x+3)(x-4)} + \frac{3(x+3)}{(x-4)(x+3)} = \frac{2x-2}{x^2-x-12}$$

$$2x-8+3x+9=2x-2$$

$$3x=-3$$

$$x=-1$$

PTS: 2

REF: 011717aai

NAT: A.REI.A.2

TOP: Solving Rationals

KEY: rational solutions

657 ANS: 1

(2) is not recursive

PTS: 2

REF: 081608aai

NAT: F.BF.A.2

TOP: Sequences

KEY: recursive

658 ANS:

$$\frac{1}{x} - \frac{1}{3} = -\frac{1}{3x}$$

$$\frac{3-x}{3x} = -\frac{1}{3x}$$

$$3-x=-1$$

$$x=4$$

PTS: 2

REF: 061625aai

NAT: A.REI.A.2

TOP: Solving Rationals

KEY: rational solutions

659 ANS:

$$100 = 140 \left(\frac{1}{2} \right)^{\frac{5}{h}} \quad \log \frac{100}{140} = \log \left(\frac{1}{2} \right)^{\frac{5}{h}}$$

$$40 = 140 \left(\frac{1}{2} \right)^{\frac{t}{10.3002}}$$

$$\log \frac{5}{7} = \frac{5}{h} \log \frac{1}{2}$$

$$\log \frac{2}{7} = \log \left(\frac{1}{2} \right)^{\frac{t}{10.3002}}$$

$$h = \frac{5 \log \frac{1}{2}}{\log \frac{5}{7}} \approx 10.3002$$

$$\log \frac{2}{7} = \frac{t \log \left(\frac{1}{2} \right)}{10.3002}$$

$$t = \frac{10.3002 \log \frac{2}{7}}{\log \frac{1}{2}} \approx 18.6$$

PTS: 6

REF: 061737aai

NAT: F.LE.A.4

TOP: Exponential Decay

660 ANS: 2

$$\sqrt{x+14} = \sqrt{2x+5} + 1 \qquad \sqrt{22+14} - \sqrt{2(22)+5} = 1$$

$$x+14 = 2x+5+2\sqrt{2x+5}+1 \qquad 6-7 \neq 1$$

$$-x+8 = 2\sqrt{2x+5}$$

$$x^2 - 16x + 64 = 8x + 20$$

$$x^2 - 24x + 44 = 0$$

$$(x-22)(x-2) = 0$$

$$x = 2, 22$$

PTS: 2

REF: 081704aai

NAT: A.REI.A.2

TOP: Solving Radicals

KEY: advanced

661 ANS: 3



PTS: 2

REF: 081604aai

NAT: S.ID.A.4

TOP: Normal Distributions

KEY: probability

662 ANS: 3

PTS: 2

REF: 061623aai

NAT: F.BF.A.2

TOP: Sequences

KEY: recursive

663 ANS: 4

PTS: 2

REF: 061601aai

NAT: N.RN.A.2

TOP: Radicals and Rational Exponents

KEY: variables

664 ANS:

$A(t) = 100(0.5)^{\frac{t}{63}}$, where t is time in years, and $A(t)$ is the amount of titanium-44 left after t years.

$\frac{A(10) - A(0)}{10 - 0} = \frac{89.58132 - 100}{10} = -1.041868$ The estimated mass at $t = 40$ is $100 - 40(-1.041868) \approx 58.3$. The

actual mass is $A(40) = 100(0.5)^{\frac{40}{63}} \approx 64.3976$. The estimated mass is less than the actual mass.

PTS: 6

REF: fall1517aai

NAT: F.LE.A.2

TOP: Modeling Exponential Functions

665 ANS:

Randomly assign participants to two groups. One group uses the toothpaste with ingredient X and the other group uses the toothpaste without ingredient X .

PTS: 2

REF: 061626aai

NAT: S.IC.B.3

TOP: Analysis of Data

666 ANS: 3

(3) repeats 3 times over 2π .

PTS: 2

REF: 011722aai

NAT: F.IF.C.7

TOP: Graphing Trigonometric Functions

KEY: recognize | bimodalgraph

667 ANS: 1 PTS: 2 REF: 081609aia NAT: F.BF.B.6
TOP: Sigma Notation KEY: represent

668 ANS:

$$P(W/D) = \frac{P(W \cap D)}{P(D)} = \frac{.4}{.5} = .8$$

PTS: 2 REF: 081726aia NAT: S.CP.A.3 TOP: Conditional Probability
669 ANS:

$$\frac{3p}{p-5} = \frac{p+2}{p+3}$$

$$3p^2 + 9p = p^2 - 3p - 10$$

$$2p^2 + 12p + 10 = 0$$

$$p^2 + 6p + 5 = 0$$

$$(p+5)(p+1) = 0$$


$$p = -5, -1$$

PTS: 4 REF: 081733aia NAT: A.REI.A.2 TOP: Solving Rationals
KEY: rational solutions
670 ANS:

$\csc \theta = \frac{1}{\sin \theta}$, and $\sin \theta$ on a unit circle represents the y value of a point on the unit circle. Since $y = \sin \theta$,

$$\csc \theta = \frac{1}{y}.$$

PTS: 2 REF: 011727aia NAT: F.TF.A.2 TOP: Reciprocal Trigonometric Relationships
671 ANS: 3



Define $r(a) = \frac{1}{0.0105 \cdot \ln\left(\frac{a}{5000}\right)}$ Done

$\frac{r(8000) - r(6000)}{8000 - 6000} = 0.013699$

$\frac{r(12000) - r(9000)}{12000 - 9000} = 0.009133$

PTS: 2 REF: 081922aia NAT: F.IF.B.6 TOP: Rate of Change
KEY: logarithmic

672 ANS:

$$\left(x^{\frac{5}{3}}\right)^{\frac{6}{5}} = \left(y^{\frac{5}{6}}\right)^{\frac{6}{5}}$$


$$x^2 = y$$

PTS: 2 REF: 011730aia NAT: N.RN.A.2 TOP: Radicals and Rational Exponents
KEY: variables

673 ANS: 3 PTS: 2 REF: 011708aia NAT: F.BF.B.5
TOP: Inverse of Functions KEY: exponential

674 ANS: 1 PTS: 2 REF: 011704aia NAT: F.TF.C.8
TOP: Proving Trigonometric Identities KEY: basic

675 ANS:



PTS: 2 REF: 061729aia NAT: F.IF.C.7 TOP: Graphing Exponential Functions
676 ANS: 4 PTS: 2 REF: 081622aia NAT: F.BF.A.1
TOP: Modeling Exponential Functions

677 ANS:

$$xi(-6i)^2 = xi(36i^2) = 36xi^3 = -36xi$$


PTS: 2 REF: 081627aia NAT: N.CN.A.2 TOP: Operations with Complex Numbers
678 ANS:

$$\frac{f(4) - f(-2)}{4 - (-2)} = \frac{80 - 1.25}{6} = 13.125 \quad g(x) \text{ has a greater rate of change}$$

$$\frac{g(4) - g(-2)}{4 - (-2)} = \frac{179 - (-49)}{6} = 38$$

PTS: 4 REF: 061636aia NAT: F.IF.B.6 TOP: Rate of Change
KEY: exponential
679 ANS: 1 PTS: 2 REF: 081722aia NAT: S.IC.B.6
TOP: Analysis of Data KEY: draw conclusions

680 ANS: 4



A parabola with a focus of (0,4) and a directrix of $y = 2$ is sketched as follows: By inspection, it is determined that the vertex of the parabola is (0,3). It is also evident that the distance, p , between the vertex and the focus is 1. It is possible to use the formula $(x - h)^2 = 4p(y - k)$ to derive the equation of the parabola as follows: $(x - 0)^2 = 4(1)(y - 3)$

$$x^2 = 4y - 12$$

$$x^2 + 12 = 4y$$

$$\frac{x^2}{4} + 3 = y$$

or A point (x, y) on the parabola must be the same distance from the focus as it is from the directrix. For any such point (x, y) , the distance to the focus is $\sqrt{(x - 0)^2 + (y - 4)^2}$ and the distance to the directrix is $y - 2$. Setting this equal leads to: $x^2 + y^2 - 8y + 16 = y^2 - 4y + 4$

$$x^2 + 16 = 4y + 4$$

$$\frac{x^2}{4} + 3 = y$$

PTS: 2 REF: spr1502aii NAT: G.GPE.A.2 TOP: Graphing Quadratic Functions
681 ANS: 3 PTS: 2 REF: 061607aii NAT: S.IC.A.2
TOP: Analysis of Data

682 ANS:

$$\sin^2 \theta + (-0.7)^2 = 1 \quad \text{Since } \theta \text{ is in Quadrant II, } \sin \theta = \sqrt{.51} \text{ and } \tan \theta = \frac{\sin \theta}{\cos \theta} = \frac{\sqrt{.51}}{-0.7} \approx -1.02$$

$$\sin^2 \theta = .51$$


$$\sin \theta = \pm \sqrt{.51}$$

PTS: 2 REF: 081628aii NAT: F.TF.C.8 TOP: Determining Trigonometric Functions
683 ANS:

sample: pails of oranges; population: truckload of oranges. It is likely that about 5% of all the oranges are unsatisfactory.

PTS: 2 REF: 011726aii NAT: S.IC.B.6 TOP: Analysis of Data
KEY: draw conclusions

684 ANS:



$$0 = x^2(x+1) - 4(x+1)$$

$$0 = (x^2 - 4)(x+1)$$

$$0 = (x+2)(x-2)(x+1)$$

$$x = -2, -1, 2$$


PTS: 4

REF: 081633aai

NAT: F.IF.C.7

TOP: Graphing Polynomial Functions

685 ANS:



$$-2x + 1 = -2x^2 + 3x + 1$$

$$2x^2 - 5x = 0$$

$$x(2x - 5) = 0$$

$$x = 0, \frac{5}{2}$$


PTS: 2

REF: fall1507aai

NAT: A.REI.C.7

TOP: Quadratic-Linear Systems

686 ANS: 4



As the range is $[4, 10]$, the midline is $y = \frac{4+10}{2} = 7$.

PTS: 2

REF: fall1506aai

NAT: F.IF.C.7

TOP: Graphing Trigonometric Functions

KEY: mixed

687 ANS: 4

PTS: 2


REF: 061716aai

NAT: N.RN.A.2

TOP: Radicals and Rational Exponents

KEY: variables

688 ANS: 4



$$y = 5^{-x} = \left(\frac{1}{5}\right)^x$$


PTS: 2

REF: 061615aai

NAT: F.IF.C.7

TOP: Graphing Exponential Functions

689 ANS:



$$100 = 325 + (68 - 325)e^{-2k} \quad T = 325 - 257e^{-0.066t}$$

$$-225 = -257e^{-2k} \quad T = 325 - 257e^{-0.066(7)} \approx 163$$

$$k = \frac{\ln\left(\frac{-225}{-257}\right)}{-2}$$

$$k \approx 0.066$$

PTS: 4

REF: fall1513aai

NAT: F.LE.A.4

TOP: Exponential Growth

690 ANS: 4

$$y = g(x) = (x - 2)^2 \quad (x - 2)^2 = 3x - 2 \quad y = 3(6) - 2 = 16$$

$$x^2 - 4x + 4 = 3x - 2 \quad y = 3(1) - 2 = 1$$

$$x^2 - 7x + 6 = 0$$

$$(x - 6)(x - 1) = 0$$

$$x = 6, 1$$

PTS: 2

REF: 011705aai

NAT: A.REI.C.7

TOP: Quadratic-Linear Systems

691 ANS: 4

	Bar Harbor	Phoenix
Minimum	31.386	66.491
Midline	55.3	86.729
Maximum	79.214	106.967
Range	47.828	40.476

PTS: 2

REF: 061715aai

NAT: F.IF.B.4

TOP: Graphing Trigonometric Functions

KEY: maximum/minimum

692 ANS: 3

PTS: 2

REF: 061710aai

NAT: S.IC.A.2

TOP: Analysis of Data

693 ANS:

The mean difference between the students' final grades in group 1 and group 2 is -3.64 . This value indicates that students who met with a tutor had a mean final grade of 3.64 points less than students who used an on-line subscription. One can infer whether this difference is due to the differences in intervention or due to which students were assigned to each group by using a simulation to rerandomize the students' final grades many (500) times. If the observed difference -3.64 is the result of the assignment of students to groups alone, then a difference of -3.64 or less should be observed fairly regularly in the simulation output. However, a difference of -3 or less occurs in only about 2% of the rerandomizations. Therefore, it is quite unlikely that the assignment to groups alone accounts for the difference; rather, it is likely that the difference between the interventions themselves accounts for the difference between the two groups' mean final grades.


PTS: 4

REF: fall1514aii

NAT: S.IC.B.5

TOP: Analysis of Data

694 ANS: 1



PTS: 2

REF: 061617aii

NAT: F.TF.A.2

TOP: Determining Trigonometric Functions

KEY: extension to reals

695 ANS:

$$a_1 = 4 \quad a_8 = 639$$

$$a_n = 2a_{n-1} + 1$$

PTS: 2

REF: 081729aii

NAT: F.BF.A.2

TOP: Sequences

KEY: recursive

696 ANS: 1

$$P(28) = 5(2)^{\frac{98}{28}} \approx 56$$

PTS: 2

REF: 011702aii

NAT: F.LE.A.2

TOP: Modeling Exponential Functions

697 ANS:

$$P(S \cap M) = P(S) + P(M) - P(S \cup M) = \frac{649}{1376} + \frac{433}{1376} - \frac{974}{1376} = \frac{108}{1376}$$

PTS: 2

REF: 061629aii

NAT: S.CP.B.7

TOP: Addition Rule

698 ANS:

$$P(A \cup B) = P(A) + P(B) - P(A \cap B) \quad A \text{ and } B \text{ are independent since } P(A \cap B) = P(A) \cdot P(B)$$

$$0.8 = 0.6 + 0.5 - P(A \cap B)$$

$$0.3 = 0.6 \cdot 0.5$$

$$P(A \cap B) = 0.3$$

$$0.3 = 0.3$$

PTS: 2

REF: 081632aii

NAT: S.CP.B.7

TOP: Addition Rule

699 ANS: 3

$$-33t^2 + 360t = 700 + 5t$$

$$-33t^2 + 355t - 700 = 0$$

$$t = \frac{-355 \pm \sqrt{355^2 - 4(-33)(-700)}}{2(-33)} \approx 3,8$$

PTS: 2 REF: 081606aia NAT: A.REI.D.11 TOP: Quadratic-Linear Systems

700 ANS:

$$S_n = \frac{33000 - 33000(1.04)^n}{1 - 1.04} \quad S_{15} = \frac{33000 - 33000(1.04)^{15}}{1 - 1.04} \approx 660778.39$$

PTS: 4 REF: 061634aia NAT: F.BF.B.7 TOP: Series

KEY: geometric

701 ANS: 3

Self selection causes bias.

PTS: 2 REF: 061703aia NAT: S.IC.B.6 TOP: Analysis of Data

KEY: bias

702 ANS: 4

$$x = \frac{8 \pm \sqrt{(-8)^2 - 4(6)(29)}}{2(6)} = \frac{8 \pm \sqrt{-632}}{12} = \frac{8 \pm i\sqrt{4}\sqrt{158}}{12} = \frac{2}{3} \pm \frac{1}{6}i\sqrt{158}$$

PTS: 2 REF: 011711aia NAT: A.REI.B.4 TOP: Solving Quadratics

KEY: complex solutions | quadratic formula

703 ANS: 3 PTS: 2 REF: 061722aia NAT: A.CED.A.1

TOP: Modeling Rationals

704 ANS:

$$M = 172600 \bullet \frac{0.00305(1 + 0.00305)^{12 \cdot 15}}{(1 + 0.00305)^{12 \cdot 15} - 1} \approx 1247 \quad 1100 = (172600 - x) \bullet \frac{0.00305(1 + 0.00305)^{12 \cdot 15}}{(1 + 0.00305)^{12 \cdot 15} - 1}$$

$$1100 \approx (172600 - x) \bullet (0.007228)$$

$$152193 \approx 172600 - x$$

$$20407 \approx x$$

PTS: 4 REF: 061734aia NAT: F.IF.B.4 TOP: Evaluating Exponential Expressions

705 ANS:

$$x^2(4x - 1) + 4(4x - 1) = (x^2 + 4)(4x - 1)$$

PTS: 2 REF: 061727aia NAT: A.SSE.A.2 TOP: Factoring Polynomials

KEY: factoring by grouping

706 ANS: 3

$$\frac{f(7)-f(-7)}{7-(-7)} = \frac{2^{-0.25(7)} \bullet \sin\left(\frac{\pi}{2}(7)\right) - 2^{-0.25(-7)} \bullet \sin\left(\frac{\pi}{2}(-7)\right)}{14} \approx -0.26$$

PTS: 2 REF: 061721aii NAT: F.IF.B.6 TOP: Rate of Change
KEY: trigonometric

707 ANS: 1

$$\frac{157}{25+47+157}$$

PTS: 2 REF: 081607aii NAT: S.CP.A.4 TOP: Conditional Probability

708 ANS: 4

$$\left(\frac{-54x^9}{y^4}\right)^{\frac{2}{3}} = \frac{(2 \cdot -27)^{\frac{2}{3}} x^{\frac{18}{3}}}{y^{\frac{8}{3}}} = \frac{2^{\frac{2}{3}} \cdot 9x^6}{y^2 \cdot y^{\frac{2}{3}}} = \frac{9x^6 \sqrt[3]{4}}{y^2 \sqrt[3]{y^2}}$$

PTS: 2 REF: 081723aii NAT: N.RN.A.2 TOP: Radicals and Rational Exponents
KEY: variables

709 ANS: 2

$$\begin{array}{r|rrrr} -4 & 1 & -11 & 16 & 84 \\ & & -4 & 60 & -304 \\ \hline & 1 & -15 & 76 & \end{array}$$

Since there is a remainder when the cubic is divided by $x + 4$, this binomial is not a factor.

PTS: 2 REF: 081720aii NAT: A.APR.B.2 TOP: Remainder and Factor Theorems

710 ANS:



69

PTS: 2 REF: 061726aii NAT: S.ID.A.4 TOP: Normal Distributions
KEY: percent

711 ANS: 3

$$2d(d^3 + 3d^2 - 9d - 27)$$

$$2d(d^2(d + 3) - 9(d + 3))$$

$$2d(d^2 - 9)(d + 3)$$

$$2d(d + 3)(d - 3)(d + 3)$$

$$2d(d + 3)^2(d - 3)$$

PTS: 2 REF: 081615aai NAT: A.SSE.A.2 TOP: Factoring Polynomials
KEY: factoring by grouping

712 ANS: 4

$$m^5 + m^3 - 6m = m(m^4 + m^2 - 6) = m(m^2 + 3)(m^2 - 2)$$

PTS: 2 REF: 011703aai NAT: A.SSE.A.2 TOP: Factoring Polynomials
KEY: higher power

713 ANS: 3 PTS: 2 REF: 011706aai NAT: S.IC.B.3
TOP: Analysis of Data

714 ANS:

$$0 = 6(-5)^3 + b(-5)^2 - 52(-5) + 15 \quad z(x) = 6x^3 + 19x^2 - 52x + 15$$

$$0 = -750 + 25b + 260 + 15$$

$$475 = 25b$$

$$19 = b$$

-5	6	19	-52	15
		-30	55	15
	6	-11	3	0

$$6x^2 - 11x + 3 = 0$$

$$(2x - 3)(3x - 1) = 0$$

$$x = \frac{3}{2}, \frac{1}{3}, -5$$

PTS: 4 REF: fall1515aai NAT: A.APR.B.3 TOP: Solving Polynomial Equations

715 ANS: 2

$$x = -\frac{3}{4}y + 2$$


$$-4x = 3y - 8$$

$$-4x + 8 = 3y$$

$$-\frac{4}{3}x + \frac{8}{3} = y$$

PTS: 2 REF: 061616aai NAT: F.BF.B.4 TOP: Inverse of Functions
KEY: linear

716 ANS:



$$720 = \frac{120000 \left(\frac{.048}{12} \right) \left(1 + \frac{.048}{12} \right)^n}{\left(1 + \frac{.048}{12} \right)^n - 1} \quad \frac{275.2}{12} \approx 23 \text{ years}$$

$$720(1.004)^n - 720 = 480(1.004)^n$$

$$240(1.004)^n = 720$$

$$1.004^n = 3$$

$$n \log 1.004 = \log 3$$

$$n \approx 275.2 \text{ months}$$

PTS: 4

REF: spr1509aii

NAT: F.LE.A.4

TOP: Exponential Growth

717 ANS: 3

$$1.0525^{\frac{1}{12}} \approx 1.00427$$



PTS: 2

REF: 061621aii

NAT: A.SSE.B.3

TOP: Modeling Exponential Functions

718 ANS:



At 1.95 years, the value of the car equals the loan balance. Zach can cancel the policy after 6 years.

PTS: 4

REF: 081737aii

NAT: A.REI.D.11

TOP: Other Systems

KEY: exponential

719 ANS:

$$7 = 20(0.5)^{\frac{t}{8.02}}$$

$$\log 0.35 = \log 0.5^{\frac{t}{8.02}}$$

$$\log 0.35 = \frac{t \log 0.5}{8.02}$$

$$\frac{8.02 \log 0.35}{\log 0.5} = t$$

$$t \approx 12$$


PTS: 4

REF: 081634aai

NAT: F.LE.A.4

TOP: Exponential Decay

720 ANS: 3



$H(t)$ is at a minimum at $70(-1) + 80 = 10$

PTS: 2

REF: 061613aai

NAT: F.IF.B.4

TOP: Graphing Trigonometric Functions

KEY: maximum/minimum

721 ANS:

Using a 95% level of confidence, $x \pm 2$ standard deviations sets the usual wait time as 150-302 seconds. 360 seconds is unusual.

PTS: 2

REF: 081629aai

NAT: S.IC.B.6

TOP: Analysis of Data

KEY: draw conclusions

722 ANS: 2

The events are independent because $P(A \text{ and } B) = P(A) \cdot P(B)$.

$$0.125 = 0.5 \cdot 0.25$$

If $P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B) = 0.25 + 0.5 - .125 = 0.625$, then the events are not mutually exclusive because $P(A \text{ or } B) = P(A) + P(B)$

$$0.625 \neq 0.5 + 0.25$$


PTS: 2

REF: 061714aai

NAT: S.CP.A.3

TOP: Conditional Probability

723 ANS:



$$\begin{array}{r}
 2x^2 + 6x + 23 \\
 x - 5 \overline{) 2x^3 - 4x^2 - 7x - 10} \\
 \underline{2x^3 - 10x^2} \\
 6x^2 - 7x \\
 \underline{6x^2 - 30x} \\
 23x - 10 \\
 \underline{23x - 115} \\
 105
 \end{array}$$

Since there is a remainder, $x - 5$ is not a factor.

PTS: 2 REF: 061627aia NAT: A.APR.B.2 TOP: Remainder and Factor Theorems

724 ANS:

$$\sqrt{x-4} = -x+6 \quad \sqrt{x-4} = -8+6 = -2 \text{ is extraneous.}$$

$$x-4 = x^2 - 12x + 36$$

$$0 = x^2 - 13x + 40$$


$$0 = (x-8)(x-5)$$

$$x = 5, 8$$

PTS: 2 REF: 061730aia NAT: A.REI.A.2 TOP: Solving Radicals

KEY: extraneous solutions


725 ANS: 1



PTS: 2 REF: 061618aia NAT: F.IF.C.7 TOP: Graphing Logarithmic Functions

726 ANS: 1

A reference triangle can be sketched using the coordinates $(-4, 3)$ in the second quadrant to find the value of $\sin \theta$.



PTS: 2


REF: spr1503aaii

NAT: F.TF.A.2

TOP: Determining Trigonometric Functions

KEY: extension to reals

727 ANS:




PTS: 2

REF: 011729aaii

NAT: F.IF.C.7

TOP: Graphing Polynomial Functions

728 ANS: 1



$$x^4 - 4x^3 - 9x^2 + 36x = 0$$

$$x^3(x - 4) - 9x(x - 4) = 0$$

$$(x^3 - 9x)(x - 4) = 0$$

$$x(x^2 - 9)(x - 4) = 0$$

$$x(x + 3)(x - 3)(x - 4) = 0$$

$$x = 0, \pm 3, 4$$

PTS: 2

REF: 061606aaii

NAT: A.APR.B.3

TOP: Solving Polynomial Equations

729 ANS:

Based on these data, the two events do not appear to be independent. $P(F) = \frac{106}{200} = 0.53$, while

$P(F|T) = \frac{54}{90} = 0.6$, $P(F|R) = \frac{25}{65} = 0.39$, and $P(F|C) = \frac{27}{45} = 0.6$. The probability of being female are not the same as the conditional probabilities. This suggests that the events are not independent.

PTS: 2

REF: fall1508aaii

NAT: S.CP.A.4

TOP: Conditional Probability

730 ANS: 3

The graph shows three real zeros, and has end behavior matching the given end behavior.

PTS: 2 REF: 061604aai NAT: F.IF.B.4 TOP: Graphing Polynomial Functions
KEY: bimodalgraph

731 ANS:

$j(-x) = (-x)^4 - 3(-x)^2 - 4 = x^2 - 3x^2 - 4$ Since $j(x) = j(-x)$, the function is even.

PTS: 2 REF: 081731aai NAT: F.BF.B.3 TOP: Even and Odd Functions

732 ANS:

$$\frac{x^3 + 9}{x^3 + 8} = \frac{x^3 + 8}{x^3 + 8} + \frac{1}{x^3 + 8}$$

$$\frac{x^3 + 9}{x^3 + 8} = \frac{x^3 + 9}{x^3 + 8}$$

PTS: 2 REF: 061631aai NAT: A.APR.D.7 TOP: Addition and Subtraction of Rationals

733 ANS: 4

$$(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3 \neq x^3 + 3xy + y^3$$

PTS: 2 REF: 081620aai NAT: A.APR.C.4 TOP: Polynomial Identities

734 ANS:

$$(1 - i)(1 - i)(1 - i) = (1 - 2i + i^2)(1 - i) = -2i(1 - i) = -2i + 2i^2 = -2 - 2i$$

PTS: 2 REF: 011725aai NAT: N.CN.A.2 TOP: Operations with Complex Numbers

735 ANS: 4 PTS: 2 REF: 061706aai NAT: F.IF.B.4

TOP: Graphing Trigonometric Functions

736 ANS:

Some of the students who did not drink energy drinks read faster than those who did drink energy drinks.

$17.7 - 19.1 = -1.4$ Differences of -1.4 and less occur $\frac{25}{232}$ or about 10% of the time, so the difference is not unusual.

PTS: 4 REF: 081636aai NAT: S.IC.B.5 TOP: Analysis of Data

737 ANS: 4

$$(1) \frac{B(60) - B(10)}{60 - 10} \approx 28\% \quad (2) \frac{B(69) - B(19)}{69 - 19} \approx 33\% \quad (3) \frac{B(72) - B(36)}{72 - 36} \approx 38\% \quad (4) \frac{B(73) - B(60)}{73 - 60} \approx 46\%$$

PTS: 2 REF: 011721aai NAT: F.IF.B.6 TOP: Rate of Change

KEY: exponential

738 ANS:

A student is more likely to jog if both siblings jog. 1 jogs: $\frac{416}{2239} \approx 0.19$. both jog: $\frac{400}{1780} \approx 0.22$

PTS: 2 REF: 061732aai NAT: S.CP.A.4 TOP: Conditional Probability

739 ANS: 4

The scenario represents a decreasing geometric sequence with a common ratio of 0.80.

PTS: 2

REF: 061610aai

NAT: F.BF.A.2

TOP: Sequences

KEY: recursive

740 ANS:

$$6x - 3y + 2z = -10 \quad x + 3y + 5z = 45 \quad 4x + 10z = 62 \quad 4x + 4(7) = 20 \quad 6(-2) - 3y + 2(7) = -10$$

$$-2x + 3y + 8z = 72 \quad 6x - 3y + 2z = -10 \quad 4x + 4z = 20 \quad 4x = -8 \quad -3y = -12$$

$$4x + 10z = 62 \quad 7x + 7z = 35 \quad 6z = 42 \quad x = -2 \quad y = 4$$

$$4x + 4z = 20 \quad z = 7$$

PTS: 4

REF: spr1510aai

NAT: A.REI.C.6

TOP: Solving Linear Systems

KEY: three variables

741 ANS: 2

 $\bar{x} + 2\sigma$ represents approximately 48% of the data.

PTS: 2

REF: 061609aai

NAT: S.ID.A.4

TOP: Normal Distributions

KEY: percent

742 ANS:

$$\frac{306.25 - 156.25}{70 - 50} = \frac{150}{20} = 7.5 \quad \text{Between 50-70 mph, each additional mph in speed requires 7.5 more feet to stop.}$$

PTS: 2


REF: 081631aai

NAT: F.IF.B.6

TOP: Rate of Change

KEY: table

743 ANS:

Part a sketch is shifted $\frac{\pi}{3}$ units right.

PTS: 4

REF: 081735aai

NAT: F.IF.C.7

TOP: Graphing Trigonometric Functions

KEY: graph

744 ANS: 2

$$6xi^3(-4xi+5) = -24x^2i^4 + 30xi^3 = -24x^2(1) + 30x(-1) = -24x^2 - 30xi$$

PTS: 2

REF: 061704aai

NAT: N.CN.A.2

TOP: Operations with Complex Numbers

745 ANS: 2


PTS: 2

REF: 011720aai

NAT: A.APR.B.2

TOP: Remainder and Factor Theorems

746 ANS:

As $x \rightarrow -3$, $y \rightarrow -\infty$. As $x \rightarrow \infty$, $y \rightarrow \infty$.


PTS: 4

REF: 061735aai

NAT: F.IF.C.7

TOP: Graphing Logarithmic Functions

747 ANS:



The vertex of the parabola is $(4, -3)$. The x -coordinate of the focus and the vertex is the same. Since the distance from the vertex to the directrix is 3, the distance from the vertex to the focus is 3, so the y -coordinate of the focus is 0. The coordinates of the focus are $(4, 0)$.

PTS: 2

REF: 061630aai

NAT: G.GPE.A.2

TOP: Graphing Quadratic Functions

748 ANS: 1

PTS: 2

REF: 081616aai

NAT: F.TF.A.1

TOP: Unit Circle

KEY: bimodalgraph

749 ANS: 4

$$S_n = \frac{32 - 32(.8)^{12}}{1 - .8} \approx 149$$

PTS: 2

REF: 081721aai

NAT: F.BF.B.7

TOP: Series

KEY: geometric

750 ANS: 1

The zeros of the polynomial are at $-b$, and c . The sketch of a polynomial of degree 3 with a negative leading coefficient should have end behavior showing as x goes to negative infinity, $f(x)$ goes to positive infinity. The multiplicities of the roots are correctly represented in the graph.

PTS: 2

REF: spr1501aai

NAT: A.APR.B.3

TOP: Graphing Polynomial Functions

KEY: bimodalgraph

751 ANS: 1

$$x = \frac{-3 \pm \sqrt{3^2 - 4(2)(2)}}{2(2)} = \frac{-3 \pm \sqrt{-7}}{4} = -\frac{3}{4} \pm \frac{i\sqrt{7}}{4}$$

PTS: 2 REF: 061612aia NAT: A.REI.B.4 TOP: Solving Quadratics

KEY: complex solutions | quadratic formula


752 ANS: 1

II. Ninth graders drive to school less often; III. Students know little about adults; IV. Calculus students love math!

PTS: 2 REF: 081602aia NAT: S.IC.B.6 TOP: Analysis of Data

KEY: bias

753 ANS:

 $f(4) = 2(4)^3 - 5(4)^2 - 11(4) - 4 = 128 - 80 - 44 - 4 = 0$ Any method that demonstrates 4 is a zero of $f(x)$ confirms
that $x - 4$ is a factor, as suggested by the Remainder Theorem.

PTS: 2 REF: spr1507aia NAT: A.APR.B.2 TOP: Remainder and Factor Theorems

754 ANS:

$$\frac{x^{\frac{8}{3}}}{x^{\frac{4}{3}}} = x^y$$

$$x^{\frac{4}{3}} = x^y$$

$$\frac{4}{3} = y$$

PTS: 2 REF: spr1505aia NAT: N.RN.A.2 TOP: Radicals and Rational Exponents

KEY: numbers

755 ANS: 1

$$\begin{array}{r}
 2x^2 + x + 5 \\
 2x - 1 \overline{) 4x^3 + 0x^2 + 9x - 5} \\
 \underline{4x^3 - 2x^2} \\
 2x^2 + 9x \\
 \underline{2x^2 - x} \\
 10x - 5 \\
 \underline{10x - 5} \\
 0
 \end{array}$$

PTS: 2 REF: 081713aai NAT: A.APR.D.6 TOP: Rational Expressions
KEY: division

756 ANS: 1

The probability of rain equals the probability of rain, given that Sean pitches.

PTS: 2 REF: 061611aai NAT: S.CP.A.3 TOP: Conditional Probability

757 ANS:

$$\begin{array}{rclclcl}
 x + y + z = 1 & x + y + z = 1 & x + y + z = 1 & -2z - z = 3 & y - (-1) = 3 & x + 2 - 1 = 1 \\
 x + 2y + 3z = 1 & \underline{x + 2y + 3z = 1} & \underline{-x + 3y - 5z = 11} & -3z = 3 & y = 2 & x = 0 \\
 -x + 3y - 5z = 11 & y + 2z = 0 & 4y - 4z = 12 & z = -1 & & \\
 & y = -2z & y - z = 3 & & &
 \end{array}$$

PTS: 4 REF: 061733aai NAT: A.REI.C.6 TOP: Solving Linear Systems
KEY: three variables

758 ANS: 2

$$\left(m^{\frac{5}{3}} \right)^{-\frac{1}{2}} = m^{-\frac{5}{6}} = \frac{1}{\sqrt[6]{m^5}}$$

PTS: 2 REF: 011707aai NAT: N.RN.A.2 TOP: Radicals and Rational Exponents
KEY: variables

759 ANS: 1

$$\begin{array}{l}
 (x+3)^2 + (2x-4)^2 = 8 \quad b^2 - 4ac \\
 x^2 + 6x + 9 + 4x^2 - 16x + 16 = 8 \quad 100 - 4(5)(17) < 0 \\
 5x^2 - 10x + 17 = 0
 \end{array}$$

PTS: 2 REF: 081719aai NAT: A.REI.C.7 TOP: Quadratic-Linear Systems

760 ANS: 2

$$ME = \left(z \sqrt{\frac{p(1-p)}{n}} \right) = \left(1.96 \sqrt{\frac{(0.55)(0.45)}{900}} \right) \approx 0.03 \text{ or } \frac{1}{\sqrt{900}} \approx 0.03$$

PTS: 2

REF: 081612aai

NAT: S.IC.B.4

TOP: Analysis of Data

761 ANS:

$$\text{normcdf}(510, 540, 480, 24) = 0.0994 \quad z = \frac{510 - 480}{24} = 1.25 \quad 1.25 = \frac{x - 510}{20} \quad 2.5 = \frac{x - 510}{20} \quad 535 - 560$$

$$z = \frac{540 - 480}{24} = 2.5 \quad x = 535 \quad x = 560$$

PTS: 4

REF: fall1516aai

NAT: S.ID.A.4

TOP: Normal Distributions

KEY: probability

762 ANS: 3

$f(x) = -f(x)$, so $f(x)$ is odd. $g(-x) \neq g(x)$, so $g(x)$ is not even. $g(-x) \neq -g(x)$, so $g(x)$ is not odd. $h(-x) = h(x)$, so $h(x)$ is even.

PTS: 2

REF: fall1502aai

NAT: F.BF.B.3

TOP: Even and Odd Functions

763 ANS: 2

The 2010 population is 110 million.


PTS: 2

REF: 061718aai

NAT: F.LE.B.5

TOP: Modeling Exponential Functions

764 ANS: 2



PTS: 2

REF: 081603aai

NAT: A.REI.D.11

TOP: Other Systems

KEY: exponential

765 ANS: 4

$$\text{period} = \frac{2\pi}{B}$$

$$\frac{1}{60} = \frac{2\pi}{B}$$

$$B = 120\pi$$

PTS: 2

REF: 061624aai

NAT: F.TF.B.5

TOP: Modeling Trigonometric Functions

766 ANS: 2

$h(x)$ does not have a y-intercept.

PTS: 2

REF: 011719aai

NAT: F.IF.C.9

TOP: Comparing Functions

767 ANS: 4

The vertex is (1,0) and $p = 2$. $y = \frac{1}{4(2)}(x-1)^2 + 0$

PTS: 2 REF: 061717aai NAT: G.GPE.A.2 TOP: Graphing Quadratic Functions

768 ANS: 1

$$x^2 + 2x - 8 = 0$$

$$(x+4)(x-2) = 0$$

$$x = -4, 2$$

PTS: 2 REF: 081701aai NAT: A.APR.D.6 TOP: Undefined Rationals

769 ANS: 2

PTS: 2

REF: 061724aai

NAT: F.BF.B.7

TOP: Series

KEY: geometric

770 ANS: 3

PTS: 2

REF: 081705aai

NAT: F.IF.B.4

TOP: Graphing Trigonometric Functions KEY: increasing/decreasing

771 ANS: 3

$$x^2 + 2x + 1 = -5 + 1$$

$$(x+1)^2 = -4$$

$$x+1 = \pm 2i$$

$$x = -1 \pm 2i$$

PTS: 2 REF: 081703aai NAT: A.REI.B.4 TOP: Solving Quadratics

KEY: complex solutions | completing the square

772 ANS: 2

$$x = \frac{y+1}{y-2}$$

$$xy - 2x = y + 1$$

$$xy - y = 2x + 1$$

$$y(x-1) = 2x + 1$$

$$y = \frac{2x+1}{x-1}$$

PTS: 2 REF: 081714aai NAT: F.BF.B.4 TOP: Inverse of Functions

KEY: rational

773 ANS: 3

$$d = 10 \log \frac{6.3 \times 10^{-3}}{1.0 \times 10^{-12}} \approx 98$$

PTS: 2 REF: 011715aai NAT: F.IF.B.4 TOP: Evaluating Logarithmic Expressions

774 ANS: 3

$$0.75^{\frac{1}{10}} \approx .9716$$


PTS: 2

REF: 061713aai

NAT: A.SSE.B.3

TOP: Modeling Exponential Functions

775 ANS: 2



PTS: 2

REF: 061705aai

NAT: A.REI.D.11

TOP: Other Systems

KEY: polynomial

776 ANS: 2

$$(2 - yi)(2 - yi) = 4 - 4yi + y^2 i^2 = -y^2 - 4yi + 4$$

PTS: 2

REF: 061603aai

NAT: N.CN.A.2

TOP: Operations with Complex Numbers

777 ANS: 1



PTS: 2

REF: 081711aai

NAT: S.ID.A.4

TOP: Normal Distributions

KEY: percent

778 ANS: 3

$$\frac{1}{J} = \frac{1}{F} - \frac{1}{W}$$

$$\frac{1}{J} = \frac{W - F}{FW}$$

$$J = \frac{FW}{W - F}$$

PTS: 2

REF: 081617aai

NAT: A.REI.A.2

TOP: Solving Rationals

KEY: rational solutions

779 ANS: 2

PTS: 2

REF: 081610aai

NAT: F.IF.B.4

TOP: Graphing Trigonometric Functions KEY: increasing/decreasing

780 ANS:

$$\left(\ln \frac{1}{2} \right)$$

$$0 < e^{\frac{1}{1590}} < 1, \text{ so } M(t) \text{ represents decay.}$$

PTS: 2

REF: 011728aai

NAT: F.IF.C.7

TOP: Graphing Exponential Functions

781 ANS: 3

PTS: 2

REF: 061602aai

NAT: A.CED.A.1

TOP: Modeling Rationals

782 ANS: 3

$$-2 \left(-\frac{1}{2}x^2 = -6x + 20 \right)$$

$$x^2 - 12x = -40$$

$$x^2 - 12x + 36 = -40 + 36$$

$$(x - 6)^2 = -4$$

$$x - 6 = \pm 2i$$

$$x = 6 \pm 2i$$

PTS: 2


REF: fall1504aai

NAT: A.REI.B.4

TOP: Solving Quadratics

KEY: complex solutions | completing the square

783 ANS:



PTS: 2

REF: fall1510aai

NAT: A.REI.D.11

TOP: Other Systems

KEY: polynomial

784 ANS: 3

PTS: 2

REF: 081618aai

NAT: F.BF.A.2

TOP: Sequences

KEY: recursive

785 ANS: 3

$$\log_{0.8} \left(\frac{V}{17000} \right) = t \quad \frac{17,000(0.8)^3 - 17,000(0.8)^1}{3 - 1} \approx -2450$$

$$0.8^t = \frac{V}{17000}$$

$$V = 17000(0.8)^t$$

PTS: 2 REF: 081709aia NAT: F.IF.B.6 TOP: Rate of Change

KEY: logarithmic

786 ANS: 2 PTS: 2 REF: 081717aia NAT: S.IC.B.3

TOP: Analysis of Data


787 ANS: 2

$$B(t) = 750 \left(1.16^{\frac{1}{12}} \right)^{12t} \approx 750(1.012)^{12t} \quad B(t) = 750 \left(1 + \frac{0.16}{12} \right)^{12t} \text{ is wrong, because the growth is an annual rate}$$

that is not compounded monthly.

PTS: 2 REF: spr1504aia NAT: A.SSE.B.3 TOP: Modeling Exponential Functions

788 ANS:



This scenario can be modeled with a Venn Diagram:

Since $P(S \cup I) = 0.2$, $P(S \cup I) = 0.8$. Then, $P(S \cap I) = P(S) + P(I) - P(S \cup I)$ If S and I are independent, then the

$$= 0.5 + 0.7 - 0.8$$

$$= 0.4$$

Product Rule must be satisfied. However, $(0.5)(0.7) \neq 0.4$. Therefore, salary and insurance have not been treated independently.

PTS: 4 REF: spr1513aia NAT: S.CP.B.7 TOP: Addition Rule

789 ANS:


$$a_n = x^{n-1}(x+1) \quad x^{n-1} = 0 \quad x+1 = 0$$

$$x = 0 \quad x = -1$$

PTS: 4 REF: spr1511aia NAT: F.BF.A.2 TOP: Sequences

KEY: recursive

790 ANS:



$$\sqrt{x-5} = -x+7 \quad \sqrt{x-5} = -9+7 = -2 \text{ is extraneous.}$$

$$x-5 = x^2 - 14x + 49$$

$$0 = x^2 - 15x + 54$$

$$0 = (x-6)(x-9)$$

$$x = 6, 9$$

PTS: 2 REF: spr1508aai NAT: A.REI.A.2 TOP: Solving Radicals

KEY: extraneous solutions

791 ANS: 1

The graph of $y = \sin x$ is unchanged when rotated 180° about the origin.

PTS: 2 REF: 081614aai NAT: F.BF.B.3 TOP: Even and Odd Functions

792 ANS: 3

$$\sqrt{56-x} = x \quad -8 \text{ is extraneous.}$$

$$56-x = x^2$$

$$0 = x^2 + x - 56$$

$$0 = (x+8)(x-7)$$

$$x = 7$$

PTS: 2 REF: 061605aai NAT: A.REI.A.2 TOP: Solving Radicals

KEY: extraneous solutions


793 ANS:

$$2x^3 - 10x^2 + 11x - 7 = 2x^3 + hx^2 + 3x - 8x^2 - 4hx - 12 + k \quad h = -2$$

$$-2x^2 + 8x + 5 = hx^2 - 4hx + k \quad k = 5$$

PTS: 4 REF: 011733aai NAT: A.APR.C.4 TOP: Polynomial Identities

794 ANS: 4



PTS: 2 REF: 061622aai NAT: A.REI.D.11 TOP: Other Systems
KEY: polynomial

795 ANS: 1 PTS: 2 REF: 061701aai NAT: A.APR.B.3
TOP: Graphing Polynomial Functions

796 ANS: 3

$$(m-2)^2(m+3) = (m^2 - 4m + 4)(m+3) = m^3 + 3m^2 - 4m^2 - 12m + 4m + 12 = m^3 - m^2 - 8m + 12$$

PTS: 2 REF: 081605aai NAT: A.SSE.A.2 TOP: Factoring Polynomials
KEY: factoring by grouping

797 ANS:
 Jillian's plan, because distance increases by one mile each week. $a_1 = 10$ $a_n = n + 12$

$$a_n = a_{n-1} + 1$$

PTS: 4 REF: 011734aai NAT: F.BF.A.2 TOP: Sequences
KEY: recursive

798 ANS:
 Since there are six flavors, each flavor can be assigned a number, 1-6. Use the simulation to see the number of times the same number is rolled 4 times in a row.

PTS: 2 REF: 081728aai NAT: S.IC.A.2 TOP: Analysis of Data

799 ANS: 4

$$4(x^2 - 6x + 9) + 4(y^2 + 18y + 81) = 76 + 36 + 324$$

$$4(x-3)^2 + 4(y+9)^2 = 436$$

PTS: 2 REF: 061619aai NAT: G.GPE.A.1 TOP: Equations of Circles
KEY: completing the square

800 ANS:

$$P(P/K) = \frac{P(P \wedge K)}{P(K)} = \frac{1.9}{2.3} \approx 82.6\%$$
 A key club member has an 82.6% probability of being enrolled in AP Physics.

PTS: 4 REF: 011735aai NAT: S.CP.A.3 TOP: Conditional Probability
801 ANS: 1 PTS: 2 REF: 061708aai NAT: F.TF.B.5
TOP: Modeling Trigonometric Functions

802 ANS: 2 PTS: 2 REF: 061620aai NAT: F.IF.B.4
TOP: Graphing Polynomial Functions

803 ANS:

Yes. The margin of error from this simulation indicates that 95% of the observations fall within ± 0.12 of the simulated proportion, 0.25. The margin of error can be estimated by multiplying the standard deviation, shown to

be 0.06 in the dotplot, by 2, or applying the estimated standard error formula, $\left(\sqrt{\frac{p(1-p)}{n}} \right)$ or $\left(\sqrt{\frac{(0.25)(0.75)}{50}} \right)$

and multiplying by 2. The interval 0.25 ± 0.12 includes plausible values for the true proportion of people who prefer Stephen's new product. The company has evidence that the population proportion could be at least 25%. As seen in the dotplot, it can be expected to obtain a sample proportion of 0.18 (9 out of 50) or less several times, even when the population proportion is 0.25, due to sampling variability. Given this information, the results of the survey do not provide enough evidence to suggest that the true proportion is not at least 0.25, so the development of the product should continue at this time.

PTS: 4 REF: spr1512aai NAT: S.IC.A.2 TOP: Analysis of Data

804 ANS:

Amplitude, because the height of the graph shows the volume of the air.

PTS: 2 REF: 081625aai NAT: F.IF.C.7 TOP: Graphing Trigonometric Functions
KEY: mixed

805 ANS: 3

Since $x + 4$ is a factor of $p(x)$, there is no remainder.

PTS: 2 REF: 081621aai NAT: A.APR.B.2 TOP: Remainder and Factor Theorems

806 ANS: 2

Combining (1) and (3): $-6c = -18$ Combining (1) and (2): $5a + 3c = -1$ Using (3): $-(-2) - 5b - 5(3) = 2$

$$c = 3$$

$$5a + 3(3) = -1$$

$$2 - 5b - 15 = 2$$

$$5a = -10$$

$$b = -3$$

$$a = -2$$

PTS: 2 REF: 081623aai NAT: A.REI.C.6 TOP: Solving Linear Systems
KEY: three variables

807 ANS: 3

$$\left(\frac{1}{2} \right)^{\frac{1}{73.83}} \approx 0.990656$$

PTS: 2 REF: 081710aai NAT: A.SSE.B.3 TOP: Modeling Exponential Functions

808 ANS: 1

$$(1) \frac{9-0}{2-1} = 9 \quad (2) \frac{17-0}{3.5-1} = 6.8 \quad (3) \frac{0-0}{5-1} = 0 \quad (4) \frac{17-5}{3.5-1} \approx 6.3$$

PTS: 2 REF: 011724aai NAT: F.IF.B.6 TOP: Rate of Change
KEY: graph