


F.BF.B.3: Transformations with Functions 4


- 1 The graph below represents $f(x)$.



Which of the following is the graph of $-f(x)$?

- 1)
- 2)
- 3)
- 4)


- 2 The accompanying graph represents the equation $y = f(x)$.



Which graph represents $g(x)$ if $g(x) = -f(x)$?


- 1)
- 2)
- 3)
- 4)

- 3 The graph below represents $f(x)$.




Which graph best represents $f(-x)$?


1)




2)



3)




4)




- 4 Given the parent function $f(x) = x^3$, the function $g(x) = (x - 1)^3 - 2$ is the result of a shift of $f(x)$

- 1) 1 unit left and 2 units down
- 2) 1 unit left and 2 units up
- 3) 1 unit right and 2 units down
- 4) 1 unit right and 2 units up

- 5 Consider the function $y = h(x)$, defined by the graph below.




Which equation could be used to represent the graph shown below?



- 1) $y = h(x) - 2$
- 2) $y = h(x - 2)$
- 3) $y = -h(x)$
- 4) $y = h(-x)$


- 6 The graph below represents national and New York State average gas prices.



If New York State's gas prices are modeled by $G(x)$ and $C > 0$, which expression best approximates the national average x months from August 2014?

- 1) $G(x + C)$
- 2) $G(x) + C$
- 3) $G(x - C)$
- 4) $G(x) - C$

- 7 The height of a running trail is modeled by the quartic function $y = f(x)$ shown below, where x is the distance in miles from the start of the trail and y is the height in feet relative to sea level.




If this trail has a minimum height of 16 feet below sea level, which function(s) could represent a running trail whose minimum height is half of the minimum height of the original trail?

I. $y = f\left(\frac{1}{2}x\right)$ II. $y = f(x) + 8$ III. $y = \frac{1}{2}f(x)$


- 1) I, only
- 2) II, only
- 3) I and III
- 4) II and III

- 8 Richard is asked to transform the graph of $b(x)$ below.




The graph of $b(x)$ is transformed using the equation $h(x) = b(x - 2) - 3$. Describe how the graph of $b(x)$ changed to form the graph of $h(x)$.

- 9 In the diagram below, $f(x) = x^3 + 2x^2$ is graphed. Also graphed is $g(x)$, the result of a translation of $f(x)$.



Determine an equation of $g(x)$. Explain your reasoning.

- 10 The graph of $y = f(x)$ is shown below. The function has a leading coefficient of 1.



Write an equation for $f(x)$. The function g is formed by translating function f left 2 units. Write an equation for $g(x)$.

F.BF.B.3: Transformations with Functions 4**Answer Section**

1 ANS: 3 REF: fall9903b

2 ANS: 1 REF: 060701b

3 ANS: 4 REF: 080406b

4 ANS: 3 REF: 011910ai

5 ANS: 3 REF: 062205aii

6 ANS: 4 REF: 081817aii

7 ANS: 4

I. Minimum does not change, only period; II. $-16 + 8 = -8$; III. $\frac{1}{2}(-16) = -8$

REF: 012523aii

8 ANS:

2 units right and 3 units down.

REF: 081626ai

9 ANS:

 $g(x) = x^3 + 2x^2 - 4$, because $g(x)$ is a translation down 4 units.

REF: 061632ai

10 ANS:

 $f(x) = x^2(x + 4)(x - 3)$; $g(x) = (x + 2)^2(x + 6)(x - 1)$

REF: 011836aii